

Masataka Seike

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5576677/publications.pdf>

Version: 2024-02-01

26
papers

776
citations

687363

13
h-index

552781

26
g-index

26
all docs

26
docs citations

26
times ranked

1218
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Tocopherol suppresses hepatic steatosis by increasing CPT1 expression in a mouse model of diet-induced nonalcoholic fatty liver disease.</i> <i>Obesity Science and Practice</i> , 2021, 7, 91-99.	1.9	8
2	Maximum Plasma Concentration of Lenvatinib Is Useful for Predicting Thrombocytopenia in Patients Treated for Hepatocellular Carcinoma. <i>World Journal of Oncology</i> , 2021, 12, 165-172.	1.5	2
3	PNPLA3 and HLA-DQB1 polymorphisms are associated with hepatocellular carcinoma after hepatitis C virus eradication. <i>Journal of Gastroenterology</i> , 2020, 55, 1162-1170.	5.1	9
4	Supplementation of branched-chain amino acids decreases fat accumulation in the liver through intestinal microbiota-mediated production of acetic acid. <i>Scientific Reports</i> , 2020, 10, 18768.	3.3	31
5	A randomized controlled trial of lusutrombopag in Japanese patients with chronic liver disease undergoing radiofrequency ablation. <i>Journal of Gastroenterology</i> , 2019, 54, 171-181.	5.1	67
6	POGLUT1, the putative effector gene driven by rs2293370 in primary biliary cholangitis susceptibility locus chromosome 3q13.33. <i>Scientific Reports</i> , 2019, 9, 102.	3.3	23
7	Clinical features of hepatocellular carcinoma in nonalcoholic fatty liver disease patients without advanced fibrosis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2019, 34, 1626-1632.	2.8	19
8	Analysis of amino acid profiles of blood over time and biomarkers associated with non-alcoholic steatohepatitis in STAM mice. <i>Experimental Animals</i> , 2019, 68, 417-428.	1.1	8
9	Predictors of hepatocellular carcinoma recurrence associated with the use of direct-acting antiviral agent therapy for hepatitis C virus after curative treatment: A prospective multicenter cohort study. <i>Cancer Medicine</i> , 2019, 8, 2646-2653.	2.8	27
10	Lusutrombopag Reduces Need for Platelet Transfusion in Patients With Thrombocytopenia Undergoing Invasive Procedures. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 1192-1200.	4.4	92
11	Direct-acting antiviral agents do not increase the incidence of hepatocellular carcinoma development: a prospective, multicenter study. <i>Hepatology International</i> , 2019, 13, 293-301.	4.2	38
12	A nationwide survey on non-B, non-C hepatocellular carcinoma in Japan: 2011–2015 update. <i>Journal of Gastroenterology</i> , 2019, 54, 367-376.	5.1	156
13	Efficacy and safety of glecaprevir/pibrentasvir in Japanese patients with chronic genotype 1 hepatitis C virus infection with and without cirrhosis. <i>Journal of Gastroenterology</i> , 2018, 53, 557-565.	5.1	103
14	Usefulness of semiquantitative PCR-invaser assay for selecting candidates for daclatasvir plus asunaprevir combination therapy among patients with hepatitis C virus genotype 1b. <i>Hepatology Research</i> , 2018, 48, 255-263.	3.4	2
15	A Data Mining-based Prognostic Algorithm for NAFLD-related Hepatoma Patients: A Nationwide Study by the Japan Study Group of NAFLD. <i>Scientific Reports</i> , 2018, 8, 10434.	3.3	32
16	Genome-wide association studies identify PRKCB as a novel genetic susceptibility locus for primary biliary cholangitis in the Japanese population. <i>Human Molecular Genetics</i> , 2017, 26, ddw406.	2.9	46
17	Reduced phosphorylation of ribosomal protein S6 is associated with sensitivity to MEK inhibition in gastric cancer cells. <i>Cancer Science</i> , 2016, 107, 1919-1928.	3.9	14
18	Hepatocellular carcinoma in Japanese patients with nonalcoholic fatty liver disease and alcoholic liver disease: multicenter survey. <i>Journal of Gastroenterology</i> , 2016, 51, 586-596.	5.1	49

#	ARTICLE	IF	CITATIONS
19	Risk factors for deterioration of long-term liver function after radiofrequency ablation therapy. <i>World Journal of Hepatology</i> , 2016, 8, 597.	2.0	3
20	Effects of branched-chain amino acids and zinc-enriched nutrients on prognosticators in HCV-infected patients: A multicenter randomized controlled trial. <i>Molecular Medicine Reports</i> , 2015, 11, 2159-2166.	2.4	13
21	Focal nodular hyperplasia-like lesion of progressive type in a patient with congenital hepatic fibrosis: a case report. <i>Acta Hepatologica Japonica</i> , 2015, 56, 461-468.	0.1	1
22	Alpha-fetoprotein and des-gamma-carboxy-prothrombin at twenty-four weeks after interferon-based therapy predict hepatocellular carcinoma development. <i>World Journal of Hepatology</i> , 2015, 7, 2757.	2.0	4
23	Benefits of nucleos(t)ide analog treatments for hepatitis B virus-related cirrhosis. <i>World Journal of Hepatology</i> , 2015, 7, 2404.	2.0	10
24	High alanine aminotransferase level as a predictor for the incidence of macrovascular disease in type 2 diabetic patients with fatty liver disease. <i>Hepatology International</i> , 2013, 7, 555-561.	4.2	1
25	Heterozygosity for leptin receptor (<i>fa</i>) accelerates hepatic triglyceride accumulation without hyperphagia in Zucker rats. <i>Obesity Research and Clinical Practice</i> , 2009, 3, 29-34.	1.8	6
26	H2-Blocker modulates heart rate variability. <i>Heart and Vessels</i> , 1999, 14, 137-142.	1.2	12