Sevket Balta

List of Publications by Citations

Source: https://exaly.com/author-pdf/5581894/sevket-balta-publications-by-citations.pdf

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

2,966 46 27 372 h-index g-index citations papers 3,368 402 2.4 5.33 L-index avg, IF ext. citations ext. papers

| # | Paper | IF | Citations |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|-----------|
| 372 | The Relation Between Atherosclerosis and the Neutrophil-Lymphocyte Ratio. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2016 , 22, 405-11 | 3.3 | 176 |
| 371 | Endocan: A novel inflammatory indicator in cardiovascular disease?. <i>Atherosclerosis</i> , 2015 , 243, 339-43 | 3.1 | 123 |
| 370 | The platelet-lymphocyte ratio: A simple, inexpensive and rapid prognostic marker for cardiovascular events. <i>Platelets</i> , 2015 , 26, 680-1 | 3.6 | 116 |
| 369 | Endocana novel inflammatory indicator in newly diagnosed patients with hypertension: a pilot study. <i>Angiology</i> , 2014 , 65, 773-7 | 2.1 | 112 |
| 368 | Serum endocan levels as a marker of disease activity in patients with Behat disease. <i>Journal of the American Academy of Dermatology</i> , 2014 , 70, 291-6 | 4.5 | 107 |
| 367 | Comparative effects of nebivolol and metoprolol on red cell distribution width and neutrophil/lymphocyte ratio in patients with newly diagnosed essential hypertension. <i>Journal of Cardiovascular Pharmacology</i> , 2013 , 62, 388-93 | 3.1 | 90 |
| 366 | The comparative effects of valsartan and amlodipine on vWf levels and N/L ratio in patients with newly diagnosed hypertension. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 516-22 | 2.2 | 80 |
| 365 | Association between coronary artery ectasia and neutrophil-lymphocyte ratio. <i>Angiology</i> , 2013 , 64, 627 | -32 1 | 80 |
| 364 | Neutrophil-lymphocyte ratio and carotid-intima media thickness in patients with Behlet disease without cardiovascular involvement. <i>Angiology</i> , 2015 , 66, 291-6 | 2.1 | 70 |
| 363 | Evaluation of the mean platelet volume in patients with cardiac syndrome X. <i>Clinics</i> , 2012 , 67, 1019-22 | 2.3 | 70 |
| 362 | The platelet lymphocyte ratio may be useful inflammatory indicator in clinical practice. Hemodialysis International, 2013 , 17, 668-9 | 1.7 | 68 |
| 361 | Endocan, a novel marker of endothelial dysfunction in patients with essential hypertension: comparative effects of amlodipine and valsartan. <i>Blood Pressure</i> , 2015 , 24, 55-60 | 1.7 | 59 |
| 360 | Aortic arterial stiffness is a moderate predictor of cardiovascular disease in patients with psoriasis vulgaris. <i>Angiology</i> , 2014 , 65, 74-8 | 2.1 | 58 |
| 359 | Mean platelet volume in recurrent aphthous stomatitis and Behät disease. <i>Angiology</i> , 2014 , 65, 161-5 | 2.1 | 57 |
| 358 | Neutrophils/lymphocytes ratio in patients with cardiac syndrome X and its association with carotid intima-media thickness. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2014 , 20, 250-5 | 3.3 | 56 |
| 357 | Predictive value of admission platelet volume indices for in-hospital major adverse cardiovascular events in acute ST-segment elevation myocardial infarction. <i>Angiology</i> , 2015 , 66, 155-62 | 2.1 | 51 |
| 356 | Endocan Levels and Subclinical Atherosclerosis in Patients With Systemic Lupus Erythematosus. <i>Angiology</i> , 2016 , 67, 749-55 | 2.1 | 48 |

(2013-2013)

| 355 | Higher neutrophil to lymhocyte ratio in patients with metabolic syndrome. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2013 , 19, 579 | 3.3 | 44 |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 354 | Red cell distribution width: a novel inflammatory marker in clinical practice. <i>Cardiology Journal</i> , 2013 , 20, 209 | 1.4 | 38 |
| 353 | Neutrophil-lymphocyte ratio as an important assessment tool. <i>Expert Review of Cardiovascular Therapy</i> , 2014 , 12, 537-8 | 2.5 | 37 |
| 352 | Red cell distribution width is a predictor of mortality in patients with severe sepsis and septic shock. <i>American Journal of Emergency Medicine</i> , 2013 , 31, 989-90 | 2.9 | 35 |
| 351 | Red cell distribution width is a predictor of mortality in patients undergoing coronary artery bypass surgery. <i>European Journal of Cardio-thoracic Surgery</i> , 2013 , 44, 396-7 | 3 | 34 |
| 350 | Predictive value of admission red cell distribution width-platelet ratio for no-reflow phenomenon in acute ST segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. <i>Cardiology Journal</i> , 2016 , 23, 84-92 | 1.4 | 31 |
| 349 | Assessment of the relationship between red cell distribution width and cardiac syndrome X. <i>Kardiologia Polska</i> , 2013 , 71, 480-4 | 0.9 | 31 |
| 348 | Predictors of No-Reflow Phenomenon in Young Patients With Acute ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. <i>Angiology</i> , 2016 , 67, 683-9 | 2.1 | 29 |
| 347 | Carotid intima-media thickness in patients with slow coronary flow and its association with neutrophil-to-lymphocyte ratio: a preliminary report. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2014 , 20, 393-9 | 3.3 | 29 |
| 346 | Bilirubin levels and their association with carotid intima media thickness and high-sensitivity C-reactive protein in patients with psoriasis vulgaris. <i>American Journal of Clinical Dermatology</i> , 2014 , 15, 137-42 | 7.1 | 27 |
| 345 | The relation between monocyte to HDL ratio and no-reflow phenomenon in the patients with acute ST-segment elevation myocardial infarction. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 1542-7 | 2.9 | 26 |
| 344 | The Relation Between No-Reflow Phenomenon and Complete Blood Count Parameters. <i>Angiology</i> , 2017 , 68, 381-388 | 2.1 | 25 |
| 343 | The Neutrophil-Lymphocyte Ratio and Inflammation. <i>Angiology</i> , 2016 , 67, 298-9 | 2.1 | 24 |
| 342 | The relationship between some of the cardiovascular risk factors and arterial stiffness parameters in essentially hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 444-8 | 2.2 | 23 |
| 341 | Novel myokine: irisin may be an independent predictor for subclinic atherosclerosis in Beheltß disease. <i>Journal of Investigative Medicine</i> , 2016 , 64, 875-81 | 2.9 | 23 |
| 340 | The comparative effects of valsartan and amlodipine on vascular microinflammation in newly diagnosed hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 418-23 | 2.2 | 22 |
| 339 | Neutrophil-lymphocyte ratio may predict left atrial thrombus in patients with nonvalvular atrial fibrillation. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015 , 21, 166-71 | 3.3 | 21 |
| 338 | Other inflammatory markers should not be forgetten when assessing the neutrophil-to-lymphocyte ratio. Clinical and Applied Thrombosis/Hemostasis, 2013, 19, 693-4 | 3.3 | 20 |

| 337 | Arterial stiffness itself without other inflammatory markers may not provide information to clinicians. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 303 | 2.3 | 20 |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|----|
| 336 | Carotid Intima Media Thickness and Its Association With Total Bilirubin Levels in Patients With Coronary Artery Ectasia. <i>Angiology</i> , 2020 , 71, 425-430 | 2.1 | 19 |
| 335 | Red cell distribution width as a novel, simple, inexpensive predictor of mortality in patients with chronic heart failure. <i>International Journal of Cardiology</i> , 2013 , 168, 3049-50 | 3.2 | 18 |
| 334 | Endocan and atherosclerosis. <i>Angiology</i> , 2015 , 66, 490 | 2.1 | 17 |
| 333 | Ivabradine improves heart rate variability in patients with nonischemic dilated cardiomyopathy. <i>Arquivos Brasileiros De Cardiologia</i> , 2014 , 103, 308-14 | 1.2 | 16 |
| 332 | Red cell distribution width: a novel and simple predictor of mortality in acute pancreatitis. <i>American Journal of Emergency Medicine</i> , 2013 , 31, 991-2 | 2.9 | 16 |
| 331 | The lymphocyte-monocyte ratio in clinical practice. <i>Journal of Clinical Pathology</i> , 2016 , 69, 88-9 | 3.9 | 15 |
| 330 | The impact of mitral stenosis on left atrial function assessed by two-dimensional speckle tracking echocardiography. <i>Echocardiography</i> , 2012 , 29, 1064-70 | 1.5 | 15 |
| 329 | Asymmetric dimethylarginine and augmentation index in newly diagnosed patients with hypertension. <i>Angiology</i> , 2015 , 66, 43-8 | 2.1 | 14 |
| 328 | Neutrophil-lymphocyte ratio as an useful mortality marker. <i>American Journal of Emergency Medicine</i> , 2014 , 32, 1546-7 | 2.9 | 14 |
| 327 | Behlet disease and risk of vascular events. Current Opinion in Cardiology, 2016, 31, 451-7 | 2.1 | 14 |
| 326 | Ischemia-Modified Albumin and Atherosclerosis in Patients With Familial Mediterranean Fever. <i>Angiology</i> , 2016 , 67, 456-60 | 2.1 | 13 |
| 325 | The Relationship Between Atherogenic Index and Carotid Artery Atherosclerosis in Familial Mediterranean Fever. <i>Angiology</i> , 2017 , 68, 315-321 | 2.1 | 13 |
| 324 | Neutrophil-to-Lymphocyte Ratio in Prognosis of Gastric Cancer. <i>Journal of Gastric Cancer</i> , 2013 , 13, 196 | -73.2 | 13 |
| 323 | Other inflammatory markers ought to be kept in mind when assessing the mean platelet volume in clinical practice. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013 , 270, 2373-4 | 3.5 | 12 |
| 322 | Carotid intima media thickness can predict coronary artery disease. <i>International Journal of Cardiology</i> , 2015 , 201, 331 | 3.2 | 12 |
| 321 | Hemostatic markers can be pivotal roles of risk factors for new-onset atrial fibrillation. <i>Platelets</i> , 2014 , 25, 554-5 | 3.6 | 12 |
| 320 | The relation between ischemia modified albumin levels and carotid intima media thickness in patients with rheumatoid arthritis. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 32-37 | 2.3 | 12 |

(2014-2014)

| 319 | Comparision of effects of rosuvastatin versus atorvastatin treatment on plasma levels of asymmetric dimethylarginine in patients with hyperlipidemia having coronary artery disease. <i>Angiology</i> , 2014 , 65, 788-93 | 2.1 | 11 |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----|
| 318 | The neutrophil lymphocyte ratio in coronary heart disease. <i>International Journal of Cardiology</i> , 2014 , 176, 267 | 3.2 | 11 |
| 317 | A new anomaly of the left anterior descending artery: Type X dual LAD. <i>Indian Heart Journal</i> , 2015 , 67 Suppl 3, S14-7 | 1.6 | 11 |
| 316 | Arterial stiffness and endothelial inflammation in prediabetes and newly diagnosed diabetes patients. <i>Archives of Endocrinology and Metabolism</i> , 2015 , 59, 407-13 | 2.2 | 11 |
| 315 | Endothelial Dysfunction and Inflammatory Markers of Vascular Disease. <i>Current Vascular Pharmacology</i> , 2021 , 19, 243-249 | 3.3 | 11 |
| 314 | Whole Blood Viscosity and Cardiovascular Diseases: A Forgotten Old Player of the Game. <i>Medical Principles and Practice</i> , 2016 , 25, 499-500 | 2.1 | 11 |
| 313 | The relation between lymphocyte-monocyte ratio and renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 421 | 2.8 | 10 |
| 312 | Epicardial adipose tissue should be evaluated with other inflammatory markers in patients with subclinical hypothyroidism. <i>Medical Principles and Practice</i> , 2013 , 22, 603 | 2.1 | 10 |
| 311 | The Relation Between Endocan Levels and Subclinic Atherosclerosis. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2016 , 22, 495-6 | 3.3 | 10 |
| 310 | Relationship of Systolic Blood Pressure and Body Mass Index With Left Ventricular Mass and Mass Index in Adolescents. <i>Angiology</i> , 2016 , 67, 58-65 | 2.1 | 9 |
| 309 | Renal dysfunction may predict new onset heart failure. American Heart Journal, 2013, 166, e5 | 4.9 | 9 |
| 308 | Red cell distribution width: a novel and simple predictor of mortality in chronic obstructive pulmonary disease. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2014 , 11, 475-6 | 2 | 9 |
| 307 | Nonalcoholic Fatty liver disease may be associated with coronary artery disease complexity. <i>Angiology</i> , 2013 , 64, 639-40 | 2.1 | 9 |
| 306 | Epicardial adipose tissue measurement: inexpensive, easy accessible and rapid practical method. <i>Anatolian Journal of Cardiology</i> , 2013 , 13, 611 | | 9 |
| 305 | Neutrophil to lymphocyte ratio may predict mortality in breast cancer patients. <i>Journal of Breast Cancer</i> , 2013 , 16, 354-5 | 3 | 9 |
| 304 | The association between red cell distribution width and non-small-cell lung cancer. <i>European Journal of Cardio-thoracic Surgery</i> , 2014 , 45, 954 | 3 | 8 |
| 303 | Serum gamma-glutamyltransferase (GGT) should be evaluated together with other inflammatory markers in clinical practice. <i>Angiology</i> , 2013 , 64, 401 | 2.1 | 8 |
| 302 | Epicardial fat thickness in psoriasis patients. <i>Dermatology</i> , 2014 , 228, 132-3 | 4.4 | 8 |

| 301 | The neutrophil-lymphocyte ratio is not enough to describe inflammatory condition. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 1839-40 | 3.5 | 8 |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 300 | Red cell distribution width should be assessed together with other inflammatory markers in daily clinical practice. <i>Cardiology</i> , 2013 , 124, 60 | 1.6 | 8 |
| 299 | Subclinical peripheral arterial disease and ankle-brachial index. <i>Angiology</i> , 2013 , 64, 395-6 | 2.1 | 8 |
| 298 | Mobile mass lesion in the aorta after transcatheter aortic valve implantation: Thrombus or residue calcification. <i>International Journal of Cardiology</i> , 2015 , 198, 45-6 | 3.2 | 7 |
| 297 | Carotid intima media thickness and subclinical early atherosclerosis. <i>International Journal of Cardiology</i> , 2016 , 203, 1146 | 3.2 | 7 |
| 296 | Other inflammatory indicators should be kept in mind when assessing red cell distribution width in patients with pneumonia. <i>American Journal of Emergency Medicine</i> , 2013 , 31, 1144-5 | 2.9 | 7 |
| 295 | Mean platelet volume as a surrogate marker of long-term mortality in patients undergoing percutaneous coronary intervention. <i>American Journal of Cardiology</i> , 2013 , 112, 142 | 3 | 7 |
| 294 | Letter to the editor: predictive value of the neutrophil-lymphocyte ratio and mean platelet volume in testicular torsion. <i>Korean Journal of Urology</i> , 2015 , 56, 601-2 | | 7 |
| 293 | The relationship between red cell distribution width and homozygous M694V mutation in familial Mediterranean fever patients. <i>Annals of Saudi Medicine</i> , 2015 , 35, 151-6 | 1.6 | 7 |
| 292 | The Relationship Among the Level of Serum Amyloid A, High-Density Lipoprotein and Microalbuminuria in Patients With Familial Mediterranean Fever. <i>Journal of Clinical Laboratory Analysis</i> , 2016 , 30, 1003-1008 | 3 | 7 |
| 291 | Assessment of Platelet-Lymphocyte Ratio Based on EDTA-Dependent Pseudothrombocytopenia. <i>Angiology</i> , 2016 , 67, 96-7 | 2.1 | 6 |
| 290 | The neutrophil-lymphocyte ratio and atherosclerosis. <i>Expert Review of Cardiovascular Therapy</i> , 2016 , 14, 885 | 2.5 | 6 |
| 289 | The progressed atrioventricular block associated with ticagrelor therapy may not require permanent pacemaker after acute coronary syndrome; it may be reversible. <i>International Journal of Cardiology</i> , 2016 , 203, 822-4 | 3.2 | 6 |
| 288 | Ischemia-modified albumin in patients with seizure. <i>American Journal of Emergency Medicine</i> , 2014 , 32, 1282 | 2.9 | 6 |
| 287 | Phase analysis in patients with reversible perfusion defects and normal coronary arteries at angiography. <i>Annals of Nuclear Medicine</i> , 2013 , 27, 416-22 | 2.5 | 6 |
| 286 | Higher N-terminal pro-B-type natriuretic peptide may be related to very different conditions. <i>Journal of the American College of Cardiology</i> , 2013 , 62, 1634-5 | 15.1 | 6 |
| 285 | The pivotal roles of risk factors for incident atrial fibrillation: interweaving pieces of puzzle. <i>International Journal of Cardiology</i> , 2013 , 168, 2937-8 | 3.2 | 6 |
| 284 | Mean platelet volume as a surrogate marker of low-grade inflammation in osteoarthritis. <i>Platelets</i> , 2014 , 25, 643-4 | 3.6 | 6 |

| 283 | Other inflammatory markers should be kept in mind when assessing the mean platelet volume. <i>Platelets</i> , 2014 , 25, 552-3 | 3.6 | 6 |
|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|---|
| 282 | Neutrophil-to-lymphocyte ratio may predict contrast-induced nephropathy. <i>Angiology</i> , 2014 , 65, 57-8 | 2.1 | 6 |
| 281 | The relation between decreased glomerular filtration rate and nonvalvular atrial fibrillation. <i>Cardiology</i> , 2013 , 124, 219 | 1.6 | 6 |
| 280 | Epicardial fat thickness and cardio-ankle vascular index without other inflammatory markers may not provide information to clinicians about the systemic inflammation. <i>Cardiology</i> , 2013 , 125, 13-4 | 1.6 | 6 |
| 279 | The relationship between ankle-brachial index and estimated glomerular filtration rate in type 2 diabetes. <i>Angiology</i> , 2013 , 64, 242 | 2.1 | 6 |
| 278 | Peripheral arterial disease assessment with photoplethysmography and continuous-wave Doppler ultrasound in addition to ankle-brachial index may loss time and funds. <i>Angiology</i> , 2013 , 64, 321 | 2.1 | 6 |
| 277 | Only ankle-brachial index may not be an accurate information about the prevalence of peripheral arterial disease. <i>Angiology</i> , 2013 , 64, 481-2 | 2.1 | 6 |
| 276 | Serum uric acid still carries controversies about its role in endothelial dysfunction. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 296 | 2.3 | 6 |
| 275 | The Relationship Between Plasma Whole Blood Viscosity and Cardiovascular Events in Patients With Chronic Kidney Disease. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017 , 23, 663-670 | 3.3 | 5 |
| 274 | Red cell distribution width is related to stroke in patients with heart failure. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015 , 21, 190 | 3.3 | 5 |
| 273 | The Relation Between Neutrophil-Lymphocyte Ratio and Hypertension. <i>American Journal of Hypertension</i> , 2015 , 28, 1386 | 2.3 | 5 |
| 272 | Thrombotic events in Behetß disease. American Journal of Emergency Medicine, 2014, 32, 184-5 | 2.9 | 5 |
| 271 | Amlodipine increases vitamin D levels more than valsartan in newly diagnosed hypertensive patients: pointing to an additional effect on bone metabolism or a novel marker of inflammation?. <i>Renal Failure</i> , 2013 , 35, 691-6 | 2.9 | 5 |
| 270 | Red Cell Distribution Width in Myocardial Infarction. <i>Medical Principles and Practice</i> , 2015 , 24, 584 | 2.1 | 5 |
| 269 | Computed tomography guided percutaneous transapical closure of cardiac apex after prosthetic mitral paravalvular leak closure. <i>International Journal of Cardiology</i> , 2015 , 198, 37-9 | 3.2 | 5 |
| 268 | Peripheral arterial disease in patients with Behetß disease. Rheumatology International, 2014 , 34, 589-9 | 90 3.6 | 5 |
| 267 | Epicardial fat thickness should be evaluated with other inflammatory markers and cardiovascular risk factors. <i>Echocardiography</i> , 2013 , 30, 739 | 1.5 | 5 |
| 266 | Cell-free circulating DNA as a novel biomarker in patients with the acute coronary syndrome. <i>Cardiology</i> , 2013 , 126, 122-3 | 1.6 | 5 |

| 265 | The red cell distribution width may be affected by many factors in the clinical practice. <i>Journal of Clinical and Diagnostic Research JCDR</i> , 2013 , 7, 1830 | Ο | 5 |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 264 | Relationship Between Neck Circumference and Epicardial Fat Thickness in a Healthy Male Population. <i>Arquivos Brasileiros De Cardiologia</i> , 2016 , 107, 266-270 | 1.2 | 5 |
| 263 | Waist-Hip Ratio in Patients with Acute Myocardial Infarction. <i>Journal of Clinical and Diagnostic Research JCDR</i> , 2015 , 9, OL01 | О | 5 |
| 262 | The best treatment option in the management of patients with acute massive pulmonary embolism: surgery or thrombolysis?. <i>International Journal of Cardiology</i> , 2016 , 214, 286-7 | 3.2 | 5 |
| 261 | Re: Effect of renin-angiotensin-aldosterone system blockade therapy on incidence of contrast-induced nephropathy in patients with chronic kidney disease. <i>Iranian Journal of Kidney Diseases</i> , 2013 , 7, 166-7 | 0.9 | 5 |
| 260 | The main argument about the etiology of coronary artery ectasia: is it inflammation or not?. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2014 , 42, 119-20 | | 5 |
| 259 | The healing of spontaneous coronary artery dissection with conservative treatment: When to stop. <i>International Journal of Cardiology</i> , 2015 , 189, 249-51 | 3.2 | 4 |
| 258 | Mean platelet volume in patients undergoing percutaneous coronary intervention. <i>Platelets</i> , 2015 , 26, 269-70 | 3.6 | 4 |
| 257 | Significant left hemothorax after transapical closure of cardiac apex with minithoracotomy and transapical transcatheter prosthetic mitral paravalvular leak closure. <i>International Journal of Cardiology</i> , 2015 , 199, 274-6 | 3.2 | 4 |
| 256 | Arterial Stiffness: Good Predictor for Hypertensive Patients. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 596 | 2.3 | 4 |
| 255 | The spontaneous coronary artery dissection may need intervention in the proximal segment of the arteries. <i>International Journal of Cardiology</i> , 2016 , 202, 943-4 | 3.2 | 4 |
| 254 | Red cell distribution width in organophosphate exposure patients. <i>American Journal of Emergency Medicine</i> , 2014 , 32, 1132 | 2.9 | 4 |
| 253 | Neutrophil-Lymphocyte Ratio in Behët Disease. <i>Angiology</i> , 2015 , 66, 695 | 2.1 | 4 |
| 252 | Neutrophil-Lymphocyte Ratio May Predict the Degree of Coronary Collateral Circulation. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015 , 21, 586-7 | 3.3 | 4 |
| 251 | Mean platelet volume as an inflammatory indicator in behcet disease. Angiology, 2014, 65, 167 | 2.1 | 4 |
| 250 | Three-dimensional echocardiographic imaging of biventricular false tendons mimicking hypertrophic cardiomyopathy. <i>Echocardiography</i> , 2012 , 29, E178-9 | 1.5 | 4 |
| 249 | Response to Red blood cell distribution width is a predictor of readmission in cardiac patients? <i>Clinical Cardiology</i> , 2013 , 36, 364-5 | 3.3 | 4 |
| 248 | Red blood cell distribution width is predictive of mortality in trauma patients. <i>Journal of Trauma and Acute Care Surgery</i> , 2013 , 75, 345-6 | 3.3 | 4 |

| 247 | Endocan: a new marker of endothelial function. Current Opinion in Cardiology, 2021, 36, 462-468 | 2.1 | 4 |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|---|
| 246 | Endocan and Erectile Dysfunction. American Journal of Menls Health, 2019, 13, 1557988319893889 | 2.2 | 4 |
| 245 | The relationship between neutrophil-lymphocyte ratio and acute aortic dissection. <i>Perfusion (United Kingdom)</i> , 2017 , 32, 336-337 | 1.9 | 3 |
| 244 | Plasma B-type natriuretic peptide in congenital heart disease. <i>American Journal of Emergency Medicine</i> , 2015 , 33, 1701 | 2.9 | 3 |
| 243 | The combination of percutaneous transapical and transseptal closure of cardiac apex and prosthetic mitral paravalvular leak with loop technique. <i>International Journal of Cardiology</i> , 2016 , 203, 221-4 | 3.2 | 3 |
| 242 | Red blood cell distribution width: Just one of many things to consider. <i>International Journal of Cardiology</i> , 2016 , 203, 438-9 | 3.2 | 3 |
| 241 | Risk factors for new-onset atrial fibrillation. <i>International Journal of Cardiology</i> , 2014 , 171, e46 | 3.2 | 3 |
| 240 | High BNP levels in rheumatoid arthritis may be related with right ventricular functions. <i>International Journal of Cardiology</i> , 2014 , 174, 149-50 | 3.2 | 3 |
| 239 | Inflammatory markers should be assessed together with cardiovascular risk factors by clinicians in masked hypertension. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 443-4 | 2.3 | 3 |
| 238 | Other inflammatory markers and related factors should be kept in mind in metabolic syndrome with psoriasis patients. <i>Archives of Dermatological Research</i> , 2013 , 305, 459-60 | 3.3 | 3 |
| 237 | Atrial fibrillation in patients with acute coronary syndromes. <i>International Journal of Cardiology</i> , 2013 , 168, 5049 | 3.2 | 3 |
| 236 | Red cell distribution width is a predictor of mortality in patients with major bleeding. <i>Revista Portuguesa De Cardiologia (English Edition)</i> , 2013 , 32, 843-844 | О | 3 |
| 235 | Amlodipine seems to be superior to valsartan in decreasing microalbuminuria in newly diagnosed hypertensive patients: a novel effect to be explained with hyperfiltration?. <i>Renal Failure</i> , 2013 , 35, 357- | -6 2 .9 | 3 |
| 234 | Mean platelet volume may indicate early diagnosed gastric cancer based on inflammation. <i>Platelets</i> , 2015 , 26, 99-100 | 3.6 | 3 |
| 233 | Anaemia is a predictor of mortality in patients undergoing aortic valve surgery. <i>European Journal of Cardio-thoracic Surgery</i> , 2014 , 45, 205-6 | 3 | 3 |
| 232 | Vitamin D levels in patients with Behätß disease. <i>International Journal of Rheumatic Diseases</i> , 2014 , 17, 479-80 | 2.3 | 3 |
| 231 | Inflammatory status as a major role of risk factor for atrial fibrillation. <i>Journal of Thrombosis and Thrombolysis</i> , 2014 , 37, 540-1 | 5.1 | 3 |
| 230 | Incremental value of three-dimensional transesophageal echocardiography in a patient with parachute mitral valve. <i>Echocardiography</i> , 2012 , 29, E24-5 | 1.5 | 3 |

| 229 | First report of endocarditis by Alloiococcus otitidis spp. in a patient with a history of chronic otitis. Journal of Infection and Public Health, 2013 , 6, 494-5 | 7.4 | 3 |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|---|
| 228 | Inflammatory markers may predict long-term cardiovascular mortality in patients with acute coronary syndrome. <i>Cardiology</i> , 2013 , 125, 88-9 | 1.6 | 3 |
| 227 | The reasons of higher NT-proBNP depend on very different conditions. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, e17 | 2.4 | 3 |
| 226 | YKL-40 levels in patients with coronary artery ectasia. <i>Anatolian Journal of Cardiology</i> , 2014 , 14, 97-8 | | 3 |
| 225 | Pivotal roles of risk factors for incident atrial fibrillation in patients with newly diagnosed hyperthyroidism. <i>Journal of Geriatric Cardiology</i> , 2013 , 10, 119-20 | 1.7 | 3 |
| 224 | Carotid intima-media thickness and other inflammatory markers in clinical practice. <i>Arquivos Brasileiros De Cardiologia</i> , 2013 , 100, 585 | 1.2 | 3 |
| 223 | The effect of standard therapy on mean platelet volume in patients with chronic hepatitis C. <i>Przeglad Gastroenterologiczny</i> , 2016 , 11, 200-205 | 6 | 3 |
| 222 | Can a novel scoring system derived from hemodynamic and anthropometric variables predict sympathetic drive in young patients?. <i>Blood Pressure Monitoring</i> , 2016 , 21, 21-6 | 1.3 | 3 |
| 221 | Red cell distribution width is associated with albuminuria in adults with familial Mediterranean fever. <i>Kaohsiung Journal of Medical Sciences</i> , 2016 , 32, 216-20 | 2.4 | 3 |
| 220 | The relation between inflammation and coronary artery ectasia. <i>Revista Portuguesa De Cardiologia</i> , 2016 , 35, 553-4 | 1 | 3 |
| 219 | ADMA is a useful marker, but many confounding factors should be considered!. <i>Anatolian Journal of Cardiology</i> , 2015 , 15, 81 | | 2 |
| 218 | Percutaneous mitral annuloplasty in a patient with coronary sinus stenosis and coronary artery compression during procedure; they will not interfere. <i>International Journal of Cardiology</i> , 2015 , 191, 84-6 | 3.2 | 2 |
| 217 | The relation between neutrophil-lymphocyte ratio and acute kidney injury. Renal Failure, 2015, 37, 1527 | ' -8 .9 | 2 |
| 216 | Percutaneous closure of ventricular septal defect with membraneous pouch and changing the device due to severe aortic regurgitation. <i>International Journal of Cardiology</i> , 2016 , 209, 42-5 | 3.2 | 2 |
| 215 | Hypercholesterolemia is Accounted for Atherosclerosis at the Proximal Arterial Segments of Myocardial Bridging: A Pilot Study. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2016 , 22, 297-302 | 3.3 | 2 |
| 214 | Right ventricular function in patients undergoing surgical or transcatheter aortic valve replacement. <i>European Journal of Cardio-thoracic Surgery</i> , 2016 , 49, 1296 | 3 | 2 |
| 213 | The Neutrophil-to-Lymphocyte Ratio Is Valuable at All Stages of Coronary Artery Disease. <i>Cardiology</i> , 2016 , 133, 56 | 1.6 | 2 |
| 212 | Occupational exposure to hand-arm vibration. <i>International Journal of Cardiology</i> , 2016 , 203, 959 | 3.2 | 2 |

| 211 | Is it possible to prevent acute kidney injury in patients who underwent contrast medium?. <i>Angiology</i> , 2014 , 65, 224 | 2.1 | 2 |
|-------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 210 | Neutrophil-to-lymphocyte ratio as a novel independent prognostic factor in urothelial carcinoma. <i>Clinical Genitourinary Cancer</i> , 2014 , 12, e69-70 | 3.3 | 2 |
| 209 | Atrial fibrillation presented with syncope in a jet pilot during daily briefing on squadron. <i>Aviation, Space, and Environmental Medicine</i> , 2014 , 85, 965-9 | | 2 |
| 208 | To maintain a proper follow-up of the patients with coronary artery disease is as nearly important as medications. <i>International Journal of Cardiology</i> , 2013 , 167, 2372 | 3.2 | 2 |
| 207 | Carotid intima-media thickness: a novel inflammatory marker which should not be assessed alone!. <i>Cardiology</i> , 2013 , 124, 207 | 1.6 | 2 |
| 206 | Red cell distribution width without additional cost compared with a relatively expensive test measurement in clinical practice. <i>International Journal of Cardiology</i> , 2013 , 168, 4899-900 | 3.2 | 2 |
| 205 | Red cell distribution width is a predictor of mortality in patients with major bleeding. <i>Revista Portuguesa De Cardiologia</i> , 2013 , 32, 843-4 | 1 | 2 |
| 204 | Levels of vitamin D and its effects on bone metabolism and cardiovascular system should be assessed after isolation of confounding factors. <i>International Journal of Cardiology</i> , 2013 , 168, 628 | 3.2 | 2 |
| 203 | Do we just assess the left ventricle in pregnant women with structural heart disease?. <i>International Journal of Cardiology</i> , 2013 , 168, 591 | 3.2 | 2 |
| 202 | The relation between N-terminal pro-B-type natriuretic peptide and heart failure. <i>American Journal of Emergency Medicine</i> , 2013 , 31, 1533 | 2.9 | 2 |
| 201 | Role of Screening Tests in the Detection and Management of Blood Pressure Abnormalities Among Young Population. <i>Angiology</i> , 2017 , 68, 441-446 | 2.1 | 2 |
| 2 00 | Red cell distribution width in pulmonary embolism. Wiener Klinische Wochenschrift, 2014 , 126, 553-4 | 2.3 | 2 |
| 199 | Non-alcoholic fatty liver disease may be associated with endothelial dysfunction. <i>Upsala Journal of Medical Sciences</i> , 2014 , 119, 57 | 2.8 | 2 |
| 198 | Different imaging modalities in quantification of epicardial adipose tissue thickness. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 616 | 2.3 | 2 |
| 197 | Coronary artery ectasia as a histopathological pattern of atherosclerosis. <i>Angiology</i> , 2014 , 65, 86 | 2.1 | 2 |
| 196 | Red cell distribution width in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2014 , 26, 361 | 2.2 | 2 |
| 195 | Other factors ought to be kept in mind when analyzing plasma asymmetric dimethylarginine levels. <i>American Journal of Hypertension</i> , 2014 , 27, 500 | 2.3 | 2 |
| 194 | CARDIOVASCULAR COMORBIDITIES IN PATIENTS WITH PSORIASIS. <i>Gulhane Medical Journal</i> , 2014 , 56, 61 | | 2 |

| 193 | Mean platelet volume can be affected by many factors and should be assessed together with other inflammatory markers. <i>Platelets</i> , 2014 , 25, 388-9 | 3.6 | 2 |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|---|
| 192 | Parachute tricuspid valve in a patient with atrial septal defect detected by two- and three-dimensional echocardiography. <i>Echocardiography</i> , 2012 , 29, E255-7 | 1.5 | 2 |
| 191 | Arterial stiffness in patients with peripheral arterial disease. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 938 | 2.3 | 2 |
| 190 | The protectiveness of the treatment of vitamin D insufficiency in the development of diabetes. <i>Arquivos Brasileiros De Endocrinologia E Metabologia</i> , 2013 , 57, 157-8 | | 2 |
| 189 | Arterial stiffness should be evaluated with other inflammatory markers in patients with subclinical hypothyroidism. <i>Arquivos Brasileiros De Endocrinologia E Metabologia</i> , 2013 , 57, 754-5 | | 2 |
| 188 | Mean platelet volume as one part of platelet function determining inflammation. <i>Annals of Saudi Medicine</i> , 2016 , 36, 234 | 1.6 | 2 |
| 187 | Helicobacter pylori infection and homocysteine levels in patients with cardiac syndrome X. <i>Cardiology Journal</i> , 2013 , 20, 329 | 1.4 | 2 |
| 186 | Increased red cell distribution width in patients with slow coronary flow. Clinics, 2013, 68, 1288 | 2.3 | 2 |
| 185 | The relation between inflammation and coronary artery ectasia. <i>Revista Portuguesa De Cardiologia</i> (English Edition), 2016 , 35, 553-554 | Ο | 2 |
| 184 | Survival of the young patients with acute ST segment elevation myocardial infarction treated with primary percutaneous coronary intervention: Does gender matters?. <i>International Journal of Cardiology</i> , 2016 , 210, 54-5 | 3.2 | 2 |
| 183 | Ischemia modified albumin in cardiovascular disease. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 1699 | 2.9 | 2 |
| 182 | Left subclavian artery dissection and repair after transcatheter aortic valve implantation; the diameter of the axillary artery is important for axillary intervention. <i>International Journal of Cardiology</i> , 2016 , 204, 77-80 | 3.2 | 2 |
| 181 | Coronary microvascular dysfunction in patients with cardiac syndrome X: Ongoing debate. <i>International Journal of Cardiology</i> , 2016 , 218, 233-234 | 3.2 | 2 |
| 180 | Inter-Arm Blood Pressure Differences May Be Important for Predicting Mortality. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 164 | 2.3 | 2 |
| 179 | Asymmetric dimethylarginine levels in patients with schizophrenia. <i>Psychiatry Research</i> , 2015 , 227, 369- | 79 .9 | 1 |
| 178 | Electrocardiographic findings in patients with pulmonary embolism. <i>American Journal of Emergency Medicine</i> , 2015 , 33, 838-9 | 2.9 | 1 |
| 177 | Antistaphylococcal action of statins is worthy in infections after coronary artery bypass grafting. <i>International Journal of Cardiology</i> , 2015 , 201, 544 | 3.2 | 1 |
| 176 | Qrs prolongation and long-term prognosis in patients with dilated cardiomyopathy: Reflections from diphteritic myocarditis. <i>International Journal of Cardiology</i> , 2016 , 212, 295 | 3.2 | 1 |

(2013-2016)

| 175 | Acute anterior myocardial infarction after heavy exercise in a young sportsman: Importance of intravascular ultrasonography on differential diagnosis. <i>International Journal of Cardiology</i> , 2016 , 215, 169-72 | 3.2 | 1 |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 174 | The relation between coronary artery ectasia and psychological-environmental factors. <i>Journal of the Saudi Heart Association</i> , 2016 , 28, 127-8 | 0.7 | 1 |
| 173 | Duration of antiplatelet or anticoagulant therapy after transcatheter aortic valve implantation in high risk patients; longer or shorter. <i>International Journal of Cardiology</i> , 2016 , 210, 14-5 | 3.2 | 1 |
| 172 | The association of severe aortic stenosis and narrow aortic root in a young patientWhat is the etiology: Rheumatic valvulitis or LamblB excrescences?. <i>International Journal of Cardiology</i> , 2016 , 208, 87-91 | 3.2 | 1 |
| 171 | Long distance walking man after a combination of percutaneous mitral annuloplasty and coronary revascularization. <i>International Journal of Cardiology</i> , 2016 , 203, 1122-4 | 3.2 | 1 |
| 170 | Acceleration forces can effect cardiovascular structure. <i>Journal of Physiological Sciences</i> , 2014 , 64, 157-8 | 32.3 | 1 |
| 169 | The neutrophil lymphocyte ratio in patients with glioblastoma multiforme. <i>Journal of Neuro-Oncology</i> , 2014 , 117, 195-6 | 4.8 | 1 |
| 168 | Admission hyperglycemia may be the result of counterregulatory hormones during acute coronary syndrome events. <i>Angiology</i> , 2014 , 65, 160 | 2.1 | 1 |
| 167 | Higher mean platelet volume level in patients with pulmonary embolism. <i>Clinical Respiratory Journal</i> , 2014 , 8, 251-2 | 1.7 | 1 |
| 166 | Parameters influencing LVEF improvement with intracoronary bone marrow stem cell delivery in acute myocardial infarction. <i>International Journal of Cardiology</i> , 2014 , 177, 644-5 | 3.2 | 1 |
| 165 | Red cell distribution width may be related to the degree of coronary collateral circulation. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2014 , 20, 107 | 3.3 | 1 |
| 164 | Red cell distribution width in renal transplant patients. <i>International Urology and Nephrology</i> , 2014 , 46, 1465-6 | 2.3 | 1 |
| 163 | Heart rate variability in patients with chest pain at the ED. <i>American Journal of Emergency Medicine</i> , 2014 , 32, 284-5 | 2.9 | 1 |
| 162 | Vegetable and fruit intake protects from chronic heart failure. <i>International Journal of Cardiology</i> , 2013 , 168, 2985 | 3.2 | 1 |
| 161 | Closest friends: chronic pulmonary disease and systolic heart failure. <i>International Journal of Cardiology</i> , 2013 , 168, 2965 | 3.2 | 1 |
| 160 | Coronary lesions complexity in patients with stable coronary artery disease. <i>Angiology</i> , 2013 , 64, 310 | 2.1 | 1 |
| 159 | Inflammatory condition in coronary artery ectasia. <i>Angiology</i> , 2013 , 64, 637-8 | 2.1 | 1 |
| 158 | Is there any correlation between serum uric acid levels and right ventricular function parameters in patients with cardiovascular risk factors?. <i>International Journal of Cardiology</i> , 2013 , 168, 3086 | 3.2 | 1 |

| 157 | N-terminal pro-B-type natriuretic peptide should be used in all dyspneic patients in the ED. <i>American Journal of Emergency Medicine</i> , 2013 , 31, 1711 | 2.9 | 1 |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 156 | Renal failure in patients with acute heart failure. <i>International Journal of Cardiology</i> , 2013 , 168, e131 | 3.2 | 1 |
| 155 | Age and Aortic Diameters in Pilots. <i>Military Medicine</i> , 2015 , 180, 1262-7 | 1.3 | 1 |
| 154 | Red cell distribution width in patients with cerebral venous sinus thrombosis. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2015 , 21, 389 | 3.3 | 1 |
| 153 | Ambulatory Blood Pressure Monitoring and Body Mass Index: Good but Not Standardized!. <i>Medical Principles and Practice</i> , 2015 , 24, 500 | 2.1 | 1 |
| 152 | P-wave duration dispersion in patients with lichen planus. <i>Clinics</i> , 2014 , 69, 304 | 2.3 | 1 |
| 151 | Multiple pseudoaneurysms of aortic arch in a patient with Behcetß disease. <i>European Heart Journal Cardiovascular Imaging</i> , 2014 , 15, 1028 | 4.1 | 1 |
| 150 | Arterial stiffness parameters should be evaluated with different types of psoriasis. <i>Angiology</i> , 2014 , 65, 83 | 2.1 | 1 |
| 149 | Whether taken medication can improve arterial stiffness or not. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 693 | 2.3 | 1 |
| 148 | Arterial stiffness parameters in patients with chronic kidney disease. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 157 | 2.3 | 1 |
| 147 | The neutrophil lymphocyte ratio level in patients with end-stage renal disease. <i>Hemodialysis International</i> , 2014 , 18, 216-7 | 1.7 | 1 |
| 146 | Inter-arm blood pressure differences in aviators: is it important?. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 387 | 2.3 | 1 |
| 145 | Masked hypertension as an unrecognized destructive condition. <i>Journal of Clinical Hypertension</i> , 2014 , 16, 155 | 2.3 | 1 |
| 144 | Arterial stiffness in patients with ischemic attack. <i>International Journal of Stroke</i> , 2014 , 9, E5 | 6.3 | 1 |
| 143 | The etiology of Behët disease: is it really inflammation or not!. <i>Angiology</i> , 2014 , 65, 327 | 2.1 | 1 |
| 142 | eComment. Thrombus strikes back: promising role of thromboelastography for thromboembolic risk prediction in HeartMate II recipients. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2014 , 18, 465 | 1.8 | 1 |
| 141 | Increased platelet indices in acute stent thrombosis. <i>Angiology</i> , 2014 , 65, 744 | 2.1 | 1 |
| 140 | Mitral anterolateral papillary muscle rupture in an asymptomatic patient with mitral stenosis after percutaneous mitral balloon valvuloplasty. <i>Echocardiography</i> , 2012 , 29, E250-2 | 1.5 | 1 |

| 139 | The crush and culotte: two different stent techniques but same results in coronary bifurcations. <i>International Journal of Cardiology</i> , 2013 , 168, 2894-5 | 3.2 | 1 |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---|
| 138 | Letter by Kurtoglu et al regarding article, "Asymmetric dimethylarginine in response to recombinant tissue-type plasminogen activator and erythropoietin in acute stroke". <i>Stroke</i> , 2013 , 44, e229 | 6.7 | 1 |
| 137 | A new piece of puzzle: inflammation in the prediction of recurrence after successful electrical cardioversion in patients with nonvalvular atrial fibrillation. <i>Anatolian Journal of Cardiology</i> , 2013 , 13, 403-4 | | 1 |
| 136 | May chest pain describe coronary heart disease?. <i>Croatian Medical Journal</i> , 2013 , 54, 411 | 1.6 | 1 |
| 135 | The relationship between microalbuminuria, left ventricular hypertrophy, retinopathy, and sex hormone status in newly diagnosed hypertensive women. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 325-9 | 2.2 | 1 |
| 134 | Diastolic blood pressure may be related to the degree of coronary collateral circulation. <i>Angiology</i> , 2013 , 64, 553 | 2.1 | 1 |
| 133 | Mean platelet volume and mitral annular calcification. <i>Blood Coagulation and Fibrinolysis</i> , 2013 , 24, 899 | 1 | 1 |
| 132 | Diastolic dysfunction in patients with peripheral arterial disease: a frequent association. <i>Angiology</i> , 2013 , 64, 544-5 | 2.1 | 1 |
| 131 | Higher pentraxin-3 level in patients with metabolic syndrome. <i>Medical Principles and Practice</i> , 2013 , 22, 513-4 | 2.1 | 1 |
| 130 | Mean platelet volume as a surrogate marker of inflammation in patients with chronic hepatitis B. <i>European Journal of Gastroenterology and Hepatology</i> , 2013 , 25, 1364 | 2.2 | 1 |
| 129 | Triple antiplatelet therapy in obese patients undergoing stent implantation. <i>Angiology</i> , 2013 , 64, 559-6 | 02.1 | 1 |
| 128 | eComment. Video-assisted thoracoscopic surgery may be an alternative diagnosis and treatment option. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 16, 274 | 1.8 | 1 |
| 127 | Diastolic dyssynchrony should be assessed as well as systolic dyssynchrony in patients with chronic kidney disease. <i>Hemodialysis International</i> , 2013 , 17, 465 | 1.7 | 1 |
| 126 | Multimodality imaging of a right atrial thrombus obliterating inferior vena cava. <i>Echocardiography</i> , 2013 , 30, E145-7 | 1.5 | 1 |
| 125 | Lifestyle change important for patients with hypertension and cardiovascular diseases. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 858 | 2.3 | 1 |
| 124 | Response to Pmprovement of arterial stiffness in the transition from acute decompensated heart failure to chronic compensated heart failure? Clinical Cardiology, 2013, 36, E49 | 3.3 | 1 |
| 123 | Quadricuspid aortic valve without severe regurgitation in pectus excavatum. <i>Asian Cardiovascular and Thoracic Annals</i> , 2013 , 21, 240 | 0.6 | 1 |
| 122 | Response to Impact of lesion length on functional significance in intermediate coronary lesions. <i>Clinical Cardiology</i> , 2013 , 36, 301 | 3.3 | 1 |

| 121 | The importance of iodine supplementation and smoking for maternal and fetal thyroid health in pregnant women. <i>European Thyroid Journal</i> , 2013 , 2, 211-2 | 4.2 | 1 |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 120 | Hyperuricemia may be related to contrast-induced nephropathy after percutaneous coronary intervention. <i>Clinics</i> , 2013 , 68, 1071 | 2.3 | 1 |
| 119 | Pediatric guillain-barr syndrome and autonomic dysfunction. <i>Journal of Cardiovascular and Thoracic Research</i> , 2013 , 5, 179 | 1.3 | 1 |
| 118 | Eat as much as you burn - a good diet and eating less should be more important than an intense exercise program for decreasing morbidity and mortality. <i>Clinics</i> , 2013 , 68, 419 | 2.3 | 1 |
| 117 | Ankle-brachial index in coronary artery disease. <i>Clinics</i> , 2014 , 69, 653 | 2.3 | 1 |
| 116 | Right ventricular diastolic function in patients with thalassemia major. <i>Arquivos Brasileiros De Cardiologia</i> , 2013 , 101, 93-4 | 1.2 | 1 |
| 115 | AuthorsPcomments on: the platelet volume in patients with cardiac syndrome X. <i>Clinics</i> , 2013 , 68, 117 | 2.3 | 1 |
| 114 | All risk factors and confounders should be discussed in order to precisely describe the ankle-brachial index and albuminuria. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 601 | 4 | 1 |
| 113 | Which device should be chosen for the percutaneous closure of post-traumatic ventricular septal defects?. <i>Clinics</i> , 2013 , 68, 423 | 2.3 | 1 |
| 112 | Epicardial fat thickness and cardiovascular involvements. <i>African Health Sciences</i> , 2015 , 15, 1354-5 | 1.1 | 1 |
| 111 | Apelin Levels in Patients with Coronary Artery Ectasia. Korean Circulation Journal, 2016, 46, 431 | 2.2 | 1 |
| 110 | Vitamin D Levels in Patients with Atrial Fibrillation. <i>Annals of Noninvasive Electrocardiology</i> , 2016 , 21, 622-623 | 1.5 | 1 |
| 109 | Percutaneous transapical closure of cardiac apex and mitral prosthetic paravalvular leak is feasible and alternative approach to the septal approach. <i>General Thoracic and Cardiovascular Surgery</i> , 2016 , 64, 633 | 1.6 | 1 |
| 108 | The best treatment option in the management of patients with chronic total occlusion of a single coronary artery: Debate continues. <i>International Journal of Cardiology</i> , 2016 , 214, 79-80 | 3.2 | 1 |
| 107 | The role of combined electrocardiogram criteria in differential diagnosis of acute pericarditis: PR segment and QT interval. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 1309 | 2.9 | 1 |
| 106 | The more proximal implantation of percutaneous mitral annuloplasty device due to the circumflex artery compression may be less beneficial. <i>International Journal of Cardiology</i> , 2016 , 209, 46-8 | 3.2 | 1 |
| 105 | Blood Pressure Variability Provides Useful and Prognostic Information on the White-Coat Effect Among Older Patients. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 825 | 2.3 | 1 |
| 104 | The Relation Between C-Reactive Protein-Albumin Ratio and Carotid Intima-Media Thickness in Psoriasis <i>Angiology</i> , 2022 , 33197211072599 | 2.1 | 1 |

(2014-2016)

| 103 | The relation between flow mediated dilation and atrial fibrillation. <i>International Journal of Cardiology</i> , 2016 , 203, 1157 | 3.2 | О |
|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|---|
| 102 | The evaluation of renal hemodynamics changes in Familial Mediterranean fever with color Doppler sonography. <i>Renal Failure</i> , 2016 , 38, 1161-6 | 2.9 | O |
| 101 | The neutrophil lymphocyte ratio may be useful inflammatory indicator before applying other expensive and invasive procedures. <i>Indian Journal of Ophthalmology</i> , 2013 , 61, 685 | 1.6 | O |
| 100 | A new route to life in patients with circulatory shock: intraosseous route. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 922-3 | 2.9 | О |
| 99 | Is left ventricular diastolic function impaired in patients with ankylosing spondylitis?. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1802 | 2.3 | |
| 98 | Mean platelet volume and glomerular filtration rate: two important risk determinants in coronary artery disease. <i>Platelets</i> , 2015 , 26, 97-8 | 3.6 | |
| 97 | High altitude may affect cardiac structure in smoking subjects. <i>International Journal of Cardiology</i> , 2015 , 184, 416 | 3.2 | |
| 96 | Red cell distribution width: a predictor of all-cause mortality in patients with coronary artery disease. <i>Cardiology</i> , 2015 , 130, 23-4 | 1.6 | |
| 95 | Red cell distribution width in primary glomerulonephritides. Renal Failure, 2015, 37, 358 | 2.9 | |
| 94 | Subclinical atherosclerosis in patients with psoriasis. <i>Journal of the Saudi Heart Association</i> , 2015 , 27, 222-3 | 0.7 | |
| 93 | Right ventricular diastolic function in patients with community-acquired pneumonia. <i>American Journal of Emergency Medicine</i> , 2015 , 33, 1521-2 | 2.9 | |
| 92 | Increased frequency of mitral valve prolapse in patients with deviated nasal septum. <i>European Archives of Oto-Rhino-Laryngology</i> , 2015 , 272, 1667-71 | 3.5 | |
| 91 | Mean platelet volume in patients with systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2016 , 35, 547-8 | 3.9 | |
| 90 | Red Cell Distribution Width in Patients with Intermediate Coronary Artery Disease. <i>Medical Principles and Practice</i> , 2016 , 25, 397-8 | 2.1 | |
| 89 | The successful management of early rupture of abdominal aortic aneurysm with endovascular stenting and instent stenting for endoleak following EVAR; a case report. <i>International Journal of Cardiology</i> , 2016 , 207, 152-6 | 3.2 | |
| 88 | Subclinical Left Ventricular Dysfunction in Patients with Obstructive Sleep Apnea. <i>Medical Principles and Practice</i> , 2016 , 25, 299-300 | 2.1 | |
| 87 | The Pivotal Roles of Risk Factors for Acute Respiratory Distress Syndrome. <i>Journal of Emergency Medicine</i> , 2016 , 50, e67-8 | 1.5 | |
| 86 | Other factors may effect the relationship between heart rate variability indices and coronary atherosclerosis. <i>Clinical Autonomic Research</i> , 2014 , 24, 89 | 4.3 | |

| 85 | Ankle brachial index in hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2014 , 18, 113-4 | 1.9 |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 84 | The association between inflammatory markers and the degree of psoriasis vulgaris. <i>Angiology</i> , 2014 , 65, 81 | 2.1 |
| 83 | Carotid intima-media thickness as a novel inflammatory marker in psoriasis: comment on the article by Lin et al. <i>Arthritis Care and Research</i> , 2014 , 66, 793 | 4.7 |
| 82 | Adiponectin levels in hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2014 , 18, 214-5 | 1.9 |
| 81 | Three dimensional insight to Austin Flint murmur. <i>International Journal of Cardiology</i> , 2014 , 174, e24-5 | 3.2 |
| 80 | Multimodality imaging in the diagnosis of caseous calcification of mitral annulus. <i>Archives of Cardiovascular Diseases</i> , 2014 , 107, 140-1 | 2.7 |
| 79 | Uric acid level in patients with kidney disease. <i>Cardiology</i> , 2014 , 127, 25 | 1.6 |
| 78 | Red cell distribution width in patients with coronary artery disease. <i>Internal Medicine</i> , 2014 , 53, 1239 | 1.1 |
| 77 | Bicuspid aortic valve may affect aortic dimensions in aviators. <i>Aviation, Space, and Environmental Medicine</i> , 2014 , 85, 867 | |
| 76 | Response to Incremental predictive value of red cell distribution width for 12-month clinical outcome after acute myocardial infarction. <i>Clinical Cardiology</i> , 2013 , 36, E34-5 | 3.3 |
| 75 | Serum prolidase may not accurately provide information to clinicians about the psoriasis activity. <i>Archives of Dermatological Research</i> , 2013 , 305, 551-2 | 3.3 |
| 74 | Increased pulse wave velocity in patients with ulcerative colitis. <i>Digestive Diseases and Sciences</i> , 2013 , 58, 2738-9 | 4 |
| 73 | Plasma B-type natriuretic peptide level in patients with coronary artery bypass surgery. <i>Kaohsiung Journal of Medical Sciences</i> , 2013 , 29, 648-9 | 2.4 |
| 72 | Improvement in immunosuppressive treatment should decrease atherosclerosis and inflammation in renal transplant patients. <i>Renal Failure</i> , 2013 , 35, 432 | 2.9 |
| 71 | The relationship between hypertensive crisis and patient outcome. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 608 | 2.3 |
| 70 | Further studies on diastolic dysfunction in patients with airway obstruction should be kept in mind. <i>International Journal of Cardiology</i> , 2013 , 168, 2992 | 3.2 |
| 69 | Plasma homocysteine level in patients with Behët® disease. <i>International Journal of Rheumatic Diseases</i> , 2015 , 18, 582 | 2.3 |
| 68 | Utility of Isovolumic Contraction Peak Velocity for Evaluation of Adult Patient Status after Transcatheter Closure of Atrial Septal Defect. <i>Echocardiography</i> , 2015 , 32, 1742 | 1.5 |

(2013-2015)

| 67 | Body mass indexes should be stratified clearly in order to determine increased arterial stiffness in subjects with impaired fasting glucose. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 612 | 3.2 |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 66 | Mean platelet volume as a mirror of all inflammatory conditions. <i>European Journal of Ophthalmology</i> , 2014 , 24, 454-5 | 1.9 |
| 65 | Neutrophil-lymphocyte ratio is a predictor of saphenous vein graft patency in patients undergoing coronary artery bypass surgery. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2014 , 20, 219-20 | 3.3 |
| 64 | Arterial stiffness in patients with lower urinary tract symptoms. <i>Scandinavian Journal of Urology</i> , 2014 , 48, 225-6 | 1.6 |
| 63 | eComment. Doctor Jekyll and Mr Heyde: a vague association between angiodysplasia and aortic stenosis. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2014 , 19, 932 | 1.8 |
| 62 | Facial purpura as a complication of upper gastrointestinal endoscopy. <i>International Journal of Dermatology</i> , 2014 , 53, e134 | 1.7 |
| 61 | The effect of different circadian blood pressure rhythms on left ventricular systolic dyssynchrony in patients with newly diagnosed essential hypertension. <i>Echocardiography</i> , 2014 , 31, 120-1 | 1.5 |
| 60 | Arterial stiffness and pentraxin 3 levels in obese patients. <i>American Journal of Hypertension</i> , 2014 , 27, 881 | 2.3 |
| 59 | Current opinion: mean platelet volume is one of the most important parameters at the first glance. <i>Medical Principles and Practice</i> , 2014 , 23, 189 | 2.1 |
| 58 | Thrombotic therapy in patients with Behätß disease. <i>Expert Review of Cardiovascular Therapy</i> , 2014 , 12, 413-4 | 2.5 |
| 57 | Arterial stiffness may be associated with mitral annular calcification based on inflammatory condition. <i>Medical Principles and Practice</i> , 2014 , 23, 98 | 2.1 |
| 56 | Tiny magnetomers and future of magnetocardiography. <i>International Journal of Cardiology</i> , 2014 , 172, e268 | 3.2 |
| 55 | The QT measurements after biventricular pacing, is it reliable?. <i>Journal of the Saudi Heart Association</i> , 2014 , 26, 231-2 | 0.7 |
| 54 | Arterial stiffness in patients with bronchial asthma. Respiratory Medicine, 2014, 108, 940 | 4.6 |
| 53 | Mean platelet volume may be confused in many conditions. Wiener Klinische Wochenschrift, 2014 , 126, 248-9 | 2.3 |
| 52 | eComment. Cardiac venous arterialization in acute myocardial infarction. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 16, 313 | 1.8 |
| 51 | Diffuse mural thrombus in a patient with thoracic aortic aneurysm. <i>Asian Cardiovascular and Thoracic Annals</i> , 2013 , 21, 490 | 0.6 |
| 50 | eComment. Three-dimensional printers remodelling cardiac interventions. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 17, 1050 | 1.8 |

| 49 | eComment. Qualitative assesment of mitral annular calcification. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 17, 125-6 | 1.8 |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 48 | eComment. The benefits and risks of blood transfusions in patients undergoing cardiac surgery. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 17, 102 | 1.8 |
| 47 | eComment. Surgical or percutaneous pericardiocentesis in symptomatic pericardial effusion. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 16, 500 | 1.8 |
| 46 | Current opinions about coronary artery ectasia. <i>Angiology</i> , 2013 , 64, 634-5 | 2.1 |
| 45 | eComment. Renal dysfunction may predict early and late cardiovascular events. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 17, 643 | 1.8 |
| 44 | Size and subclasses of low-density lipoproteins: the role of smoking. <i>Angiology</i> , 2013 , 64, 562 | 2.1 |
| 43 | eComment. Atrial fibrillation in patients undergoing thoracic surgery. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2013 , 17, 686-7 | 1.8 |
| 42 | Systolic blood pressure can be lowered by strict volume control in hemodialysis patients. <i>Journal of Clinical Hypertension</i> , 2013 , 15, 607 | 2.3 |
| 41 | Patients should undergo conventional angiography to detect any suspected coronary artery lesions. <i>Internal Medicine</i> , 2013 , 52, 699 | 1.1 |
| 40 | Nocturnal intermittent hypoxemia and metabolic dyslipidemia. <i>Chest</i> , 2013 , 144, 357 | 5.3 |
| 39 | Smoking and the renal functional status of patients should be defined clearly in order to assess atherosclerosis and metabolic syndrome. <i>Internal Medicine</i> , 2013 , 52, 2389 | 1.1 |
| 38 | Chicken or egg? Which one is first? Myocardial infarction or low thyroid hormones?. <i>Internal Medicine</i> , 2013 , 52, 645 | 1.1 |
| 37 | Lifestyle change programs in the management of hyperlipidemia. <i>Oman Medical Journal</i> , 2012 , 27, 511 | 1.4 |
| 36 | C-reactive protein-albumin ratio and erectile dysfunction <i>Andrologia</i> , 2022 , e14386 | 2.4 |
| 35 | The relation between epicardial fat thickness and metabolic syndrome. <i>Journal of Geriatric Cardiology</i> , 2016 , 13, 369-70 | 1.7 |
| 34 | Red cell distribution width in subclinical hypothyroidism. <i>Arquivos Brasileiros De Endocrinologia E Metabologia</i> , 2014 , 58, 974-5 | |
| 33 | Anemia and renal dysfunction: Two different conditions but same results in cardiovascular disease. <i>Cardiology Journal</i> , 2015 , 22, 121 | 1.4 |
| 32 | Neutrophil-lymphocyte ratio: Can clinicians really trust it as an inflammatory indicator?. <i>Cardiology Journal</i> , 2015 , 22, 475 | 1.4 |

(2013-2016)

| 31 | Relation between epicardial fat thickness and chronic obstructive pulmonary disease. <i>Anatolian Journal of Cardiology</i> , 2016 , 16, 640 | 0.8 |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| 30 | Relationship between epicardial fat thickness and hypertension. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2016 , 44, 355-6 | 0.3 |
| 29 | Do spontaneous coronary artery dissections always need intervention in patients with no atherosclerosis?. <i>Anatolian Journal of Cardiology</i> , 2016 , 16, 365-6 | 0.8 |
| 28 | The Relation between Epicardial Fat Thickness and Prognostic Risk Scores. <i>Arquivos Brasileiros De Cardiologia</i> , 2016 , 107, 607 | 1.2 |
| 27 | Whole blood cell parameters help us predict the prognosis in patients with acute ST segment elevation myocardial infarction: Myth or reality?. <i>Cardiology Journal</i> , 2016 , 23, 357-8 | 1.4 |
| 26 | Gamma-glutamyl transferase and pulmonary embolism. Sao Paulo Medical Journal, 2016, 0 | 1.6 |
| 25 | Neutrophil-lymphocyte ratio and right ventricular dysfunction. Cardiology Journal, 2016, 23, 583 | 1.4 |
| 24 | Response to the letter regarding the article "Predictive value of admission red cell distribution width-platelet ratio for no-reflow phenomenon in acute ST segment elevation myocardial infarction undergoing primary percutaneous coronary intervention". <i>Cardiology Journal</i> , 2016 , 23, 589-5 | 1.4 90 |
| 23 | Differences in the mechanisms that induce obesity and metabolic syndrome in experimental animal models and humans may cause treatment failure. <i>Clinics</i> , 2013 , 68, 119 | 2.3 |
| 22 | Carotid intima-media thickness is a relatively inexpensive and favorable prognostic marker in patients with spondyloarthritis. <i>Sao Paulo Medical Journal</i> , 2013 , 131, 436-8 | 1.6 |
| 21 | Prediction of hospital events based on the severity of illness. <i>Clinics</i> , 2013 , 68, 121 | 2.3 |
| 20 | Mean platelet volume may be associated with extent of coronary artery disease. <i>Arquivos Brasileiros De Cardiologia</i> , 2013 , 101, 284-5 | 1.2 |
| 19 | Any variation existing in pathways inducing hyperlipidemia in experimental animal models and in real humans may cause difference in treatment success. <i>Oman Medical Journal</i> , 2013 , 28, 75 | 1.4 |
| 18 | Early age at onset of diabetes mellitus may be worse in terms of metabolic side effects. <i>Oman Medical Journal</i> , 2013 , 28, 151 | 1.4 |
| 17 | It is important to control for confounders when examining the role of diet in cardiovascular disease prevention. <i>Clinics</i> , 2013 , 68, 575 | 2.3 |
| 16 | P2 scallop prolapsus resulting from chordae tendineae rupture detected by threedimensional echocardiography. <i>Kardiologia Polska</i> , 2013 , 71, 429 | 0.9 |
| 15 | Further studies should evaluate multiple predispositions in heart failure prognosis. <i>Cardiology Journal</i> , 2013 , 20, 211 | 1.4 |
| 14 | Future studies should consider multiple predisposing conditions in predicting weaning failure from mechanical ventilation in patients after cardiac surgery. <i>Clinics</i> , 2013 , 68, 725 | 2.3 |

| 13 | Modified Hodge test as screening test for spreading carbapenemase resistance has become more important. <i>Clinics</i> , 2013 , 68, 1175 | 2.3 |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| 12 | The association between vitamin D level and extent of coronary stenotic lesions. <i>Cardiology Journal</i> , 2014 , 21, 206-7 | 1.4 |
| 11 | Red Cell Distribution Width and Coronary Artery Calcification. <i>Korean Circulation Journal</i> , 2016 , 46, 270 | 0-2.2 |
| 10 | Nebivolol and Endothelial Dysfunction in Patients With Essential Hypertension: A Reputation Saver of Eblockers?. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 1258-1259 | 2.3 |
| 9 | Coronary Obstruction During Transcatheter Aortic Valve Replacement: Related to Calcification or Thrombus?. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2016 , 69, 456-7 | 0.7 |
| 8 | The relation between signal peptide-CUB-EGF domain-containing protein 1 and coronary artery disease. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 647-8 | 2.9 |
| 7 | The effectiveness of bilirubin for the treatment of ischemia-reperfusion injury in an experimental model: Can it be extrapolated to humans?. <i>International Journal of Cardiology</i> , 2016 , 215, 173-4 | 3.2 |
| 6 | eComment. Rise of Ticagrelor and the vital necessity for specific reversal agents. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2016 , 22, 410 | 1.8 |
| 5 | Sodium bicarbonate dilemma in patients with out-of-hospital cardiac arrest: A double-edged sword. <i>American Journal of Emergency Medicine</i> , 2016 , 34, 1314-5 | 2.9 |
| 4 | Red cell distribution width in patients with sarcoidosis. <i>Clinical Respiratory Journal</i> , 2016 , 10, 129-30 | 1.7 |
| 3 | Red cell distribution width and mean platelet volume in carbon monoxide poisoning. <i>American Journal of Emergency Medicine</i> , 2019 , 37, 1196-1197 | 2.9 |
| 2 | Re: correlation between ankle-brachial index and microalbuminuria in type 2 diabetes mellitus. <i>Iranian Journal of Kidney Diseases</i> , 2013 , 7, 415-6 | 0.9 |
| 1 | Atherosclerosis and Non-Alcoholic Fatty Liver Disease <i>Angiology</i> , 2022 , 33197221091317 | 2.1 |