
Wolfram Jochum

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/560066/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Rapid mass spectrometric conversion of tissue biopsy samples into permanent quantitative digital proteome maps. Nature Medicine, 2015, 21, 407-413.	30.7	358
2	Cancer genetics-guided discovery of serum biomarker signatures for diagnosis and prognosis of prostate cancer. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108, 3342-3347.	7.1	175
3	Golgi phosphoprotein 2 (GOLPH2) expression in liver tumors and its value as a serum marker in hepatocellular carcinomas. Hepatology, 2009, 49, 1602-1609.	7.3	110
4	Highâ€ŧhroughput proteomic analysis of <scp>FFPE</scp> tissue samples facilitates tumor stratification. Molecular Oncology, 2019, 13, 2305-2328.	4.6	100
5	Screening for ALK in non-small cell lung carcinomas: 5A4 and D5F3 antibodies perform equally well, but combined use with FISH is recommended. Lung Cancer, 2015, 89, 104-109.	2.0	69
6	IMP3 expression in lesions of the biliary tract: a marker for high-grade dysplasia and an independent prognostic factor in bile duct carcinomas. Human Pathology, 2009, 40, 1377-1383.	2.0	66
7	Frequent expression of the novel cancer testis antigen MAGEâ€C2/CTâ€10 in hepatocellular carcinoma. International Journal of Cancer, 2009, 124, 352-357.	5.1	63
8	Expression of the extracellular matrix protein periostin in liver tumours and bile duct carcinomas. Histopathology, 2010, 56, 600-606.	2.9	52
9	Multi-region proteome analysis quantifies spatial heterogeneity of prostate tissue biomarkers. Life Science Alliance, 2018, 1, e201800042.	2.8	51
10	Prognostic significance of nuclear DNA content and proliferative activity in renal cell carcinomas: A clinicopathologic study of 58 patients using mitotic count, MIB-1 staining, and DNA cytophotometry. Cancer, 1996, 77, 514-521.	4.1	44
11	Cell adhesion molecules P-cadherin and CD24 are markers for carcinoma and dysplasia in the biliary tract. Human Pathology, 2010, 41, 1558-1565.	2.0	36
12	Evaluation of type-specific antibodies to high risk-human papillomavirus (HPV) proteins in patients with oropharyngeal cancer. Oral Oncology, 2017, 70, 43-50.	1.5	28
13	Expression and Clinicopathological Significance of Notch Signaling and Cell-Fate Genes in Biliary Tract Cancer. American Journal of Gastroenterology, 2012, 107, 126-132.	0.4	25
14	NGS-based BRCA1/2 mutation testing of high-grade serous ovarian cancer tissue: results and conclusions of the first international round robin trial. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2016, 468, 697-705.	2.8	24
15	Prognostic significance of cell cycle-associated proteins p16, pRB, cyclin D1 and p53 in resected oropharyngeal carcinoma. Journal of Otolaryngology - Head and Neck Surgery, 2018, 47, 53.	1.9	22
16	Comparison of automated and manual FISH for evaluation of HER2 gene status on breast carcinoma core biopsies. BMC Clinical Pathology, 2013, 13, 13.	1.8	18
17	Brush cytology for the detection of highâ€risk HPV infection in oropharyngeal squamous cell carcinoma. Cancer Cytopathology, 2015, 123, 732-738.	2.4	17
18	Impact of human papillomavirus on outcome in patients with oropharyngeal cancer treated with primary surgery. Head and Neck, 2017, 39, 2004-2015.	2.0	14

#	Article	IF	CITATIONS
19	Multi-laboratory proficiency testing of clinical cancer genomic profiling by next-generation sequencing. Pathology Research and Practice, 2018, 214, 957-963.	2.3	11
20	Cost-Efficient and Easy to Perform PCR-Based Assay to Identify Met Exon 14 Skipping in Formalin-Fixed Paraffin-Embedded (FFPE) Non-Small Cell Lung Cancer (NSCLC) Samples. Diagnostics, 2019, 9, 13.	2.6	10
21	Prognostic significance of nuclear DNA content and proliferative activity in renal cell carcinomas: A clinicopathologic study of 58 patients using mitotic count, MIBâ€1 staining, and DNA cytophotometry. Cancer, 1996, 77, 514-521.	4.1	4