Damien Réveillon

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5605254/publications.pdf

Version: 2024-02-01

20 453
papers citations

759233 752698 20
h-index g-index

20 20 all docs citations

20 times ranked 491 citing authors

#	Article	IF	CITATIONS
1	Toxicity of palytoxin, purified ovatoxin-a, ovatoxin-d and extracts of Ostreopsis cf. ovata on the Caco-2 intestinal barrier model. Environmental Toxicology and Pharmacology, 2022, 94, 103909.	4.0	3
2	Molecular networking as a novel approach to unravel toxin diversity of four strains of the dominant Dinophysis species from French coastal waters. Harmful Algae, 2021, 103, 102026.	4.8	4
3	Combined effects of temperature and light intensity on growth, metabolome and ovatoxin content of a Mediterranean Ostreopsis cf. ovata strain. Harmful Algae, 2021, 106, 102060.	4.8	6
4	Effect of a short-term salinity stress on the growth, biovolume, toxins, osmolytes and metabolite profiles on three strains of the Dinophysis acuminata-complex (Dinophysis cf. sacculus). Harmful Algae, 2021, 107, 102009.	4.8	8
5	Deeper insight into Gambierdiscus polynesiensis toxin production relies on specific optimization of high-performance liquid chromatography-high resolution mass spectrometry. Talanta, 2021, 232, 122400.	5.5	7
6	Tetrodotoxins in French Bivalve Mollusks—Analytical Methodology, Environmental Dynamics and Screening of Bacterial Strain Collections. Toxins, 2021, 13, 740.	3.4	12
7	Sulfo-Gambierones, Two New Analogs of Gambierone Produced by Gambierdiscus excentricus. Marine Drugs, 2021, 19, 657.	4.6	13
8	Toxin content of Ostreopsis cf. ovata depends on bloom phases, depth and macroalgal substrate in the NW Mediterranean Sea. Harmful Algae, 2020, 92, 101727.	4.8	23
9	Centrodinium punctatum (Dinophyceae) produces significant levels of saxitoxin and related analogs. Harmful Algae, 2020, 100, 101923.	4.8	16
10	Taxonomy and toxicity of a bloom-forming Ostreopsis species (Dinophyceae, Gonyaulacales) in Tahiti island (South Pacific Ocean): one step further towards resolving the identity of O. siamensis Harmful Algae, 2020, 98, 101888.	4.8	12
11	Salt Shock Responses of Microcystis Revealed through Physiological, Transcript, and Metabolomic Analyses. Toxins, 2020, 12, 192.	3.4	15
12	Combined Effects of Temperature, Irradiance, and <scp>pH</scp> on <i>Teleaulax amphioxeia</i> (Cryptophyceae) Physiology and Feeding Ratio For Its Predator <i>Mesodinium rubrum</i> (Ciliophora) ¹ . Journal of Phycology, 2020, 56, 775-783.	2.3	8
13	Cultures of Dinophysis sacculus, D.Âacuminata and pectenotoxin 2 affect gametes and fertilization success of the Pacific oyster, Crassostrea gigas. Environmental Pollution, 2020, 265, 114840.	7. 5	16
14	Physiological and Metabolic Responses of Freshwater and Brackish-Water Strains of Microcystis aeruginosa Acclimated to a Salinity Gradient: Insight into Salt Tolerance. Applied and Environmental Microbiology, 2019, 85, .	3.1	27
15	Ostreopsis lenticularis Y. Fukuyo (Dinophyceae, Gonyaulacales) from French Polynesia (South Pacific) Tj ETQq1 1 C). <u>7</u> .84314 r	iggT/Overlo
16	Exploring the chemodiversity of tropical microalgae for the discovery of natural antifouling compounds. Journal of Applied Phycology, 2019, 31, 319-333.	2.8	10
17	Production of BMAA and DAB by diatoms (Phaeodactylum tricornutum, Chaetoceros sp., Chaetoceros) Tj ETQq1 Algae, 2016, 58, 45-50.		.4 rgBT /Ove 61
18	Systematic detection of BMAA (\hat{l}^2 -N-methylamino-l-alanine) and DAB (2,4-diaminobutyric acid) in mollusks collected in shellfish production areas along the French coasts. Toxicon, 2016, 110, 35-46.	1.6	54

#	Article	IF	CITATIONS
19	\hat{l}^2 -N-methylamino-l-alanine (BMAA) and isomers: Distribution in different food web compartments of Thau lagoon, French Mediterranean Sea. Marine Environmental Research, 2015, 110, 8-18.	2.5	73
20	Beta-N-Methylamino-l-Alanine: LC-MS/MS Optimization, Screening of Cyanobacterial Strains and Occurrence in Shellfish from Thau, a French Mediterranean Lagoon. Marine Drugs, 2014, 12, 5441-5467.	4.6	56