## Xiang-Chuan Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5623725/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	A unique record of <i>Cercis</i> from the late early Miocene of interior Asia and its significance for paleoenvironments and paleophytogeography. Journal of Systematics and Evolution, 2021, 59, 1321-1338.	3.1	8
2	Ormosia (Fabaceae: Faboideae) from the Miocene of southeastern China support historical expansion of the tropical genus in East Asia. Historical Biology, 2021, 33, 3561-3578.	1.4	7
3	An Extinct Fruit Species of Fabaceae from the Early Eocene of Northwestern Wyoming, USA. International Journal of Plant Sciences, 2021, 182, 730-746.	1.3	1
4	New records of Podocarpium A. Braun ex Stizenberger (Fabaceae) from the Oligocene to Miocene of China: Reappraisal of the phylogeographical history of the genus. Review of Palaeobotany and Palynology, 2019, 260, 38-50.	1.5	14
5	An ancient example of Platycladus (Cupressceae) from the early Miocene of northern China: origin and biogeographical implications. Historical Biology, 2018, 30, 1123-1131.	1.4	2
6	The first fossil brown lacewing from the Miocene of the Tibetan Plateau (Neuroptera, Hemerobiidae). ZooKeys, 2018, 726, 145-154.	1.1	4
7	Miocene pollen assemblages from the Zeku Basin, northeastern Tibetan Plateau, and their palaeoecological and palaeoaltimetric implications. Palaeogeography, Palaeoclimatology, Palaeoecology, 2018, 511, 419-432.	2.3	14
8	First report of Cixiidae insect fossils from the Miocene of the northeastern Tibetan Plateau and their palaeoenvironmental implications. Alcheringa, 2017, 41, 54-60.	1.2	7
9	Fossil fruits of Koelreuteria (Sapindaceae) from the Miocene of northeastern Tibetan Plateau and their palaeoenvironmental, phytogeographic and phylogenetic implications. Review of Palaeobotany and Palynology, 2016, 234, 125-135.	1.5	9
10	<i>Paliurus</i> (Paliureae, Rhamnaceae) from the Miocene of East China and its Macrofossil-based Phylogenetic and Phytogeographical History. Acta Geologica Sinica, 2014, 88, 1364-1377.	1.4	13
11	Stable isotope compositions of recent and fossil sun/shade leaves and implications for palaeoenvironmental reconstruction. Review of Palaeobotany and Palynology, 2013, 190, 75-84.	1.5	10
12	Cunninghamia praelanceolata sp. nov. with associated epiphyllous fungi from the upper Miocene of eastern Zhejiang, S.E China and their palaeoecological implications. Review of Palaeobotany and Palynology, 2012, 182, 32-43.	1.5	34
13	Reconstructing Neogene vegetation and climates to infer tectonic uplift in western Yunnan, China. Palaeogeography, Palaeoclimatology, Palaeoecology, 2011, 304, 328-336.	2.3	144
14	Miocene Smilax leaves and associated epiphyllous fungi from Zhejiang, East China and their paleoecological implications. Review of Palaeobotany and Palynology, 2011, 165, 209-223.	1.5	37
15	Anatomical variations of living and fossil Liquidambar leaves: A proxy for paleoenvironmental reconstruction. Science China Earth Sciences, 2011, 54, 493-508.	5.2	19
16	Leaf macrofossils of llex protocornuta sp. nov. (Aquifoliaceae) from the Late Miocene of East China: Implications for palaeoecology. Review of Palaeobotany and Palynology, 2010, 161, 87-103.	1.5	18
17	Leaf megafossils of Betula yunnanensis sp. nov. (Betulaceae) from the Mangbang Formation, SW China and its taphonomic implications. Review of Palaeobotany and Palynology, 2010, 163, 84-103.	1.5	2
18	Response to paleoatmospheric CO2 concentration of Solenites vimineus (Phillips) Harris (Ginkgophyta) from the Middle Jurassic of the Yaojie Basin, Gansu Province, China. Science in China Series D: Earth Sciences, 2009, 52, 2029-2039.	0.9	25