

# Ayumi Fukuoka

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5627306/publications.pdf>

Version: 2024-02-01

15  
papers

427  
citations

840119

11  
h-index

1058022

14  
g-index

19  
all docs

19  
docs citations

19  
times ranked

876  
citing authors

#	ARTICLE	IF	CITATIONS
1	The role of basophils and proallergic cytokines, TSLP and IL-33, in cutaneously sensitized food allergy. <i>International Immunology</i> , 2014, 26, 539-549.	1.8	103
2	Diesel exhaust particles exacerbate allergic rhinitis in mice by disrupting the nasal epithelial barrier. <i>Clinical and Experimental Allergy</i> , 2016, 46, 142-152.	1.4	62
3	Barrier dysfunction in the nasal allergy. <i>Allergology International</i> , 2018, 67, 18-23.	1.4	46
4	Murine allergic rhinitis and nasal T h2 activation are mediated via TSLP- and IL-33-signaling pathways. <i>International Immunology</i> , 2016, 28, 65-76.	1.8	45
5	Human cystatin SN is an endogenous protease inhibitor that prevents allergic rhinitis. <i>Journal of Allergy and Clinical Immunology</i> , 2019, 143, 1153-1162.e12.	1.5	35
6	Allergen endotoxins induce T-cellâ€‘dependent and nonâ€‘IgE-mediated nasal hypersensitivity in mice. <i>Journal of Allergy and Clinical Immunology</i> , 2017, 139, 258-268.e10.	1.5	27
7	Identification of a novel type 2 innate immunocyte with the ability to enhance IgE production. <i>International Immunology</i> , 2013, 25, 373-382.	1.8	20
8	Placental chemokine compartmentalisation: A novel mammalian molecular control mechanism. <i>PLoS Biology</i> , 2019, 17, e3000287.	2.6	18
9	Regnase-1 degradation is crucial for IL-33â€‘ and IL-25â€‘mediated ILC2 activation. <i>JCI Insight</i> , 2020, 5, .	2.3	18
10	Chemokine receptors coordinately regulate macrophage dynamics and mammary gland development. <i>Development (Cambridge)</i> , 2020, 147, .	1.2	15
11	Pathogenic Th2-type follicular helper T cells contribute to the development of lupus in <i>Fas</i> -deficient mice. <i>International Immunology</i> , 2014, 26, 221-231.	1.8	12
12	Activation of group 2 innate lymphoid cells exacerbates and confers corticosteroid resistance to mouse nasal type 2 inflammation. <i>International Immunology</i> , 2017, 29, 221-233.	1.8	11
13	Fas deficiency in mice with the Balb/c background induces blepharitis with allergic inflammation and hyper-IgE production in conjunction with severe autoimmune disease. <i>International Immunology</i> , 2013, 25, 287-293.	1.8	9
14	Diverse myeloid cells are recruited to the developing and inflamed mammary gland. <i>Immunology</i> , 2022, 165, 206-218.	2.0	4
15	The roles of basophils, TSLP and ILâ€‘33 in food allergy following epicutaneous sensitisation. <i>Clinical and Translational Allergy</i> , 2015, 5, O17.	1.4	0