

Naoto Takahashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5629997/publications.pdf>

Version: 2024-02-01

125
papers

2,499
citations

218381

26
h-index

233125

45
g-index

130
all docs

130
docs citations

130
times ranked

2864
citing authors

#	ARTICLE	IF	CITATIONS
1	Lineage Involvement of Stem Cells Bearing the Philadelphia Chromosome in Chronic Myeloid Leukemia in the Chronic Phase as Shown by a Combination of Fluorescence-Activated Cell Sorting and Fluorescence In Situ Hybridization. <i>Blood</i> , 1998, 92, 4758-4763.	0.6	161
2	A Clinical Analysis of 52 Adult Patients With Hemophagocytic Syndrome: The Prognostic Significance of the Underlying Diseases. <i>International Journal of Hematology</i> , 2001, 74, 209-213.	0.7	153
3	Influence of CYP3A5 and drug transporter polymorphisms on imatinib trough concentration and clinical response among patients with chronic phase chronic myeloid leukemia. <i>Journal of Human Genetics</i> , 2010, 55, 731-737.	1.1	147
4	Discontinuation of imatinib in Japanese patients with chronic myeloid leukemia. <i>Haematologica</i> , 2012, 97, 903-906.	1.7	138
5	Treatment-Free Remission After Second-Line Nilotinib Treatment in Patients With Chronic Myeloid Leukemia in Chronic Phase. <i>Annals of Internal Medicine</i> , 2018, 168, 461.	2.0	105
6	A Clinicopathological Study of 20 Patients With T/Natural Killer (NK)-Cell Lymphoma-Associated Hemophagocytic Syndrome With Special Reference to Nasal and Nasal-Type NK/T-Cell Lymphoma. <i>International Journal of Hematology</i> , 2001, 74, 303-308.	0.7	93
7	Deeper molecular response is a predictive factor for treatment-free remission after imatinib discontinuation in patients with chronic phase chronic myeloid leukemia: the JALSG-STIM213 study. <i>International Journal of Hematology</i> , 2018, 107, 185-193.	0.7	72
8	Fluorescence In Situ Hybridization of Progenitor Cells Obtained by Fluorescence-Activated Cell Sorting for the Detection of Cells Affected by Chromosome Abnormality Trisomy 8 in Patients With Myelodysplastic Syndromes. <i>Blood</i> , 1998, 92, 2886-2892.	0.6	67
9	Treatment-free remission after two-year consolidation therapy with nilotinib in patients with chronic myeloid leukemia: STAT2 trial in Japan. <i>Haematologica</i> , 2018, 103, 1835-1842.	1.7	59
10	Tyrosine kinase inhibitor imatinib augments tumor immunity by depleting effector regulatory T cells. <i>Journal of Experimental Medicine</i> , 2020, 217, .	4.2	58
11	A synthetic double-stranded RNA, poly I:C, induces a rapid apoptosis of human CD34+ cells. <i>Experimental Hematology</i> , 2012, 40, 330-341.	0.2	52
12	Hypoxia-inducible KDM3A addiction in multiple myeloma. <i>Blood Advances</i> , 2018, 2, 323-334.	2.5	50
13	An integrative model of pathway convergence in genetically heterogeneous blast crisis chronic myeloid leukemia. <i>Blood</i> , 2020, 135, 2337-2353.	0.6	49
14	Influence of H2-receptor antagonists and proton pump inhibitors on dasatinib pharmacokinetics in Japanese leukemia patients. <i>Cancer Chemotherapy and Pharmacology</i> , 2012, 69, 999-1004.	1.1	43
15	Routine therapeutic drug monitoring of tyrosine kinase inhibitors by HPLC-UV or LC-MS/MS methods. <i>Drug Metabolism and Pharmacokinetics</i> , 2016, 31, 12-20.	1.1	41
16	Molecular heterogeneity of the NUP98/HOXA9 fusion transcript in myelodysplastic syndromes associated with t(7;11)(p15;p15). <i>British Journal of Haematology</i> , 1999, 107, 600-604.	1.2	40
17	Clonal evolution and clinical implications of genetic abnormalities in blastic transformation of chronic myeloid leukaemia. <i>Nature Communications</i> , 2021, 12, 2833.	5.8	39
18	Quantitative Determination of Imatinib in Human Plasma with High-Performance Liquid Chromatography and Ultraviolet Detection. <i>Journal of Chromatographic Science</i> , 2011, 49, 412-415.	0.7	37

#	ARTICLE	IF	CITATIONS
19	Therapeutic Drug Monitoring of Imatinib for Chronic Myeloid Leukemia Patients in the Chronic Phase. <i>Pharmacology</i> , 2011, 87, 241-248.	0.9	36
20	Influence of UGT1A1 *6, *27, and *28 Polymorphisms on Nilotinib-induced Hyperbilirubinemia in Japanese Patients with Chronic Myeloid Leukemia. <i>Drug Metabolism and Pharmacokinetics</i> , 2014, 29, 449-454.	1.1	35
21	Hypoxia-inducible hexokinase-2 enhances anti-apoptotic function via activating autophagy in multiple myeloma. <i>Cancer Science</i> , 2020, 111, 4088-4101.	1.7	34
22	Ponatinib in Japanese patients with Philadelphia chromosome-positive leukemia, a phase 1/2 study. <i>International Journal of Hematology</i> , 2017, 106, 385-397.	0.7	33
23	Involvement of natural killer cells in patients with myelodysplastic syndrome carrying monosomy 7 revealed by the application of fluorescence in situ hybridization to cells collected by means of fluorescence-activated cell sorting. <i>British Journal of Haematology</i> , 2000, 110, 876-879.	1.2	32
24	Hypoxia-inducible microRNA-210 regulates the DIMT1-IRF4 oncogenic axis in multiple myeloma. <i>Cancer Science</i> , 2017, 108, 641-652.	1.7	31
25	High-performance liquid chromatography with solid-phase extraction for the quantitative determination of nilotinib in human plasma. <i>Biomedical Chromatography</i> , 2010, 24, 789-793.	0.8	30
26	Effect of itraconazole on the concentrations of tacrolimus and cyclosporine in the blood of patients receiving allogeneic hematopoietic stem cell transplants. <i>European Journal of Clinical Pharmacology</i> , 2013, 69, 1321-1329.	0.8	30
27	A multicenter clinical study evaluating the confirmed complete molecular response rate in imatinib-treated patients with chronic phase chronic myeloid leukemia by using the international scale of real-time quantitative polymerase chain reaction. <i>Haematologica</i> , 2013, 98, 1407-1413.	1.7	29
28	Efficacy and safety of tyrosine kinase inhibitors for newly diagnosed chronic-phase chronic myeloid leukemia over a 5-year period: results from the Japanese registry obtained by the New TARGET system. <i>International Journal of Hematology</i> , 2019, 109, 426-439.	0.7	29
29	Histone deacetylase inhibitors inhibit metastasis by restoring a tumor suppressive microRNA-150 in advanced cutaneous T-cell lymphoma. <i>Oncotarget</i> , 2017, 8, 7572-7585.	0.8	27
30	Drug interaction between lenalidomide and itraconazole. <i>American Journal of Hematology</i> , 2012, 87, 338-339.	2.0	25
31	Itraconazole Oral Solution Enhanced Vincristine Neurotoxicity in Five Patients with Malignant Lymphoma. <i>Internal Medicine</i> , 2008, 47, 651-653.	0.3	24
32	Kidney-limited intravascular large B cell lymphoma: a distinct variant of IVLBCL?. <i>International Journal of Hematology</i> , 2009, 89, 533-537.	0.7	24
33	Erythroblast enucleation is a dynein-dependent process. <i>Experimental Hematology</i> , 2016, 44, 247-256.e12.	0.2	24
34	A phase 1/2 study of bosutinib in Japanese adults with Philadelphia chromosome-positive chronic myeloid leukemia. <i>International Journal of Hematology</i> , 2015, 101, 154-164.	0.7	23
35	Therapeutic drug monitoring of ponatinib using a simple high-performance liquid chromatography method in Japanese patients. <i>Leukemia Research</i> , 2018, 64, 42-45.	0.4	23
36	Pharmacokinetics of dasatinib for Philadelphia-positive acute lymphocytic leukemia with acquired T315I mutation. <i>Journal of Hematology and Oncology</i> , 2012, 5, 23.	6.9	21

#	ARTICLE	IF	CITATIONS
37	Correlation of plasma concentration and adverse effects of bosutinib: standard dose or dose-escalation regimens of bosutinib treatment for patients with chronic myeloid leukemia. <i>Experimental Hematology and Oncology</i> , 2018, 7, 9.	2.0	21
38	JSH Practical Guidelines for Hematological Malignancies, 2018: I. Leukemia-4. Chronic myeloid leukemia (CML)/myeloproliferative neoplasms (MPN). <i>International Journal of Hematology</i> , 2020, 112, 268-291.	0.7	21
39	Disruption of CCL20-CCR6 interaction inhibits metastasis of advanced cutaneous T-cell lymphoma. <i>Oncotarget</i> , 2016, 7, 13563-13574.	0.8	21
40	Multicenter phase II clinical trial of nilotinib for patients with imatinib-resistant or -intolerant chronic myeloid leukemia from the East Japan CML study group evaluation of molecular response and the efficacy and safety of nilotinib. <i>Biomarker Research</i> , 2014, 2, 6.	2.8	20
41	Rearrangements of the BCL6 Gene and Chromosome Aberrations Affecting 3q27 in 54 Patients with Non-Hodgkin's Lymphoma. <i>Leukemia and Lymphoma</i> , 1997, 27, 329-334.	0.6	18
42	Long-term treatment-free remission in patients with chronic myeloid leukemia after second-line nilotinib: ENESTop 5-year update. <i>Leukemia</i> , 2021, 35, 1631-1642.	3.3	18
43	ATP produced by anaerobic glycolysis is essential for enucleation of human erythroblasts. <i>Experimental Hematology</i> , 2019, 72, 14-26.e1.	0.2	17
44	Clinical features of adult acute leukemia with 11q23 abnormalities in Japan: a co-operative multicenter study. <i>International Journal of Hematology</i> , 2008, 87, 195-202.	0.7	16
45	The localization of α -synuclein in the process of differentiation of human erythroid cells. <i>International Journal of Hematology</i> , 2018, 108, 130-138.	0.7	16
46	Treatment-Free Remission in Patients with Chronic Myeloid Leukemia in Chronic Phase According to Reasons for Switching from Imatinib to Nilotinib: Subgroup Analysis from ENESTop. <i>Blood</i> , 2016, 128, 792-792.	0.6	16
47	BCR-ABL Activates IGF-1 Expression and Signaling in Chronic Myelogenous Leukemia Blast Crisis Cell Lines. <i>Blood</i> , 2006, 108, 1932-1932.	0.6	16
48	Molecular features of a new human lymphoma cell line carrying both BCL2 and BCL6 gene rearrangements. <i>Oncogene</i> , 1998, 17, 971-979.	2.6	15
49	11q23 Aberration is an additional chromosomal change in de novo acute leukemia after treatment with etoposide and mitoxantrone. , 1996, 53, 264-266.		14
50	Fluorescence In Situ Hybridization Monitoring of BCR-ABL-Positive Neutrophils in Chronic-Phase Chronic Myeloid Leukemia Patients during the Primary Stage of Imatinib Mesylate Therapy. <i>International Journal of Hematology</i> , 2005, 81, 235-241.	0.7	14
51	Drug interaction of (S)-warfarin, and not (R)-warfarin, with itraconazole in a hematopoietic stem cell transplant recipient. <i>Clinica Chimica Acta</i> , 2011, 412, 2002-2006.	0.5	14
52	IL-6 Generated from Human Hematopoietic Stem and Progenitor Cells through TLR4 Signaling Promotes Emergency Granulopoiesis by Regulating Transcription Factor Expression. <i>Journal of Immunology</i> , 2021, 207, 1078-1086.	0.4	14
53	Fatal hemorrhagic pneumonia caused by <i>Stenotrophomonas maltophilia</i> in a patient with non-Hodgkin lymphoma. <i>Journal of Infection and Chemotherapy</i> , 2011, 17, 858-862.	0.8	13
54	Prognostic effect of comorbidities in patients with chronic myeloid leukemia treated with a tyrosine kinase inhibitor. <i>Cancer Science</i> , 2020, 111, 3714-3725.	1.7	13

#	ARTICLE	IF	CITATIONS
55	Effects of CYP3A5 polymorphism on the pharmacokinetics of a once-daily modified-release tacrolimus formulation and acute kidney injury in hematopoietic stem cell transplantation. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 111-118.	1.1	12
56	Acute Myelogenous Leukemia Associated with a Mediastinal Tumor. <i>Leukemia and Lymphoma</i> , 1993, 12, 143-146.	0.6	11
57	Pharmacokinetics of nilotinib in imatinib-resistant/intolerant chronic myeloid leukemia patients on hemodialysis for chronic renal failure. <i>American Journal of Hematology</i> , 2012, 87, 451-451.	2.0	11
58	Cdc42 regulates cell polarization and contractile actomyosin rings during terminal differentiation of human erythroblasts. <i>Scientific Reports</i> , 2020, 10, 11806.	1.6	11
59	Nilotinib Vs. Dasatinib in Achieving MR4.5 for Newly Diagnosed Chronic Myeloid Leukemia: Results of the Prospective Randomized Phase 3 Study, JALSC CML212. <i>Blood</i> , 2020, 136, 40-41.	0.6	11
60	Phase 2 study of bosutinib in Japanese patients with newly diagnosed chronic phase chronic myeloid leukemia. <i>International Journal of Hematology</i> , 2020, 112, 24-32.	0.7	10
61	Lineage Involvement of Stem Cells Bearing the Philadelphia Chromosome in Chronic Myeloid Leukemia in the Chronic Phase as Shown by a Combination of Fluorescence-Activated Cell Sorting and Fluorescence In Situ Hybridization. <i>Blood</i> , 1998, 92, 4758-4763.	0.6	10
62	Effect of oral itraconazole on the pharmacokinetics of tacrolimus in a hematopoietic stem cell transplant recipient with CYP3A5*3/*3. <i>American Journal of Hematology</i> , 2010, 85, 634-635.	2.0	9
63	H2-receptor antagonist influences dasatinib pharmacokinetics in a patient with Philadelphia-positive acute lymphoblastic leukemia. <i>Cancer Chemotherapy and Pharmacology</i> , 2012, 70, 351-352.	1.1	9
64	Phase II Clinical Trial of Lenalidomide and Dexamethasone Therapy in Japanese Elderly Patients With Newly Diagnosed Multiple Myeloma to Determine Optimal Plasma Concentration of Lenalidomide. <i>Therapeutic Drug Monitoring</i> , 2018, 40, 301-309.	1.0	9
65	Safety and efficacy of high-dose ranimustine (MCNU) containing regimen followed by autologous stem cell transplantation for diffuse large B-cell lymphoma. <i>International Journal of Hematology</i> , 2018, 108, 510-515.	0.7	9
66	High-throughput sequencing of IgG B-cell receptors reveals frequent usage of the rearranged IGHV4*28/IGHJ4 gene in primary immune thrombocytopenia. <i>Scientific Reports</i> , 2019, 9, 8645.	1.6	9
67	Hypereosinophilic syndrome with abundant Charcot-Leyden crystals in spleen and lymph nodes. <i>Asia Pacific Allergy</i> , 2020, 10, e24.	0.6	9
68	Multiple myeloma with t(11;14)-associated immature phenotype has lower CD38 expression and higher BCL2 dependence. <i>Cancer Science</i> , 2021, 112, 3645-3654.	1.7	8
69	Effect of CYP3A5 and ABCB1 polymorphisms on the interaction between tacrolimus and itraconazole in patients with connective tissue disease. <i>European Journal of Clinical Pharmacology</i> , 2015, 71, 1091-1097.	0.8	7
70	Multiple Myeloma-Associated Ig Light Chain Crystalline Cast Nephropathy. <i>Kidney International Reports</i> , 2020, 5, 1595-1602.	0.4	7
71	The BCRP inhibitor febuxostat enhances the effect of nilotinib by regulation of intracellular concentration. <i>International Journal of Hematology</i> , 2021, 113, 100-105.	0.7	7
72	Safety and efficacy of low-dose liposomal amphotericin B as empirical antifungal therapy for patients with prolonged neutropenia. <i>International Journal of Clinical Oncology</i> , 2013, 18, 983-987.	1.0	6

#	ARTICLE	IF	CITATIONS
73	Long-term treatment with bosutinib in a phase 1/2 study in Japanese chronic myeloid leukemia patients resistant/intolerant to prior tyrosine kinase inhibitor treatment. <i>International Journal of Hematology</i> , 2017, 106, 398-410.	0.7	6
74	Effects of polymorphisms in NR112, CYP3A4, and ABC transporters on the steady-state plasma trough concentrations of bosutinib in Japanese patient with chronic myeloid leukemia. <i>Medical Oncology</i> , 2018, 35, 90.	1.2	6
75	Influence of ABCB1 polymorphisms on the pharmacokinetics and toxicity of lenalidomide in patients with multiple myeloma. <i>Medical Oncology</i> , 2019, 36, 55.	1.2	6
76	Treatment-free remission (TFR) in patients (pts) with chronic myeloid leukemia in chronic phase (CML-CP) treated with second-line nilotinib (NIL): First results from the ENESTop study.. <i>Journal of Clinical Oncology</i> , 2016, 34, 7054-7054.	0.8	6
77	ENESTop 192-week results: Treatment-free remission (TFR) in patients (pts) with chronic myeloid leukemia in chronic phase (CML-CP) after stopping second-line (2L) nilotinib (NIL).. <i>Journal of Clinical Oncology</i> , 2019, 37, 7005-7005.	0.8	6
78	Comparative proteomic analysis of renal proteins from IgA nephropathy model mice and control mice. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 666-679.	0.7	6
79	Downregulation of miR-26 promotes invasion and metastasis via targeting interleukin-22 in cutaneous T-cell lymphoma. <i>Cancer Science</i> , 2022, 113, 1208-1219.	1.7	6
80	Safety profile of bosutinib in Japanese versus non-Japanese patients with chronic myeloid leukemia: a pooled analysis. <i>International Journal of Hematology</i> , 2022, 115, 838-851.	0.7	6
81	A Limited Sampling Model to Estimate Exposure to Lenalidomide in Multiple Myeloma Patients. <i>Therapeutic Drug Monitoring</i> , 2014, 36, 505-509.	1.0	5
82	TAFRO Syndrome with Bilateral Adrenal Hemorrhage. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2017, 106, 288-294.	0.0	5
83	The potential role of clarithromycin addition to lenalidomide and dexamethasone therapy (BiRd) in multiple myeloma. <i>Annals of Hematology</i> , 2018, 97, 1097-1099.	0.8	4
84	Treatment outcomes of chronic-phase chronic myeloid leukemia with resistance and/or intolerance to a 1st-line tyrosine kinase inhibitor in Japan: the results of the New TARGET study 2nd-line. <i>International Journal of Hematology</i> , 2020, 111, 812-825.	0.7	4
85	Effects of SLC22A2 808G>T polymorphism and bosutinib concentrations on serum creatinine in patients with chronic myeloid leukemia receiving bosutinib therapy. <i>Scientific Reports</i> , 2021, 11, 6362.	1.6	4
86	Long-term treatment-free remission (TFR) in patients (pts) with chronic myeloid leukemia in chronic phase (CML-CP) after stopping second-line (2L) nilotinib: ENESTop 144-wk results.. <i>Journal of Clinical Oncology</i> , 2018, 36, 7003-7003.	0.8	4
87	Hematologic Malignancies (HM)-Screen-Japan 01: A Mutation Profiling Multicenter Study on Patients with Acute Myeloid Leukemia. <i>Blood</i> , 2021, 138, 4457-4457.	0.6	4
88	Phagocytosis of co-developing neutrophil progenitors by dendritic cells in a culture of human CD34+ cells with granulocyte colony-stimulating factor and tumor necrosis factor- α . <i>International Journal of Hematology</i> , 2008, 88, 64-72.	0.7	3
89	Switching to nilotinib is associated with deeper molecular responses in chronic myeloid leukemia chronic phase with major molecular responses to imatinib: STAT1 trial in Japan. <i>International Journal of Hematology</i> , 2018, 108, 176-183.	0.7	3
90	Effects of proprotein convertase subtilisin/kexin type 9 and nilotinib plasma concentrations on nilotinib-induced hypercholesterolaemia in patients with chronic myeloid leukaemia. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2021, 46, 382-387.	0.7	3

#	ARTICLE	IF	CITATIONS
91	The Genomic and Epigenomic Landscapes of Blast Crisis Transformation in Chronic Myeloid Leukemia. <i>Blood</i> , 2015, 126, 3737-3737.	0.6	3
92	Fluorescence In Situ Hybridization of Progenitor Cells Obtained by Fluorescence-Activated Cell Sorting for the Detection of Cells Affected by Chromosome Abnormality Trisomy 8 in Patients With Myelodysplastic Syndromes. <i>Blood</i> , 1998, 92, 2886-2892.	0.6	3
93	Regulatory T Cell as a Biomarker of Treatment-Free Remission in Patients with Chronic Myeloid Leukemia. <i>Cancers</i> , 2021, 13, 5904.	1.7	3
94	Bile acid is important for gastrointestinal absorption of nilotinib. <i>European Journal of Clinical Pharmacology</i> , 2012, 68, 1575-1576.	0.8	2
95	Effect of low platelet HLA-C expression on donor-specific antibody depletion following platelet transfusion from a corresponding HLA donor. <i>Bone Marrow Transplantation</i> , 2019, 54, 1713-1716.	1.3	2
96	Evaluation of the plasma concentration of ponatinib in a chronic myeloid leukaemia patient with ponatinib intolerance. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2021, 46, 219-222.	0.7	2
97	The Combination of Interferon-Alpha and Ponatinib Enables Faster and Deeper Molecular Responses in Patient with De Novo Blast Crisis of CML: Interferon-Alpha May Return as a CML Treatment. <i>Case Reports in Hematology</i> , 2021, 2021, 1-4.	0.3	2
98	Serial evaluation of the pharmacokinetics of ponatinib in patients with CML and Ph ⁺ ALL. <i>International Journal of Hematology</i> , 2021, 114, 509-516.	0.7	2
99	Dasatinib Cerebrospinal Fluid Concentration and Plasma Pharmacokinetics: Potential for Central Nervous System Prophylaxis In Philadelphia Chromosome-Positive Leukemia. <i>Blood</i> , 2010, 116, 1807-1807.	0.6	2
100	A multicenter phase II study of bendamustine with rituximab in patients with relapsed/refractory diffuse large B-cell lymphoma (DLBCL).. <i>Journal of Clinical Oncology</i> , 2012, 30, 8023-8023.	0.8	2
101	Safety, Feasibility and Efficacy of High Dose Ranimustine (MCNU), Carboplatin, Etoposide, and Cyclophosphamide (MCVC) Therapy Followed by Autologous Stem Cell Transplantation for Malignant Lymphoma.. <i>Blood</i> , 2010, 116, 4588-4588.	0.6	2
102	Interim Analysis of Hematologic Malignancies (HM)-Screen-Japan 01: A Mutation Profiling Multicenter Study of Patients with AML. <i>Blood</i> , 2020, 136, 2-3.	0.6	2
103	Relationship between achievement of major molecular response or deep molecular response and nilotinib plasma concentration in patients with chronic myeloid leukemia receiving first-line nilotinib therapy. <i>Cancer Chemotherapy and Pharmacology</i> , 2022, 89, 609-616.	1.1	2
104	Therapeutic drug monitoring enables safe and effective lenalidomide therapy in patients with multiple myeloma on hemodialysis. <i>Annals of Hematology</i> , 2016, 95, 2087-2088.	0.8	1
105	Drug interaction between tacrolimus and nilotinib in a patient with chronic myeloid leukemia after renal transplant. <i>Clinical Case Reports (discontinued)</i> , 2017, 5, 605-607.	0.2	1
106	Thrombocytopenia Caused by a Tea Beverage of <i>Taxus yunnanensis</i> (Chinese Yew). <i>Internal Medicine</i> , 2019, 58, 3153-3156.	0.3	1
107	RIZ1 Is Downregulated during CML Progression and Displays Tumor Suppressor Properties in CML Cell Lines.. <i>Blood</i> , 2006, 108, 2134-2134.	0.6	1
108	Clinical Significance of FLT3 Mutations in a Comprehensive NGS Multicenter Study of AML: HM-Screen-Japan 01. <i>Blood</i> , 2021, 138, 2313-2313.	0.6	1

#	ARTICLE	IF	CITATIONS
109	Early prediction of a long-term outcome by neutrophil-FISH in patients with CML receiving imatinib mesylate. <i>International Journal of Hematology</i> , 2010, 92, 559-561.	0.7	0
110	Hypereosinophilic Syndrome in the Tyrosine Kinase Inhibitor Era. <i>Internal Medicine</i> , 2015, 54, 551-552.	0.3	0
111	Clinical Analysis of Adult Acute Leukemia with Rearrangements of the 11q23/MLL: Multicenter Co-Operative Study.. <i>Blood</i> , 2006, 108, 2354-2354.	0.6	0
112	Phagocytosis of Co-Developing Neutrophil Progenitors by Dendritic Cells in Culture with Granulocyte-Colony Stimulating-Factor and Tumor Necrosis Factor- α : Induction of T Regulatory Cells by Co-Developing Dendritic Cells.. <i>Blood</i> , 2006, 108, 1720-1720.	0.6	0
113	Low Level of Serum Haptoglobin in Patients with Acquired Bone Marrow Failure (BMF) Syndromes.. <i>Blood</i> , 2007, 110, 3774-3774.	0.6	0
114	Low Dose and Standard Dose of Imatinib Therapy for Patients with Chronic Myeloid Leukemia in Akita Prefecture, Japan.. <i>Blood</i> , 2007, 110, 4575-4575.	0.6	0
115	Personalized Therapy of Tyrosine Kinase Inhibitors. <i>Japanese Journal of Clinical Pharmacology and Therapeutics</i> , 2013, 44, 225-228.	0.1	0
116	Effective Steroid Pulse Therapy for mitigate the acute phase symptoms of Human herpesvirus 6 encephalitis after allogenic hematopoietic stem cell transplantation: experience of two cases. <i>Journal of Hematopoietic Cell Transplantation</i> , 2013, 2, 75-79.	0.1	0
117	Ponatinib Safety and Efficacy in Japanese Patients with Philadelphia Positive Leukemia: Update of a Phase 1/2 Study. <i>Blood</i> , 2014, 124, 5541-5541.	0.6	0
118	Functional Analysis of the CML Blast Crisis Transcriptome and Epigenome Using Crispr-CAS9 and Pharmacologic Approaches. <i>Blood</i> , 2015, 126, 2764-2764.	0.6	0
119	7. Diagnosis and Treatment of Chronic Myeloid Leukemia and Myeloproliferative Neoplasms. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2019, 108, 547-550.	0.0	0
120	Properties and Distribution of IDH-1/2 Mutations in Acute Myeloid Leukemia By the Comprehensive Genomic Analysis. <i>Blood</i> , 2021, 138, 4447-4447.	0.6	0
121	Efficacy and Safety of Bosutinib in Japanese Patients with Newly Diagnosed Chronic Phase Chronic Myeloid Leukemia: Final 3-Year Results of a Phase 2 Study. <i>Blood</i> , 2021, 138, 2557-2557.	0.6	0
122	Genomic Analysis of <i>NPM1</i> Mutation and <i>KMT2A</i> / <i>MLL</i> -Rearrangement/Amplification in Japanese Patients with Acute Myeloid Leukemia: Hematologic Malignancies (HM)-Screen-Japan 01. <i>Blood</i> , 2021, 138, 4460-4460.	0.6	0
123	Patient and Physician Perspectives of Unmet Needs in CML - Designing the CML SUN Survey. <i>Blood</i> , 2021, 138, 4986-4986.	0.6	0
124	Genetic Features of AML with <i>MLL</i> -Rearrangement and <i>NPM1</i> Mutation: An Interim-Analysis of HM-Screen-Japan 01. <i>Blood</i> , 2020, 136, 35-36.	0.6	0
125	Genomic Analysis of <i>FLT3</i> Mutations in a Comprehensive NGS Multicenter Study of AML: HM-Screen-Japan 01. <i>Blood</i> , 2020, 136, 32-34.	0.6	0