

Gianpaolo Zerbini

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/565270/publications.pdf>

Version: 2024-02-01

16
papers

799
citations

759055

12
h-index

940416

16
g-index

18
all docs

18
docs citations

18
times ranked

1274
citing authors

#	ARTICLE	IF	CITATIONS
1	Renal dysfunction and podocyturia in pre-eclampsia may be explained by increased urinary VEGF. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, 1109-1117.	0.4	6
2	Sodium-glucose cotransporter 2 inhibition prevents renal fibrosis in cyclosporine nephropathy. <i>Acta Diabetologica</i> , 2021, 58, 1059-1070.	1.2	22
3	Angiotensin Type 2 and Mas Receptor Activation Prevents Myocardial Fibrosis and Hypertrophy through the Reduction of Inflammatory Cell Infiltration and Local Sympathetic Activity in Angiotensin II-Dependent Hypertension. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13678.	1.8	16
4	Association between On-Treatment Haemoglobin A1c and All-Cause Mortality in Individuals with Type 2 Diabetes: Importance of Personalized Goals and Type of Anti-Hyperglycaemic Treatment. <i>Journal of Clinical Medicine</i> , 2020, 9, 246.	1.0	2
5	Morphofunctional analysis of the retina in patients with type 1 diabetes without complications after 30 years of disease. <i>Scientific Reports</i> , 2020, 10, 206.	1.6	12
6	Multimodal Imaging Assessment of Vascular and Neurodegenerative Retinal Alterations in Type 1 Diabetic Patients without Fundoscopic Signs of Diabetic Retinopathy. <i>Journal of Clinical Medicine</i> , 2019, 8, 1409.	1.0	33
7	Activation of angiotensin type 2 (AT2) receptors prevents myocardial hypertrophy in Zucker diabetic fatty rats. <i>Acta Diabetologica</i> , 2019, 56, 97-104.	1.2	19
8	Increased soluble urokinase plasminogen activator receptor (suPAR) levels in neovascular age-related macular degeneration: a role for inflammation in the pathogenesis of the disease?. <i>Graefe's Archive for Clinical and Experimental Ophthalmology</i> , 2019, 257, 899-903.	1.0	2
9	Glomerular endothelial cells versus podocytes as the cellular target in diabetic nephropathy. <i>Acta Diabetologica</i> , 2018, 55, 1105-1111.	1.2	25
10	Optical coherence tomography angiography analysis of retinal vascular plexuses and choriocapillaris in patients with type 1 diabetes without diabetic retinopathy. <i>Acta Diabetologica</i> , 2017, 54, 695-702.	1.2	221
11	Angiotensin type-2 (AT-2)-receptor activation reduces renal fibrosis in cyclosporine nephropathy: evidence for blood pressure independent effect. <i>Bioscience Reports</i> , 2016, 36, .	1.1	15
12	Different regulation of miR-29a-3p in glomeruli and tubules in an experimental model of angiotensin II-dependent hypertension: potential role in renal fibrosis. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2016, 43, 335-342.	0.9	15
13	Viable Podocyturia in Healthy Individuals: Implications for Podocytopathies. <i>American Journal of Kidney Diseases</i> , 2014, 64, 1003-1005.	2.1	16
14	Rate and Determinants of Association Between Advanced Retinopathy and Chronic Kidney Disease in Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2317-2323.	4.3	106
15	Clinical significance of nonalbuminuric renal impairment in type 2 diabetes. <i>Journal of Hypertension</i> , 2011, 29, 1802-1809.	0.3	198
16	Persistent Renal Hypertrophy and Faster Decline of Glomerular Filtration Rate Precede the Development of Microalbuminuria in Type 1 Diabetes. <i>Diabetes</i> , 2006, 55, 2620-2625.	0.3	89