

# Pedro Romano

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5659378/publications.pdf>

Version: 2024-02-01

16  
papers

216  
citations

1307594

7  
h-index

1058476

14  
g-index

27  
all docs

27  
docs citations

27  
times ranked

270  
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Atolchelys lepida</i> , a new side-necked turtle from the Early Cretaceous of Brazil and the age of crown Pleurodira. <i>Biology Letters</i> , 2014, 10, 20140290.	2.3	48
2	Palynofacies as indicators of paleoenvironmental changes in a Cretaceous succession from the Larsen Basin, James Ross Island, Antarctica. <i>Sedimentary Geology</i> , 2013, 295, 53-66.	2.1	39
3	ARE EXTANT PODOCNEMIDID TURTLES RELICTS OF A WIDESPREAD CRETACEOUS ANCESTOR?. <i>South American Journal of Herpetology</i> , 2006, 1, 175-184.	0.5	28
4	New Information about Pelomedusoides (Testudines: Pleurodira) from the Cretaceous of Brazil. <i>Vertebrate Paleobiology and Paleoanthropology</i> , 2013, , 261-275.	0.5	16
5	The oldest plesiosaur (Reptilia, Sauropterygia) from Antarctica. <i>Polar Research</i> , 2011, 30, 7265.	1.6	13
6	Taxonomic identification using geometric morphometric approach and limited data: an example using the upper molars of two sympatric species of <i>Calomys</i> (Cricetidae: Rodentia). <i>Zoologia</i> , 2017, 34, 1-11.	0.5	10
7	Intra-specific variation and allometry of the skull of Late Cretaceous side-necked turtle <i>Bauruemys elegans</i> (Pleurodira, Podocnemididae) and how to deal with morphometric data in fossil vertebrates. <i>PeerJ</i> , 2017, 5, e2890.	2.0	7
8	The shape of sound: A new R package that crosses the bridge between Bioacoustics and Geometric Morphometrics. <i>Methods in Ecology and Evolution</i> , 2021, 12, 1115-1121.	5.2	7
9	Rostral morphology of Spinosauridae (Theropoda, Megalosauroidea): premaxilla shape variation and a new phylogenetic inference. <i>Historical Biology</i> , 2022, 34, 2089-2109.	1.4	7
10	Dental eruption sequence and hypsodonty index of a Pleistocene macraucheniid from the Brazilian Intertropical Region. <i>Journal of Paleontology</i> , 2017, 91, 1083-1090.	0.8	6
11	Biogeographic tools help to assess the effectiveness of protected areas for the conservation of anurans in the Mantiqueira mountain range, Southeastern Brazil. <i>Journal for Nature Conservation</i> , 2020, 54, 125799.	1.8	5
12	New outcrop with vertebrate remains from Solimões Formation (Eocene–Pliocene), Southern Solimões Basin, Acre State, Northern Brazil. <i>Journal of South American Earth Sciences</i> , 2020, 101, 102588.	1.4	4
13	The tale of the headless turtle. <i>Zootaxa</i> , 2016, 4200, 327.	0.5	3
14	Geochemical characterization of fossil turtles from Tartaruguito outcrop (Upper Cretaceous, Bauru) <i>Journal of South American Earth Sciences</i> , 2020, 104, 102843.	1.4	1
15	Species Delimitation of the Atlantic Forest Endemic <i>Inga subnuda</i> (Leguminosae,) <i>Palaeoecological Data. Systematic Botany</i> , 2022, 47, 417-430.	0.5	1
16	Georeferencing fossiliferous localities from Solimões and Acre Basins (Brazil) - what we know so far about Solimões Formation and future perspectives. <i>Anais Da Academia Brasileira De Ciencias</i> , 2021, 93, e20201642.	0.8	0