

Enrique Brandan S

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5659948/publications.pdf>

Version: 2024-02-01

120
papers

5,926
citations

41258

49
h-index

88477

70
g-index

122
all docs

122
docs citations

122
times ranked

5926
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Anchorage of collagen-tailed acetylcholinesterase to the extracellular matrix is mediated by heparan sulfate proteoglycans.. Journal of Cell Biology, 1985, 101, 985-992. | 2.3 | 156 |
| 2 | ECM is required for skeletal muscle differentiation independently of muscle regulatory factor expression. American Journal of Physiology - Cell Physiology, 2002, 282, C383-C394. | 2.1 | 144 |
| 3 | Restoration of muscle strength in dystrophic muscle by angiotensin-1-7 through inhibition of TGF- β 2 signalling. Human Molecular Genetics, 2014, 23, 1237-1249. | 1.4 | 143 |
| 4 | Decorin Core Protein Fragment Leu155-Val260 Interacts with TGF- β 2 but Does Not Compete for Decorin Binding to Type I Collagen. Archives of Biochemistry and Biophysics, 1998, 355, 241-248. | 1.4 | 138 |
| 5 | Extracellular proteoglycans modify TGF- β 2 bio-availability attenuating its signaling during skeletal muscle differentiation. Matrix Biology, 2006, 25, 332-341. | 1.5 | 127 |
| 6 | CTGF Inhibits BMP-7 Signaling in Diabetic Nephropathy. Journal of the American Society of Nephrology: JASN, 2008, 19, 2098-2107. | 3.0 | 123 |
| 7 | Mice Long-Term High-Fat Diet Feeding Recapitulates Human Cardiovascular Alterations: An Animal Model to Study the Early Phases of Diabetic Cardiomyopathy. PLoS ONE, 2013, 8, e60931. | 1.1 | 121 |
| 8 | Reducing CTGF/CCN2 slows down mdx muscle dystrophy and improves cell therapy. Human Molecular Genetics, 2013, 22, 4938-4951. | 1.4 | 118 |
| 9 | Connective tissue cells expressing fibro/adipogenic progenitor markers increase under chronic damage: relevance in fibroblast-myofibroblast differentiation and skeletal muscle fibrosis. Cell and Tissue Research, 2016, 364, 647-660. | 1.5 | 117 |
| 10 | Heparan sulfate proteoglycans are increased during skeletal muscle regeneration: requirement of syndecan-3 for successful fiber formation. Journal of Cell Science, 2004, 117, 73-84. | 1.2 | 112 |
| 11 | A Novel Modulatory Mechanism of Transforming Growth Factor- β 2 Signaling through Decorin and LRP-1. Journal of Biological Chemistry, 2007, 282, 18842-18850. | 1.6 | 112 |
| 12 | Skeletal muscle cells express the profibrotic cytokine connective tissue growth factor (CTGF/CCN2), which induces their dedifferentiation. Journal of Cellular Physiology, 2008, 215, 410-421. | 2.0 | 109 |
| 13 | Wnt Signaling in Skeletal Muscle Dynamics: Myogenesis, Neuromuscular Synapse and Fibrosis. Molecular Neurobiology, 2014, 49, 574-589. | 1.9 | 107 |
| 14 | Biglycan is a new extracellular component of the Chordin-BMP4 signaling pathway. EMBO Journal, 2005, 24, 1397-1405. | 3.5 | 104 |
| 15 | Extracellular matrix is required for skeletal muscle differentiation but not myogenin expression. , 1996, 62, 227-239. | | 103 |
| 16 | Decorin Interacts with Connective Tissue Growth Factor (CTGF)/CCN2 by LRR12 Inhibiting Its Biological Activity. Journal of Biological Chemistry, 2011, 286, 24242-24252. | 1.6 | 101 |
| 17 | A Novel Mechanism of Sequestering Fibroblast Growth Factor 2 by Glypican in Lipid Rafts, Allowing Skeletal Muscle Differentiation. Molecular and Cellular Biology, 2010, 30, 1634-1649. | 1.1 | 100 |
| 18 | ALS skeletal muscle shows enhanced TGF- β 2 signaling, fibrosis and induction of fibro/adipogenic progenitor markers. PLoS ONE, 2017, 12, e0177649. | 1.1 | 94 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Antisense Inhibition of Decorin Expression in Myoblasts Decreases Cell Responsiveness to Transforming Growth Factor β^2 and Accelerates Skeletal Muscle Differentiation. <i>Journal of Biological Chemistry</i> , 2001, 276, 3589-3596. | 1.6 | 93 |
| 20 | Transient up-regulation of biglycan during skeletal muscle regeneration: delayed fiber growth along with decorin increase in biglycan-deficient mice. <i>Developmental Biology</i> , 2004, 268, 358-371. | 0.9 | 92 |
| 21 | CTGF/CCN2 overexpression can directly induce features of skeletal muscle dystrophy. <i>Journal of Pathology</i> , 2011, 225, 490-501. | 2.1 | 92 |
| 22 | Increase in decorin and biglycan in Duchenne Muscular Dystrophy: role of fibroblasts as cell source of these proteoglycans in the disease. <i>Journal of Cellular and Molecular Medicine</i> , 2006, 10, 758-769. | 1.6 | 89 |
| 23 | Angiotensins as therapeutic targets beyond heart disease. <i>Trends in Pharmacological Sciences</i> , 2015, 36, 310-320. | 4.0 | 85 |
| 24 | Novel regulatory mechanisms for the proteoglycans decorin and biglycan during muscle formation and muscular dystrophy. <i>Matrix Biology</i> , 2008, 27, 700-708. | 1.5 | 83 |
| 25 | Novel and optimized strategies for inducing fibrosis in vivo: focus on Duchenne Muscular Dystrophy. <i>Skeletal Muscle</i> , 2014, 4, 7. | 1.9 | 80 |
| 26 | Antisense Inhibition of Syndecan-3 Expression during Skeletal Muscle Differentiation Accelerates Myogenesis through a Basic Fibroblast Growth Factor-dependent Mechanism. <i>Journal of Biological Chemistry</i> , 1999, 274, 37876-37884. | 1.6 | 73 |
| 27 | Syndecan-1 Expression Is Down-regulated during Myoblast Terminal Differentiation. <i>Journal of Biological Chemistry</i> , 1997, 272, 18418-18424. | 1.6 | 72 |
| 28 | Angiotensin II receptor type 1 blockade decreases CTGF/CCN2-mediated damage and fibrosis in normal and dystrophic skeletal muscles. <i>Journal of Cellular and Molecular Medicine</i> , 2012, 16, 752-764. | 1.6 | 72 |
| 29 | Angiotensin II-induced pro-fibrotic effects require p38MAPK activity and transforming growth factor beta 1 expression in skeletal muscle cells. <i>International Journal of Biochemistry and Cell Biology</i> , 2012, 44, 1993-2002. | 1.2 | 70 |
| 30 | Angiotensin-(1-7) decreases skeletal muscle atrophy induced by angiotensin II through a Mas receptor-dependent mechanism. <i>Clinical Science</i> , 2015, 128, 307-319. | 1.8 | 70 |
| 31 | The cross-talk between TGF- β^2 and PDGFR signaling pathways regulates stromal fibro/adipogenic progenitors' fate. <i>Journal of Cell Science</i> , 2019, 132, . | 1.2 | 70 |
| 32 | Orientation and role of nucleoside diphosphatase and 5'-nucleotidase in Golgi vesicles from rat liver. <i>Biochemistry</i> , 1982, 21, 4640-4645. | 1.2 | 69 |
| 33 | Andrographolide Ameliorates Inflammation and Fibrogenesis and Attenuates Inflammasome Activation in Experimental Non-Alcoholic Steatohepatitis. <i>Scientific Reports</i> , 2017, 7, 3491. | 1.6 | 68 |
| 34 | Inhibition of the angiotensin-converting enzyme decreases skeletal muscle fibrosis in dystrophic mice by a diminution in the expression and activity of connective tissue growth factor (CTGF/CCN-2). <i>Cell and Tissue Research</i> , 2013, 353, 173-187. | 1.5 | 67 |
| 35 | Fibrotic response induced by angiotensin-II requires NAD(P)H oxidase-induced reactive oxygen species (ROS) in skeletal muscle cells. <i>Biochemical and Biophysical Research Communications</i> , 2011, 410, 665-670. | 1.0 | 65 |
| 36 | Angiotensin-(1-7) attenuates disuse skeletal muscle atrophy via the Mas receptor. <i>DMM Disease Models and Mechanisms</i> , 2016, 9, 441-9. | 1.2 | 65 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Synthesis of proteoglycans is augmented in dystrophic mdx mouse skeletal muscle. <i>European Journal of Cell Biology</i> , 2000, 79, 173-181. | 1.6 | 63 |
| 38 | Role of skeletal muscle proteoglycans during myogenesis. <i>Matrix Biology</i> , 2013, 32, 289-297. | 1.5 | 63 |
| 39 | Syndecan-1 Expression Inhibits Myoblast Differentiation through a Basic Fibroblast Growth Factor-dependent Mechanism. <i>Journal of Biological Chemistry</i> , 1998, 273, 32288-32296. | 1.6 | 62 |
| 40 | The Internal Region Leucine-rich Repeat 6 of Decorin Interacts with Low Density Lipoprotein Receptor-related Protein-1, Modulates Transforming Growth Factor (TGF)- β 2-dependent Signaling, and Inhibits TGF- β 2-dependent Fibrotic Response in Skeletal Muscles. <i>Journal of Biological Chemistry</i> , 2012, 287, 6773-6787. | 1.6 | 60 |
| 41 | Axonal sprouting induced in the sciatic nerve by the amyloid precursor protein (APP) and other antiproteases. <i>Neuroscience Letters</i> , 1992, 144, 130-134. | 1.0 | 58 |
| 42 | Endotoxin-induced skeletal muscle wasting is prevented by angiotensin-(1 α -7) through a p38 MAPK-dependent mechanism. <i>Clinical Science</i> , 2015, 129, 461-476. | 1.8 | 57 |
| 43 | Expression of Perlecan, a Proteoglycan That Binds Myogenic Inhibitory Basic Fibroblast Growth Factor, Is Down Regulated during Skeletal Muscle Differentiation. <i>Experimental Cell Research</i> , 1997, 234, 405-412. | 1.2 | 56 |
| 44 | The formation of skeletal muscle myotubes requires functional membrane receptors activated by extracellular ATP. <i>Brain Research Reviews</i> , 2004, 47, 174-188. | 9.1 | 56 |
| 45 | The angiotensin-(1 α -7)/Mas axis reduces myonuclear apoptosis during recovery from angiotensin II-induced skeletal muscle atrophy in mice. <i>Pflugers Archiv European Journal of Physiology</i> , 2015, 467, 1975-1984. | 1.3 | 53 |
| 46 | Augmented synthesis and differential localization of heparan sulfate proteoglycans in Duchenne muscular dystrophy. <i>Journal of Cellular Biochemistry</i> , 2002, 85, 703-713. | 1.2 | 52 |
| 47 | Dermatan sulfate exerts an enhanced growth factor response on skeletal muscle satellite cell proliferation and migration. <i>Journal of Cellular Physiology</i> , 2004, 198, 169-178. | 2.0 | 52 |
| 48 | Denervation-induced skeletal muscle fibrosis is mediated by CTGF/CCN2 independently of TGF- β 2. <i>Matrix Biology</i> , 2019, 82, 20-37. | 1.5 | 52 |
| 49 | ACE2 Is Augmented in Dystrophic Skeletal Muscle and Plays a Role in Decreasing Associated Fibrosis. <i>PLoS ONE</i> , 2014, 9, e93449. | 1.1 | 51 |
| 50 | The Low Density Lipoprotein Receptor-related Protein Functions as an Endocytic Receptor for Decorin. <i>Journal of Biological Chemistry</i> , 2006, 281, 31562-31571. | 1.6 | 50 |
| 51 | Connective tissue growth factor induction by lysophosphatidic acid requires transactivation of transforming growth factor type β 2 receptors and the JNK pathway. <i>Cellular Signalling</i> , 2011, 23, 449-457. | 1.7 | 50 |
| 52 | Extracellular matrix histone H1 binds to perlecan, is present in regenerating skeletal muscle and stimulates myoblast proliferation. <i>Journal of Cell Science</i> , 2002, 115, 2041-2051. | 1.2 | 50 |
| 53 | Betaglycan induces TGF- β 2 signaling in a ligand-independent manner, through activation of the p38 pathway. <i>Cellular Signalling</i> , 2006, 18, 1482-1491. | 1.7 | 49 |
| 54 | Isolation of the heparan sulfate proteoglycans from the extracellular matrix of rat skeletal muscle. <i>Journal of Neurobiology</i> , 1987, 18, 271-282. | 3.7 | 46 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Caenorhabditis elegans syndecan (SDN-1) is required for normal egg laying and associates with the nervous system and the vulva. <i>Journal of Cell Science</i> , 2004, 117, 5179-5190. | 1.2 | 46 |
| 56 | TGF- β 2 receptors, in a Smad-independent manner, are required for terminal skeletal muscle differentiation. <i>Experimental Cell Research</i> , 2010, 316, 2487-2503. | 1.2 | 45 |
| 57 | Role of hypoxia in skeletal muscle fibrosis: Synergism between hypoxia and TGF- β 2 signaling upregulates CCN2/CTGF expression specifically in muscle fibers. <i>Matrix Biology</i> , 2020, 87, 48-65. | 1.5 | 45 |
| 58 | Changes in secreted and cell associated proteoglycan synthesis during conversion of myoblasts to osteoblasts in response to bone morphogenetic protein-2: Role of decorin in cell response to BMP-2. <i>Journal of Cellular Physiology</i> , 2006, 206, 58-67. | 2.0 | 44 |
| 59 | Betaglycan Expression Is Transcriptionally Up-regulated during Skeletal Muscle Differentiation. <i>Journal of Biological Chemistry</i> , 2003, 278, 382-390. | 1.6 | 43 |
| 60 | Interaction between Alzheimer's disease β 44 precursor protein (APP) and the extracellular matrix: Evidence for the participation of heparan sulfate proteoglycans. <i>Journal of Cellular Biochemistry</i> , 1997, 65, 145-158. | 1.2 | 42 |
| 61 | Diet-Induced Nonalcoholic Fatty Liver Disease Is Associated with Sarcopenia and Decreased Serum Insulin-Like Growth Factor-1. <i>Digestive Diseases and Sciences</i> , 2016, 61, 3190-3198. | 1.1 | 42 |
| 62 | Extracellular matrix histone H1 binds to perlecan, is present in regenerating skeletal muscle and stimulates myoblast proliferation. <i>Journal of Cell Science</i> , 2002, 115, 2041-51. | 1.2 | 42 |
| 63 | Structural and functional organization of synaptic acetylcholinesterase. <i>Brain Research Reviews</i> , 2004, 47, 96-104. | 9.1 | 41 |
| 64 | Expression and localization of proteoglycans during limb myogenic activation. <i>Developmental Dynamics</i> , 2001, 221, 106-115. | 0.8 | 40 |
| 65 | Constitutively activated dystrophic muscle fibroblasts show a paradoxical response to TGF- β 2 and CTGF/CCN2. <i>Journal of Cell Communication and Signaling</i> , 2007, 1, 205-217. | 1.8 | 40 |
| 66 | The pro-fibrotic connective tissue growth factor (CTGF/CCN2) correlates with the number of necrotic-regenerative foci in dystrophic muscle. <i>Journal of Cell Communication and Signaling</i> , 2018, 12, 413-421. | 1.8 | 40 |
| 67 | Adherent muscle connective tissue fibroblasts are phenotypically and biochemically equivalent to stromal fibro/adipogenic progenitors. <i>Matrix Biology Plus</i> , 2019, 2, 100006. | 1.9 | 37 |
| 68 | SUBCELLULAR FRACTIONATION STUDIES ON THE ORGANIZATION OF FATTY ACID OXIDATION BY LIVER PEROXISOMES. <i>Annals of the New York Academy of Sciences</i> , 1982, 386, 62-80. | 1.8 | 35 |
| 69 | A lipid-anchored heparan sulfate proteoglycan is present in the surface of differentiated skeletal muscle cells. Isolation and biochemical characterization. <i>FEBS Journal</i> , 1993, 216, 587-595. | 0.2 | 35 |
| 70 | Role of proteoglycans in the regulation of the skeletal muscle fibrotic response. <i>FEBS Journal</i> , 2013, 280, 4109-4117. | 2.2 | 35 |
| 71 | Heparin activates Wnt signaling for neuronal morphogenesis. <i>Journal of Cellular Physiology</i> , 2008, 216, 805-815. | 2.0 | 34 |
| 72 | HIF-hypoxia signaling in skeletal muscle physiology and fibrosis. <i>Journal of Cell Communication and Signaling</i> , 2020, 14, 147-158. | 1.8 | 34 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Andrographolide attenuates skeletal muscle dystrophy in mdx mice and increases efficiency of cell therapy by reducing fibrosis. <i>Skeletal Muscle</i> , 2014, 4, 6. | 1.9 | 33 |
| 74 | Extracellular matrix components and amyloid in neuritic plaques of Alzheimer's disease. <i>General Pharmacology</i> , 1993, 24, 1063-1068. | 0.7 | 31 |
| 75 | Matrix Metalloproteinase-2-deficient Fibroblasts Exhibit an Alteration in the Fibrotic Response to Connective Tissue Growth Factor/CCN2 because of an Increase in the Levels of Endogenous Fibronectin. <i>Journal of Biological Chemistry</i> , 2009, 284, 13551-13561. | 1.6 | 30 |
| 76 | Inhibition of myoblast migration via decorin expression is critical for normal skeletal muscle differentiation. <i>Developmental Biology</i> , 2003, 259, 209-224. | 0.9 | 29 |
| 77 | Transforming growth factor type beta 1 increases the expression of angiotensin II receptor type 2 by a SMAD β and p38 MAPK β dependent mechanism in skeletal muscle. <i>BioFactors</i> , 2013, 39, 467-475. | 2.6 | 29 |
| 78 | The inhibition of CTGF/CCN2 activity improves muscle and locomotor function in a murine ALS model. <i>Human Molecular Genetics</i> , 2018, 27, 2913-2926. | 1.4 | 29 |
| 79 | Glypican-1 regulates myoblast response to HGF via Met in a lipid raft-dependent mechanism: effect on migration of skeletal muscle precursor cells. <i>Skeletal Muscle</i> , 2014, 4, 5. | 1.9 | 28 |
| 80 | Nilotinib impairs skeletal myogenesis by increasing myoblast proliferation. <i>Skeletal Muscle</i> , 2018, 8, 5. | 1.9 | 28 |
| 81 | Expression of CTGF/CCN2 in response to LPA is stimulated by fibrotic extracellular matrix via the integrin/FAK axis. <i>American Journal of Physiology - Cell Physiology</i> , 2018, 314, C415-C427. | 2.1 | 28 |
| 82 | CTGF/CCN2 from Skeletal Muscle to Nervous System: Impact on Neurodegenerative Diseases. <i>Molecular Neurobiology</i> , 2019, 56, 5911-5916. | 1.9 | 27 |
| 83 | TGF β -driven downregulation of the Wnt/ β -Catenin transcription factor TCF7L2/TCF4 in PDGFR β fibroblasts. <i>Journal of Cell Science</i> , 2020, 133, . | 1.2 | 26 |
| 84 | Transforming Growth Factor β (TGF β) Signaling Is Regulated by Electrical Activity in Skeletal Muscle Cells. <i>Journal of Biological Chemistry</i> , 2006, 281, 18473-18481. | 1.6 | 25 |
| 85 | PDGF-PDGFR network differentially regulates the fate, migration, proliferation, and cell cycle progression of myogenic cells. <i>Cellular Signalling</i> , 2021, 84, 110036. | 1.7 | 24 |
| 86 | Co-solubilization of asymmetric acetylcholinesterase and dermatan sulfate proteoglycan from the extracellular matrix of rat skeletal muscles. <i>FEBS Letters</i> , 1987, 213, 159-163. | 1.3 | 23 |
| 87 | Different membrane-bound forms of acetylcholinesterase are present at the cell surface of hepatocytes. <i>FEBS Journal</i> , 1989, 182, 203-207. | 0.2 | 23 |
| 88 | Isolation and purification of human biliary vesicles with potent cholesterol-nucleation-promoting activity. <i>Clinical Science</i> , 1992, 82, 175-180. | 1.8 | 23 |
| 89 | Isolation and partial characterization of cholesterol pronucleating hydrophobic glycoproteins associated to native biliary vesicles. <i>FEBS Letters</i> , 1993, 318, 45-49. | 1.3 | 23 |
| 90 | SMAD3 and SP1/SP3 Transcription Factors Collaborate to Regulate Connective Tissue Growth Factor Gene Expression in Myoblasts in Response to Transforming Growth Factor β . <i>Journal of Cellular Biochemistry</i> , 2015, 116, 1880-1887. | 1.2 | 22 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Interaction of skeletal muscle cells with collagen type IV is mediated by perlecan associated with the cell surface. , 1999, 75, 665-674. | | 19 |
| 92 | Driving fibrosis in neuromuscular diseases: Role and regulation of Connective tissue growth factor (CCN2/CTGF). Matrix Biology Plus, 2021, 11, 100059. | 1.9 | 18 |
| 93 | Blockade of Bradykinin receptors worsens the dystrophic phenotype of mdx mice: differential effects for B1 and B2 receptors. Journal of Cell Communication and Signaling, 2018, 12, 589-601. | 1.8 | 17 |
| 94 | The linkage between inflammation and fibrosis in muscular dystrophies: The axis autotaxinâ€“lysophosphatidic acid as a new therapeutic target?. Journal of Cell Communication and Signaling, 2021, 15, 317-334. | 1.8 | 15 |
| 95 | Adenovirus-mediated hepatic syndecan-1 overexpression induces hepatocyte proliferation and hyperlipidaemia in mice. Liver International, 2007, 27, 569-581. | 1.9 | 13 |
| 96 | RECK-Mediated Î²1-Integrin Regulation by TGF-Î²1 Is Critical for Wound Contraction in Mice. PLoS ONE, 2015, 10, e0135005. | 1.1 | 13 |
| 97 | Wnt signaling pathway improves central inhibitory synaptic transmission in a mouse model of Duchenne muscular dystrophy. Neurobiology of Disease, 2016, 86, 109-120. | 2.1 | 11 |
| 98 | The Low Density Lipoprotein Receptor-related Protein Functions as an Endocytic Receptor for Decorin. Journal of Biological Chemistry, 2006, 281, 31562-31571. | 1.6 | 11 |
| 99 | Isolation and characterization of rat skeletal muscle proteoglycan decorin and comparison with the human fibroblast decorin. Comparative Biochemistry and Physiology Part B: Comparative Biochemistry, 1991, 100, 565-570. | 0.2 | 10 |
| 100 | Inhibition of extracellular matrix assembly induces the expression of osteogenic markers in skeletal muscle cells by a BMP-2 independent mechanism. BMC Cell Biology, 2009, 10, 73. | 3.0 | 10 |
| 101 | Syndecan-4 and Î²1 integrin are regulated by electrical activity in skeletal muscle: Implications for cell adhesion. Matrix Biology, 2010, 29, 383-392. | 1.5 | 10 |
| 102 | Transforming growth factor typeâ€“Î² inhibits Mas receptor expression in fibroblasts but not in myoblasts or differentiated myotubes; Relevance to fibrosis associated to muscular dystrophies. BioFactors, 2015, 41, 111-120. | 2.6 | 9 |
| 103 | Role of Matricellular CCN Proteins in Skeletal Muscle: Focus on CCN2/CTGF and Its Regulation by Vasoactive Peptides. International Journal of Molecular Sciences, 2021, 22, 5234. | 1.8 | 9 |
| 104 | Sulfation is required for bone morphogenetic protein 2-dependent Id1 induction. Biochemical and Biophysical Research Communications, 2006, 344, 1207-1215. | 1.0 | 8 |
| 105 | Angiotensin-(1-7) Prevents Lipopolysaccharide-Induced Autophagy via the Mas Receptor in Skeletal Muscle. International Journal of Molecular Sciences, 2020, 21, 9344. | 1.8 | 8 |
| 106 | Activation of the ATX/LPA/LPARs axis induces a fibrotic response in skeletal muscle. Matrix Biology, 2022, 109, 121-139. | 1.5 | 8 |
| 107 | [28] Isolation and characterization of coated vesicles from rat liver. Methods in Enzymology, 1983, 98, 326-336. | 0.4 | 7 |
| 108 | Decorin is specifically solubilized by heparin from the extracellular matrix of rat skeletal muscles. FEBS Letters, 1993, 319, 249-252. | 1.3 | 6 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Golgi Complex Function in the Excretion of Renal Kallikrein. <i>Experimental Biology and Medicine</i> , 1982, 171, 221-231. | 1.1 | 5 |
| 110 | Sulfation is required for mobility of veliger larvae of <i>Concholepas concholepas</i> (Mollusca; Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 702 Td (| 1.4 | 5 |
| 111 | A high molecular weight proteoglycan is differentially expressed during development of the mollusc <i>Concholepas concholepas</i> (Mollusca; Gastropoda; Muricidae). <i>The Journal of Experimental Zoology</i> , 1992, 264, 363-371. | 1.4 | 5 |
| 112 | Fibro/adipogenic progenitors safeguard themselves: a novel mechanism to reduce fibrosis is discovered. <i>Journal of Cell Communication and Signaling</i> , 2017, 11, 77-78. | 1.8 | 5 |
| 113 | Analysis of Pathological Activities of CCN2/CTGF in Muscle Dystrophy. <i>Methods in Molecular Biology</i> , 2017, 1489, 513-521. | 0.4 | 5 |
| 114 | Isolation of proteoglycans synthesized by rat heart: Evidence for the presence of several distinct forms. <i>General Pharmacology</i> , 1992, 23, 249-255. | 0.7 | 3 |
| 115 | Reduced RECK levels accelerate skeletal muscle differentiation, improve muscle regeneration, and decrease fibrosis. <i>FASEB Journal</i> , 2021, 35, e21503. | 0.2 | 3 |
| 116 | Uptake of Tritiated Liquids by Individual Breakfast Cereal Flakes. <i>Journal of Food Science</i> , 2010, 75, E194-200. | 1.5 | 2 |
| 117 | Skeletal Muscle System. , 2019, , 169-190. | | 2 |
| 118 | Effect of salt concentration on the synthesis of sulphated macromolecules in the brine shrimp (<i>Artemia franciscana</i>): Changes of sulphation rate during development. <i>Comparative Biochemistry and Physiology A, Comparative Physiology</i> , 1993, 105, 519-523. | 0.7 | 1 |
| 119 | Heparan sulfate provides a mechanism to respond to FGFR2b and control regenerative expansion. <i>Journal of Cell Communication and Signaling</i> , 2015, 9, 89-89. | 1.8 | 1 |
| 120 | Increase of macromolecule synthesis after hatching of <i>Concholepas concholepas</i> veliger larvae: Effect of sulfate in the synthesis of proteoglycans. <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1990, 96, 613-619. | 0.2 | 0 |