

Stephan Stilgenbauer

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5664937/stephan-stilgenbauer-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

602

papers

30,596

citations

85

h-index

167

g-index

639

ext. papers

35,081

ext. citations

5.1

avg, IF

6.57

L-index

#	Paper	IF	Citations
602	KIR2DS1-HLA-C status as a predictive marker for benefit from rituximab: a post-hoc analysis of the RICOVER-60 and CLL8 trials.. <i>Lancet Haematology</i> , 2022 , 9, e133-e142	14.6	1
601	Efficacy and Safety of the Combination of Tirabrutinib and Entospletinib With or Without Obinutuzumab in Relapsed Chronic Lymphocytic Leukemia.. <i>HemaSphere</i> , 2022 , 6, e692	0.3	1
600	Efficacy and Safety of Tirabrutinib and Idelalisib With or Without Obinutuzumab in Relapsed Chronic Lymphocytic Leukemia. <i>HemaSphere</i> , 2022 , 6, e729	0.3	
599	Interleukin-10 receptor signaling promotes the maintenance of a PD-1 TCF-1 CD8 T cell population that sustains anti-tumor immunity. <i>Immunity</i> , 2021 ,	32.3	8
598	Activation of Notch and Myc signaling via B cell-restricted depletion of Dnmt3a generates a consistent murine model of chronic lymphocytic leukemia. <i>Cancer Research</i> , 2021 ,	10.1	1
597	IgG seroprevalence of COVID-19 among people living with HIV or at high risk of HIV in south-west Germany: A seroprevalence study. <i>HIV Medicine</i> , 2021 ,	2.7	1
596	Pooled Analysis of First-Line Treatment with Targeted Agents in Patients with Chronic Lymphocytic Leukemia (CLL) Aged 80 Years and Older. <i>Blood</i> , 2021 , 138, 1552-1552	2.2	1
595	Comparison of Tumor Lysis Syndrome (TLS) Risk Reduction and Incidence in Different Venetoclax-Based Combinations within the Randomized Phase 3 GAIA (CLL13) Trial. <i>Blood</i> , 2021 , 138, 2639-2639	2.2	
594	Characterization of Bruton Tyrosine Kinase Inhibitor (BTKi)-Related Adverse Events in a Head-to-Head Trial of Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2021 , 138, 3721-3721	2.2	
593	The CLL12 trial: Ibrutinib versus placebo in treatment-naïve, early stage chronic lymphocytic leukemia. <i>Blood</i> , 2021 ,	2.2	5
592	Identification of recurrent genomic alterations in the apoptotic machinery in chronic lymphocytic leukemia patients treated with venetoclax monotherapy. <i>American Journal of Hematology</i> , 2021 ,	7.1	1
591	High Resolution Assessment of Minimal Residual Disease (MRD) By Next-Generation Sequencing (NGS) and High-Sensitivity Flow Cytometry (hsFCM) in the Phase 3 GAIA (CLL13) Trial. <i>Blood</i> , 2021 , 138, 72-72	2.2	
590	A Randomized Phase III Study of Venetoclax-Based Time-Limited Combination Treatments (RVE, GVe, GIVe) Vs Standard Chemoimmunotherapy (CIT: FCR/BR) in Frontline Chronic Lymphocytic Leukemia (CLL) of Fit Patients: First Co-Primary Endpoint Analysis of the International Intergroup GAIA (CLL13) Trial. <i>Blood</i> , 2021 , 138, 71-71	2.2	5
589	Venetoclax plus bendamustine-rituximab or bendamustine-obinutuzumab in chronic lymphocytic leukemia: final results of a phase 1b study (GO28440). <i>Haematologica</i> , 2021 , 106, 2834-2844	6.6	1
588	Significant reduced loss of bone mineral density after four vs. six cycles of R-CHOP: an analysis of the FLYER-trial. <i>Leukemia and Lymphoma</i> , 2021 , 1-9	1.9	
587	Minimal Residual Disease Dynamics after Venetoclax-Obinutuzumab Treatment: Extended Off-Treatment Follow-up From the Randomized CLL14 Study. <i>Journal of Clinical Oncology</i> , 2021 , JCO2101181	2.2	10
586	Identification of the atypically modified autoantigen Ars2 as the target of B-cell receptors from activated B-cell-type diffuse large B-cell lymphoma. <i>Haematologica</i> , 2021 , 106, 2224-2232	6.6	6

585	Integrative prognostic models predict long-term survival after immunochemotherapy in chronic lymphocytic leukemia patients. <i>Haematologica</i> , 2021 ,	6.6	1
584	EOMES is essential for antitumor activity of CD8 T cells in chronic lymphocytic leukemia. <i>Leukemia</i> , 2021 , 35, 3152-3162	10.7	4
583	MARCKS affects cell motility and response to BTK inhibitors in CLL. <i>Blood</i> , 2021 , 138, 544-556	2.2	2
582	Clonal evolution in chronic lymphocytic leukemia is scant in relapsed but accelerated in refractory cases after chemo(immune)therapy. <i>Haematologica</i> , 2021 ,	6.6	3
581	Increased B-cell activity with consumption of activated monocytes in severe COVID-19 patients. <i>European Journal of Immunology</i> , 2021 , 51, 1449-1460	6.1	3
580	B-cell acute lymphoblastic leukemia in patients with chronic lymphocytic leukemia treated with lenalidomide. <i>Blood</i> , 2021 , 137, 2267-2271	2.2	3
579	FDG PET/CT to detect bone marrow involvement in the initial staging of patients with aggressive non-Hodgkin lymphoma: results from the prospective, multicenter PETAL and OPTIMAL>60 trials. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2021 , 48, 3550-3559	8.8	4
578	Discovery of Candidate DNA Methylation Cancer Driver Genes. <i>Cancer Discovery</i> , 2021 , 11, 2266-2281	24.4	12
577	Current Treatment Options in CLL. <i>Cancers</i> , 2021 , 13,	6.6	4
576	Mutational mechanisms shaping the coding and noncoding genome of germinal center derived B-cell lymphomas. <i>Leukemia</i> , 2021 , 35, 2002-2016	10.7	3
575	Clinical, biological, and molecular genetic features of Richter syndrome and prognostic significance: A study of the French Innovative Leukemia Organization. <i>American Journal of Hematology</i> , 2021 , 96, E311-E314 ²	7.1	1
574	Durable remissions following combined targeted therapy in patients with CLL harboring TP53 deletions and/or mutations. <i>Blood</i> , 2021 , 138, 1805-1816	2.2	1
573	Measurable residual disease in chronic lymphocytic leukemia: expert review and consensus recommendations. <i>Leukemia</i> , 2021 , 35, 3059-3072	10.7	6
572	Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic Leukemia: Results of the First Randomized Phase III Trial. <i>Journal of Clinical Oncology</i> , 2021 , 39, 3441-3452	2.2	65
571	Comparative analysis of targeted next-generation sequencing panels for the detection of gene mutations in chronic lymphocytic leukemia: an ERIC multi-center study. <i>Haematologica</i> , 2021 , 106, 682-691	6.6	3
570	Combining ibrutinib and checkpoint blockade improves CD8+ T-cell function and control of chronic lymphocytic leukemia in Em-TCL1 mice. <i>Haematologica</i> , 2021 , 106, 968-977	6.6	16
569	Clinical activity of abemaciclib in patients with relapsed or refractory mantle cell lymphoma - a phase II study. <i>Haematologica</i> , 2021 , 106, 859-862	6.6	4
568	Bendamustine, followed by ofatumumab and ibrutinib in chronic lymphocytic leukemia (CLL2-BIO): primary endpoint analysis of a multicenter, open-label phase-II trial. <i>Haematologica</i> , 2021 , 106, 543-554	6.6	6

567	Insertion site of central venous catheter correlates with catheter-related infectious events in patients undergoing intensive chemotherapy. <i>Bone Marrow Transplantation</i> , 2021 , 56, 195-201	4.4	2
566	Higher-order connections between stereotyped subsets: implications for improved patient classification in CLL. <i>Blood</i> , 2021 , 137, 1365-1376	2.2	26
565	U-RT1 - A new model for Richter transformation. <i>Neoplasia</i> , 2021 , 23, 140-148	6.4	2
564	No increased bleeding events in patients with relapsed chronic lymphocytic leukemia and indolent non-Hodgkin lymphoma treated with idelalisib. <i>Leukemia and Lymphoma</i> , 2021 , 62, 837-845	1.9	
563	Allogeneic hematopoietic cell transplantation for patients with TP53 mutant or deleted chronic lymphocytic leukemia: Results of a prospective observational study. <i>Bone Marrow Transplantation</i> , 2021 , 56, 692-695	4.4	1
562	LRPAP1 autoantibodies in mantle cell lymphoma are associated with superior outcome. <i>Blood</i> , 2021 , 137, 3251-3258	2.2	1
561	EOMES and IL-10 regulate antitumor activity of T regulatory type 1 CD4 T cells in chronic lymphocytic leukemia. <i>Leukemia</i> , 2021 , 35, 2311-2324	10.7	7
560	Safety and efficacy of obinutuzumab alone or with chemotherapy in previously untreated or relapsed/refractory chronic lymphocytic leukaemia patients: Final analysis of the Phase IIIb GREEN study. <i>British Journal of Haematology</i> , 2021 , 193, 325-338	4.5	3
559	Longitudinal analyses of CLL in mice identify leukemia-related clonal changes including a Myc gain predicting poor outcome in patients. <i>Leukemia</i> , 2021 ,	10.7	1
558	Tafasitamab combined with idelalisib or venetoclax in patients with CLL previously treated with a BTK inhibitor. <i>Leukemia and Lymphoma</i> , 2021 , 1-12	1.9	1
557	Killer immunoglobulin-like receptor 2DS5 is associated with recovery from coronavirus disease 2019. <i>Intensive Care Medicine Experimental</i> , 2021 , 9, 45	3.7	2
556	Long-term survival of patients with mantle cell lymphoma after autologous haematopoietic stem-cell transplantation in first remission: a post-hoc analysis of an open-label, multicentre, randomised, phase 3 trial. <i>Lancet Haematology</i> , 2021 , 8, e648-e657	14.6	6
555	Multi-platform profiling characterizes molecular subgroups and resistance networks in chronic lymphocytic leukemia. <i>Nature Communications</i> , 2021 , 12, 5395	17.4	1
554	Phase 2 study of obinutuzumab (GA-101), ibrutinib and venetoclax (CLL2-GiVe) in patients with untreated high-risk chronic lymphocytic leukemia. <i>Blood</i> , 2021 ,	2.2	6
553	Role of Specific B-Cell Receptor Antigens in Lymphomagenesis. <i>Frontiers in Oncology</i> , 2020 , 10, 604685	5.3	4
552	DNA methylation of chronic lymphocytic leukemia with differential response to chemotherapy. <i>Scientific Data</i> , 2020 , 7, 133	8.2	1
551	Feasibility and Safety of CD19 Chimeric Antigen Receptor T Cell Treatment for B Cell Lymphoma Relapse after Allogeneic Hematopoietic Stem Cell Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2020 , 26, 1575-1580	4.7	12
550	Prognostic and predictive impact of genetic markers in patients with CLL treated with obinutuzumab and venetoclax. <i>Blood</i> , 2020 , 135, 2402-2412	2.2	43

549	International prognostic score for asymptomatic early-stage chronic lymphocytic leukemia. <i>Blood</i> , 2020 , 135, 1859-1869	2.2	45
548	Noncatalytic Bruton's tyrosine kinase activates PLC γ variants mediating ibrutinib resistance in human chronic lymphocytic leukemia cells. <i>Journal of Biological Chemistry</i> , 2020 , 295, 5717-5736	5.4	16
547	Methylome-based cell-of-origin modeling (Methyl-COOM) identifies aberrant expression of immune regulatory molecules in CLL. <i>Genome Medicine</i> , 2020 , 12, 29	14.4	8
546	Approved and emerging PI3K inhibitors for the treatment of chronic lymphocytic leukemia and non-Hodgkin lymphoma. <i>Expert Opinion on Pharmacotherapy</i> , 2020 , 21, 917-929	4	13
545	Efficacy and Safety of Duvelisib Following Disease Progression on Ofatumumab in Patients with Relapsed/Refractory CLL or SLL in the DUO Crossover Extension Study. <i>Clinical Cancer Research</i> , 2020 , 26, 2096-2103	12.9	18
544	COVID-19 among fit patients with CLL treated with venetoclax-based combinations. <i>Leukemia</i> , 2020 , 34, 2225-2229	10.7	25
543	Revolution of Chronic Lymphocytic Leukemia Therapy: the Chemo-Free Treatment Paradigm. <i>Current Oncology Reports</i> , 2020 , 22, 16	6.3	14
542	Prognostic impact of prevalent chronic lymphocytic leukemia stereotyped subsets: analysis within prospective clinical trials of the German CLL Study Group (GCLLSG). <i>Haematologica</i> , 2020 , 105, 2598-2607	6.6	20
541	Early treatment with FCR versus watch and wait in patients with stage Binet A high-risk chronic lymphocytic leukemia (CLL): a randomized phase 3 trial. <i>Leukemia</i> , 2020 , 34, 2038-2050	10.7	19
540	is an inherited risk factor for CLL through the acquisition of a single-point mutation enabling autonomous BCR signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 4320-4327	11.5	24
539	Prognostic model for newly diagnosed CLL patients in Binet stage A: results of the multicenter, prospective CLL1 trial of the German CLL study group. <i>Leukemia</i> , 2020 , 34, 1038-1051	10.7	14
538	Stromal cell protein kinase C- γ inhibition enhances chemosensitivity in B cell malignancies and overcomes drug resistance. <i>Science Translational Medicine</i> , 2020 , 12,	17.5	10
537	Prognostic and predictive role of gene mutations in chronic lymphocytic leukemia: results from the pivotal phase III study COMPLEMENT1. <i>Haematologica</i> , 2020 , 105, 2440-2447	6.6	14
536	Robust Discovery of Candidate DNA Methylation Cancer Drivers. <i>Blood</i> , 2020 , 136, 33-34	2.2	
535	Fixed-duration venetoclax-obinutuzumab for previously untreated patients with chronic lymphocytic leukemia: Follow-up of efficacy and safety results from the multicenter, open-label, randomized, phase III CLL14 trial. <i>Journal of Clinical Oncology</i> , 2020 , 38, 8027-8027	2.2	2
534	Reconstruction of rearranged T-cell receptor loci by whole genome and transcriptome sequencing gives insights into the initial steps of T-cell prolymphocytic leukemia. <i>Genes Chromosomes and Cancer</i> , 2020 , 59, 261-267	5	7
533	Genomic alterations in high-risk chronic lymphocytic leukemia frequently affect cell cycle key regulators and NOTCH1-regulated transcription. <i>Haematologica</i> , 2020 , 105, 1379-1390	6.6	7
532	Influence of obesity and gender on treatment outcomes in patients with chronic lymphocytic leukemia (CLL) undergoing rituximab-based chemoimmunotherapy. <i>Leukemia</i> , 2020 , 34, 1177-1181	10.7	3

531	TBET-expressing Th1 CD4 T cells accumulate in chronic lymphocytic leukaemia without affecting disease progression in E μ -TCL1 mice. <i>British Journal of Haematology</i> , 2020 , 189, 133-145	4.5	7
530	Elevated Hedgehog activity contributes to attenuated DNA damage responses in aged hematopoietic cells. <i>Leukemia</i> , 2020 , 34, 1125-1134	10.7	3
529	Integration of the B-Cell Receptor Antigen Neurabin-I/SAMD14 Into an Antibody Format as New Therapeutic Approach for the Treatment of Primary CNS Lymphoma. <i>Frontiers in Oncology</i> , 2020 , 10, 580364	5.3	1
528	Prolonged Course of COVID-19-Associated Pneumonia in a B-Cell Depleted Patient After Rituximab. <i>Frontiers in Oncology</i> , 2020 , 10, 1578	5.3	26
527	Venetoclax plus obinutuzumab versus chlorambucil plus obinutuzumab for previously untreated chronic lymphocytic leukaemia (CLL14): follow-up results from a multicentre, open-label, randomised, phase 3 trial. <i>Lancet Oncology, The</i> , 2020 , 21, 1188-1200	21.7	92
526	Therapeutic targeting of mutant p53 in pediatric acute lymphoblastic leukemia. <i>Haematologica</i> , 2020 , 105, 170-181	6.6	19
525	Cryptic insertion of exons 2 and 3 into the immunoglobulin heavy chain locus detected by whole genome sequencing in a case of "-negative" Burkitt lymphoma. <i>Haematologica</i> , 2020 , 105, e202-e205	6.6	15
524	The impact of complex karyotype on the overall survival of patients with relapsed chronic lymphocytic leukemia treated with idelalisib plus rituximab. <i>Leukemia</i> , 2020 , 34, 296-300	10.7	15
523	Oxidative stress as candidate therapeutic target to overcome microenvironmental protection of CLL. <i>Leukemia</i> , 2020 , 34, 115-127	10.7	9
522	Obinutuzumab plus fludarabine and cyclophosphamide in previously untreated, fit patients with chronic lymphocytic leukemia: a subgroup analysis of the GREEN study. <i>Leukemia</i> , 2020 , 34, 441-450	10.7	4
521	Characterization of an HLA-restricted and human cytomegalovirus-specific antibody repertoire with therapeutic potential. <i>Cancer Immunology, Immunotherapy</i> , 2020 , 69, 1535-1548	7.4	2
520	High efficacy of venetoclax plus obinutuzumab in patients with complex karyotype and chronic lymphocytic leukemia. <i>Blood</i> , 2020 , 135, 866-870	2.2	18
519	Telomere Dysfunction in Chronic Lymphocytic Leukemia. <i>Frontiers in Oncology</i> , 2020 , 10, 612665	5.3	2
518	Adjuvant Therapy of High-Risk (Stages IIC-IV) Malignant Melanoma in the Post Interferon-Alpha Era: A Systematic Review and Meta-Analysis. <i>Frontiers in Oncology</i> , 2020 , 10, 637161	5.3	2
517	Time-to-progression after front-line fludarabine, cyclophosphamide, and rituximab chemoimmunotherapy for chronic lymphocytic leukaemia: a retrospective, multicohort study. <i>Lancet Oncology, The</i> , 2019 , 20, 1576-1586	21.7	11
516	Dissecting the Prognostic Significance and Functional Role of Progranulin in Chronic Lymphocytic Leukemia. <i>Cancers</i> , 2019 , 11,	6.6	2
515	Mode of progression after first line treatment correlates with outcome of chronic lymphocytic leukemia (CLL). <i>American Journal of Hematology</i> , 2019 , 94, 1002-1006	7.1	3
514	Venetoclax and Obinutuzumab in Patients with CLL and Coexisting Conditions. <i>New England Journal of Medicine</i> , 2019 , 380, 2225-2236	59.2	368

513	Linking aberrant chromatin features in chronic lymphocytic leukemia to transcription factor networks. <i>Molecular Systems Biology</i> , 2019 , 15, e8339	12.2	20
512	Phase 1 first-in-human trial of the anti-CD37 antibody BI 836826 in relapsed/refractory chronic lymphocytic leukemia. <i>Leukemia</i> , 2019 , 33, 2531-2535	10.7	14
511	Venetoclax resistance and acquired mutations in chronic lymphocytic leukemia. <i>Haematologica</i> , 2019 , 104, e434-e437	6.6	81
510	IGF1R as druggable target mediating PI3K-inhibitor resistance in a murine model of chronic lymphocytic leukemia. <i>Blood</i> , 2019 , 134, 534-547	2.2	25
509	MDM4 Is Targeted by 1q Gain and Drives Disease in Burkitt Lymphoma. <i>Cancer Research</i> , 2019 , 79, 3125-3138	3.8	8
508	Final Results of a Randomized, Phase III Study of Rituximab With or Without Idelalisib Followed by Open-Label Idelalisib in Patients With Relapsed Chronic Lymphocytic Leukemia. <i>Journal of Clinical Oncology</i> , 2019 , 37, 1391-1402	2.2	109
507	Efficacy of venetoclax in relapsed chronic lymphocytic leukemia is influenced by disease and response variables. <i>Blood</i> , 2019 , 134, 111-122	2.2	94
506	Short telomeres are associated with inferior outcome, genomic complexity, and clonal evolution in chronic lymphocytic leukemia. <i>Leukemia</i> , 2019 , 33, 2183-2194	10.7	14
505	Genomic and transcriptomic changes complement each other in the pathogenesis of sporadic Burkitt lymphoma. <i>Nature Communications</i> , 2019 , 10, 1459	17.4	49
504	Spontaneous regression of a plasmablastic lymphoma with MYC rearrangement. <i>British Journal of Haematology</i> , 2019 , 186, e203-e207	4.5	5
503	Long-Term Studies Assessing Outcomes of Ibrutinib Therapy in Patients With Del(11q) Chronic Lymphocytic Leukemia. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2019 , 19, 715-722.e6	2	22
502	Prediction of venetoclax activity in precursor B-ALL by functional assessment of apoptosis signaling. <i>Cell Death and Disease</i> , 2019 , 10, 571	9.8	18
501	Obinutuzumab Alone or Combined with Chemotherapy in Previously Untreated (Fit or Unfit) or Relapsed/Refractory Chronic Lymphocytic Leukemia (CLL) Patients: Final Results from the Phase IIIb GREEN Safety Study with a Focus on Efficacy. <i>Blood</i> , 2019 , 134, 3035-3035	2.2	1
500	Comparison of Overall Survival in High Risk Patients with Minimal Residual Disease after First-Line Treatment across Three Generations of Phase 3 Trials of the German CLL Study Group. <i>Blood</i> , 2019 , 134, 3040-3040	2.2	1
499	Quantitative Analysis of Minimal Residual Disease (MRD) Shows High Rates of Undetectable MRD after Fixed-Duration Chemotherapy-Free Treatment and Serves As Surrogate Marker for Progression-Free Survival: A Prospective Analysis of the Randomized CLL14 Trial. <i>Blood</i> , 2019 , 134, 36-36	2.2	15
498	Primary Analysis of Anti-CD19 Tafasitamab (MOR208) Treatment in Combination with Idelalisib or Venetoclax in R/R CLL Patients Who Failed Prior BTK Inhibitor Therapy (COSMOS Trial). <i>Blood</i> , 2019 , 134, 1754-1754	2.2	7
497	Role of FDG PET/CT to Detect Bone Marrow Involvement in the Initial Staging of Aggressive Non-Hodgkin Lymphoma. <i>Blood</i> , 2019 , 134, 2892-2892	2.2	2
496	Bortezomib-Based Induction and Maintenance Overcomes the Negative Prognostic Impact of Renal Impairment and del17p in Transplant-Eligible Myeloma Patients: Long Term Results from the Phase III HOVON-65/GMMG-HD4 Study after Median 137 Months Follow up. <i>Blood</i> , 2019 , 134, 3308-3308	2.2	2

495	A Prospective, Open-Label, Multicenter, Phase 2 Trial to Evaluate the Safety and Efficacy of the Combination of Tirabrutinib (ONO/GS-4059) and Entospletinib with and without Obinutuzumab in Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2019 , 134, 4297-4297	2.2	5
494	A Prospective, Open-Label, Multicenter, Phase 2 Trial to Evaluate the Safety and Efficacy of the Combination of Tirabrutinib (ONO/GS-4059) and Idelalisib with and without Obinutuzumab in Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2019 , 134, 3047-3047	2.2	3
493	Inherited DNA repair and cell cycle gene defects in chronic lymphocytic leukemia.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 1508-1508	2.2	2
492	Effect of dose modifications on response to duvelisib in patients with relapsed/refractory (R/R) CLL/SLL in the DUO trial.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 7523-7523	2.2	4
491	CLL with Del (17p)/TP53 Mutation. <i>Hematologic Malignancies</i> , 2019 , 97-106	0	
490	Effect of fixed-duration venetoclax plus obinutuzumab (VenG) on progression-free survival (PFS), and rates and duration of minimal residual disease negativity (MRD) in previously untreated patients (pts) with chronic lymphocytic leukemia (CLL) and comorbidities.. <i>Journal of Clinical Oncology</i> , 2019 , 37, 7502-7502	2.2	
489	Eomes and IL-10 Regulate Anti-Tumor Activity of T Cells in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2019 , 134, 4288-4288	2.2	
488	Modelling Single Cell B-Cell Receptor Signaling Reveals Enhanced Activity in Primary CLL Cells Compared to Non-Malignant Cells While Fundamental Network Circuit Topology Remains Stable Even with Novel Therapeutic Inhibitors. <i>Blood</i> , 2019 , 134, 4275-4275	2.2	
487	Venetoclax Resistance in Mantle Cell Lymphoma Is Mediated By BCL-XL and Can be Circumvent By Inhibiting the BH4 Domain of BCL-2. <i>Blood</i> , 2019 , 134, 1507-1507	2.2	1
486	The B-Cell Receptor Antigen ARS2 Can be Integrated into a BAR-Body Format to Treat Diffuse Large B-Cell Lymphomas in Xenograft Mouse Models. <i>Blood</i> , 2019 , 134, 2860-2860	2.2	
485	Identification and Characterization of CMV-Specific, HLA-C*07:02 Restricted Antibodies. <i>Blood</i> , 2019 , 134, 5616-5616	2.2	
484	Telomere Shortening By Terc Knockout in the Eµ-TCL1 Transgenic Murine Model of CLL: Characterization of Disease Development and Survival. <i>Blood</i> , 2019 , 134, 1732-1732	2.2	
483	Characterization of Mechanisms of Acquired Venetoclax-Insensitivity in B-Cell Precursor Acute Lymphoblastic Leukemia. <i>Blood</i> , 2019 , 134, 3954-3954	2.2	
482	Response to Comment by Jonathan Weiss. <i>Haematologica</i> , 2019 , 104, e542	6.6	1
481	Four versus six cycles of CHOP chemotherapy in combination with six applications of rituximab in patients with aggressive B-cell lymphoma with favourable prognosis (FLYER): a randomised, phase 3, non-inferiority trial. <i>Lancet, The</i> , 2019 , 394, 2271-2281	4.0	85
480	From Biology to Therapy: The CLL Success Story. <i>HemaSphere</i> , 2019 , 3, e175	0.3	29
479	PI3K inhibition modulates regulatory and effector T-cell differentiation and function in chronic lymphocytic leukemia. <i>Leukemia</i> , 2019 , 33, 1427-1438	10.7	36
478	CLL2-BIG: sequential treatment with bendamustine, ibrutinib and obinutuzumab (GA101) in chronic lymphocytic leukemia. <i>Leukemia</i> , 2019 , 33, 1161-1172	10.7	26

477	The involvement of microRNA in the pathogenesis of Richter syndrome. <i>Haematologica</i> , 2019 , 104, 1004-1015	10	14
476	Prognostic value of MRD in CLL patients with comorbidities receiving chlorambucil plus obinutuzumab or rituximab. <i>Blood</i> , 2019 , 133, 494-497	2.2	18
475	mutations reduce binding of NOTCH1, leading to cleaved NOTCH1 accumulation and target gene activation in CLL. <i>Blood</i> , 2019 , 133, 830-839	2.2	33
474	New lessons learned in T-PLL: results from a prospective phase-II trial with fludarabine-mitoxantrone-cyclophosphamide-alemtuzumab induction followed by alemtuzumab maintenance. <i>Leukemia and Lymphoma</i> , 2019 , 60, 649-657	1.9	10
473	Control of chronic lymphocytic leukemia development by clonally-expanded CD8 T-cells that undergo functional exhaustion in secondary lymphoid tissues. <i>Leukemia</i> , 2019 , 33, 625-637	10.7	42
472	Anti-CD20 immunotherapy as a bridge to tolerance, after allogeneic stem cell transplantation for patients with chronic lymphocytic leukaemia: results of the CLLX4 trial. <i>British Journal of Haematology</i> , 2019 , 184, 833-836	4.5	4
471	Optimising outcomes for patients with chronic lymphocytic leukaemia on ibrutinib therapy: European recommendations for clinical practice. <i>British Journal of Haematology</i> , 2018 , 180, 666-679	4.5	38
470	Tumor necrosis factor receptor signaling is a driver of chronic lymphocytic leukemia that can be therapeutically targeted by the flavonoid wogonin. <i>Haematologica</i> , 2018 , 103, 688-697	6.6	13
469	iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. <i>Blood</i> , 2018 , 131, 2745-2760	2.2	607
468	CLL2-BXX Phase II trials: sequential, targeted treatment for eradication of minimal residual disease in chronic lymphocytic leukemia. <i>Future Oncology</i> , 2018 , 14, 499-513	3.6	20
467	Telomere length in poor-risk chronic lymphocytic leukemia: associations with disease characteristics and outcome. <i>Leukemia and Lymphoma</i> , 2018 , 59, 1614-1623	1.9	11
466	NFATC1 activation by DNA hypomethylation in chronic lymphocytic leukemia correlates with clinical staging and can be inhibited by ibrutinib. <i>International Journal of Cancer</i> , 2018 , 142, 322-333	7.5	24
465	Venetoclax: Targeting BCL2 in Hematological Cancers. <i>Recent Results in Cancer Research</i> , 2018 , 212, 215-242	13	
464	Safety of obinutuzumab alone or combined with chemotherapy for previously untreated or relapsed/refractory chronic lymphocytic leukemia in the phase IIIb GREEN study. <i>Haematologica</i> , 2018 , 103, 1889-1898	6.6	13
463	High-risk chronic lymphocytic leukemia in the era of pathway inhibitors: integrating molecular and cellular therapies. <i>Blood</i> , 2018 , 132, 892-902	2.2	64
462	Obinutuzumab plus bendamustine in previously untreated patients with CLL: a subgroup analysis of the GREEN study. <i>Leukemia</i> , 2018 , 32, 1778-1786	10.7	20
461	Bendamustine followed by obinutuzumab and venetoclax in chronic lymphocytic leukaemia (CLL2-BAG): primary endpoint analysis of a multicentre, open-label, phase 2 trial. <i>Lancet Oncology</i> , 2018 , 19, 1215-1228	21.7	70
460	Evaluation of 230 patients with relapsed/refractory deletion 17p chronic lymphocytic leukaemia treated with ibrutinib from 3 clinical trials. <i>British Journal of Haematology</i> , 2018 , 182, 504-512	4.5	32

459	Comprehensive Safety Analysis of Venetoclax Monotherapy for Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , 2018 , 24, 4371-4379	12.9	90
458	Minimal Residual Disease Status with Venetoclax Monotherapy Is Associated with Progression-Free Survival in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2018 , 132, 3134-3134	2.2	3
457	Excellent Outcome of Young Patients (18-60 years) with Favourable-Prognosis Diffuse Large B-Cell Lymphoma (DLBCL) Treated with 4 Cycles CHOP Plus 6 Applications of Rituximab: Results of the 592 Patients of the Flyer Trial of the Dshnl/GLA. <i>Blood</i> , 2018 , 132, 781-781	2.2	8
456	Durable Remissions after Discontinuation of Combined Targeted Treatment in Patients with Chronic Lymphocytic Leukemia (CLL) Harboring a High-Risk Genetic Lesion (del(17p)/TP53 Mutation). <i>Blood</i> , 2018 , 132, 694-694	2.2	6
455	Residual Abdominal Lymphadenopathy after Intensive Frontline Chemoimmunotherapy Is Associated with Inferior Outcome Regardless of MRD Status in Advanced Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2018 , 132, 4430-4430	2.2	1
454	Identification of Posttranslationally Modified Neoantigens As Targets of B Cell Receptors of Burkitt Lymphoma. <i>Blood</i> , 2018 , 132, 1588-1588	2.2	2
453	Clinical and Biological Indicators of Duvelisib Efficacy in CLL from the Phase 3 DUOTM Study. <i>Blood</i> , 2018 , 132, 1856-1856	2.2	2
452	Hypophosphatemia after High Dosage Iron Substitution with Ferric Carboxymaltose (FCM) and Iron Isomaltoside (IM) In the Randomised Controlled Home Affairs 1 Trial. <i>Blood</i> , 2018 , 132, 3627-3627	2.2	4
451	Two-Cohort Phase II Study in R/R CLL (COSMOS): First Preliminary Safety and Efficacy Results of Anti-CD19 MOR208 Treatment in Combination with Venetoclax in Patients Who Discontinued Prior BTK Inhibitor Therapy. <i>Blood</i> , 2018 , 132, 4433-4433	2.2	2
450	NOTCH1 Signaling Is Activated in CLL By Mutations of FBXW7 and Low Expression of USP28 at 11q23. <i>Blood</i> , 2018 , 132, 946-946	2.2	1
449	Vecabrutinib Is Efficacious In Vivo in a Preclinical CLL Adoptive Transfer Model. <i>Blood</i> , 2018 , 132, 1868-1868	2.2	1
448	The Efficacy and Safety of Duvelisib Following Disease Progression on Ofatumumab in Patients with Relapsed/Refractory CLL or SLL: Updated Results from the DUO Crossover Extension Study. <i>Blood</i> , 2018 , 132, 3140-3140	2.2	1
447	Safety and Efficacy of Venetoclax (VEN) in Combination with Bendamustine (B) Plus Rituximab (R) or Obinutuzumab (G) in Patients (pts) with Previously Untreated Chronic Lymphocytic Leukemia (CLL): Results from a Phase Ib Study (GO28440). <i>Blood</i> , 2018 , 132, 1859-1859	2.2	0
446	Duvelisib inhibition of chemokines in patients with CLL (DUO study) and iNHL (DYNAMO study).. <i>Journal of Clinical Oncology</i> , 2018 , 36, 12048-12048	2.2	1
445	The efficacy of duvelisib monotherapy following disease progression on ofatumumab monotherapy in patients with relapsed/refractory CLL or SLL in the DUO crossover extension study.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 7533-7533	2.2	1
444	The GAIA (CLL13) trial: An international intergroup phase III study for frontline therapy in chronic lymphocytic leukemia (CLL).. <i>Journal of Clinical Oncology</i> , 2018 , 36, TPS7582-TPS7582	2.2	4
443	A phase 1, open-label, multicenter, non-randomized study to assess the safety, tolerability, pharmacokinetics, and preliminary antitumor activity of AZD4573, a potent and selective CDK9 inhibitor, in subjects with relapsed or refractory hematological malignancies.. <i>Journal of Clinical Oncology</i> , 2018 , 36, TPS7588-TPS7588	2.2	2
442	Functional characterization of phospholipase C- μ mutant protein causing both somatic ibrutinib resistance and a germline monogenic autoinflammatory disorder. <i>Oncotarget</i> , 2018 , 9, 34357-34378	3.3	11

441	B-cell acute lymphoblastic leukemia (B-ALL) in CLL patients treated with lenalidomide.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 7531-7531	2.2	
440	Comparison of different phase II studies using sequential combinations of targeted agents for treating chronic lymphocytic leukemia.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 7513-7513	2.2	
439	Prognostic role of beta-2 microglobulin (B2M) in relapsed/refractory (R/R) chronic lymphocytic leukemia (CLL) patients (pts) treated with ibrutinib (ibr).. <i>Journal of Clinical Oncology</i> , 2018 , 36, 7521-7521 ²	2.2	1
438	Mitigation of tumor lysis syndrome (TLS) complications with venetoclax (VEN) in CLL.. <i>Journal of Clinical Oncology</i> , 2018 , 36, 7526-7526	2.2	1
437	Gene Set Enrichment Analysis Suggests That Increased Rituximab-Mediated NK Cell Cytotoxicity after Vitamin D Substitution Is Driven By Upregulation of Interferon Alpha (IFN- α) Isoforms. <i>Blood</i> , 2018 , 132, 3696-3696	2.2	
436	Integrated Proteomic and Phosphoproteomic Analysis Reveal Novel Targets and Suggest Rationale for Ibrutinib Efficacy in UM-CLL. <i>Blood</i> , 2018 , 132, 583-583	2.2	
435	Modeling of the Epigenome of the Cell-of-Origin Identifies Cancer-Specific DNA Methylation Patterns in CLL. <i>Blood</i> , 2018 , 132, 3885-3885	2.2	
434	Obesity Negatively Impacts Outcome in Female Patients with Chronic Lymphocytic Leukemia (CLL) Treated with Fludarabine, Cyclophosphamide and Rituximab (FCR): An Analysis of Three Phase III Studies of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2018 , 132, 4429-4429	2.2	
433	MYC Pathway Activation Is Frequently Observed in Treatment-Naive CLL and Defines a Subgroup with Particular Benefit from the Addition of Rituximab to Chemotherapy. <i>Blood</i> , 2018 , 132, 1866-1866	2.2	
432	Characterization of Mechanisms Underlying Acquired Venetoclax-Resistance in Mantle Cell Lymphoma: BDA-366 As a Potential Treatment Option. <i>Blood</i> , 2018 , 132, 1580-1580	2.2	
431	BAR-Bodies: B-Cell Receptor Antigens As the Targeting Moiety of Antibodies in Substitution for the Variable Region of Heavy and Light Chains. <i>Blood</i> , 2018 , 132, 2940-2940	2.2	0
430	Venetoclax for Patients With Chronic Lymphocytic Leukemia With 17p Deletion: Results From the Full Population of a Phase II Pivotal Trial. <i>Journal of Clinical Oncology</i> , 2018 , 36, 1973-1980	2.2	174
429	aberrations in chronic lymphocytic leukemia: an overview of the clinical implications of improved diagnostics. <i>Haematologica</i> , 2018 , 103, 1956-1968	6.6	43
428	Outcome of patients aged 80 years or older treated for chronic lymphocytic leukaemia. <i>British Journal of Haematology</i> , 2018 , 183, 727-735	4.5	6
427	The phase 3 DUO trial: duvelisib vs ofatumumab in relapsed and refractory CLL/SLL. <i>Blood</i> , 2018 , 132, 2446-2455	2.2	184
426	IG- neoplasms with precursor B-cell phenotype are molecularly distinct from Burkitt lymphomas. <i>Blood</i> , 2018 , 132, 2280-2285	2.2	19
425	Chronische Lymphatische Leukämie. <i>Klinikerarzt</i> , 2018 , 47, 412-417	0	
424	Associations of ofatumumab exposure and treatment outcomes in patients with untreated CLL receiving chemoimmunotherapy. <i>Leukemia and Lymphoma</i> , 2017 , 58, 348-356	1.9	2

423	Venetoclax in Patients with Previously Treated Chronic Lymphocytic Leukemia. <i>Clinical Cancer Research</i> , 2017 , 23, 4527-4533	12.9	43
422	Front-line treatment of CLL in the era of novel agents. <i>Cancer Treatment Reviews</i> , 2017 , 53, 70-78	14.4	19
421	Idelalisib or placebo in combination with bendamustine and rituximab in patients with relapsed or refractory chronic lymphocytic leukaemia: interim results from a phase 3, randomised, double-blind, placebo-controlled trial. <i>Lancet Oncology, The</i> , 2017 , 18, 297-311	21.7	173
420	Human NACHT, LRR, and PYD domain-containing protein 3 (NLRP3) inflammasome activity is regulated by and potentially targetable through Bruton tyrosine kinase. <i>Journal of Allergy and Clinical Immunology</i> , 2017 , 140, 1054-1067.e10	11.5	72
419	Chronic Lymphocytic Leukemia with Mutated IGHV4-34 Receptors: Shared and Distinct Immunogenetic Features and Clinical Outcomes. <i>Clinical Cancer Research</i> , 2017 , 23, 5292-5301	12.9	20
418	TP53 mutation and survival in aggressive B cell lymphoma. <i>International Journal of Cancer</i> , 2017 , 141, 1381-1388	7.5	39
417	Ibrutinib-resistant CLL: unwanted and unwonted!. <i>Blood</i> , 2017 , 129, 1407-1409	2.2	5
416	Venetoclax and obinutuzumab in chronic lymphocytic leukemia. <i>Blood</i> , 2017 , 129, 2702-2705	2.2	90
415	Impact of telomere length on the outcome of allogeneic stem cell transplantation for poor-risk chronic lymphocytic leukaemia: results from the GCLLSG CLL3X trial. <i>British Journal of Haematology</i> , 2017 , 179, 342-346	4.5	2
414	Whole-brain radiotherapy or autologous stem-cell transplantation as consolidation strategies after high-dose methotrexate-based chemoimmunotherapy in patients with primary CNS lymphoma: results of the second randomisation of the International Extranodal Lymphoma Study Group-32 phase 2 trial. <i>Lancet Haematology, the</i> , 2017 , 4, e510-e523	14.6	157
413	Analysis of rare germ line variants in chronic lymphocytic leukemia. <i>Blood</i> , 2017 , 130, 2443-2444	2.2	4
412	Profile of venetoclax and its potential in the context of treatment of relapsed or refractory chronic lymphocytic leukemia. <i>OncoTargets and Therapy</i> , 2017 , 10, 645-656	4.4	7
411	Chronic Lymphocytic Leukemia With TP53 Aberrations: Breakthroughs and Challenges. <i>Journal of Oncology Practice</i> , 2017 , 13, 381-382	3.1	1
410	Tumor-derived exosomes modulate PD-L1 expression in monocytes. <i>Science Immunology</i> , 2017 , 2,	2.8	170
409	Lenalidomide maintenance after first-line therapy for high-risk chronic lymphocytic leukaemia (CLLM1): final results from a randomised, double-blind, phase 3 study. <i>Lancet Haematology, the</i> , 2017 , 4, e475-e486	14.6	36
408	Allogeneic hematopoietic cell transplantation for high-risk CLL: 10-year follow-up of the GCLLSG CLL3X trial. <i>Blood</i> , 2017 , 130, 1477-1480	2.2	49
407	Two mouse models reveal an actionable PARP1 dependence in aggressive chronic lymphocytic leukemia. <i>Nature Communications</i> , 2017 , 8, 153	17.4	29
406	Rapid detection of ATM/p53 function with p27Kip FACS analysis: a novel diagnostic tool for chronic lymphocytic leukemia?. <i>Leukemia and Lymphoma</i> , 2017 , 58, 6-7	1.9	1

405	Alemtuzumab consolidation in chronic lymphocytic leukaemia: a phase I/II multicentre trial. <i>European Journal of Haematology</i> , 2017 , 98, 254-262	3.8	7
404	COSMOS: MOR208 plus idelalisib or venetoclax in patients with relapsed or refractory (R/R) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) previously treated with a Bruton's tyrosine kinase inhibitor (BTKi) A two-cohort phase II study.. <i>Journal of Clinical Oncology</i> , 2017 , 35, TPS7567-TPS7567	2.2	
403	Minimal Residual Disease Assessment Improves Prediction of Outcome in Patients With Chronic Lymphocytic Leukemia (CLL) Who Achieve Partial Response: Comprehensive Analysis of Two Phase III Studies of the German CLL Study Group. <i>Journal of Clinical Oncology</i> , 2016 , 34, 3758-3765	2.2	111
402	Ibrutinib for patients with relapsed or refractory chronic lymphocytic leukaemia with 17p deletion (RESONATE-17): a phase 2, open-label, multicentre study. <i>Lancet Oncology, The</i> , 2016 , 17, 1409-1418	21.7	233
401	The Phospholipase C Mutants R665W and L845F Identified in Ibrutinib-resistant Chronic Lymphocytic Leukemia Patients Are Hypersensitive to the Rho GTPase Rac2 Protein. <i>Journal of Biological Chemistry</i> , 2016 , 291, 22136-22148	5.4	29
400	Telomere shortening leads to an acceleration of synucleinopathy and impaired microglia response in a genetic mouse model. <i>Acta Neuropathologica Communications</i> , 2016 , 4, 87	7.3	28
399	Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. <i>Lancet, The</i> , 2016 , 388, 565-75	4.0	233
398	Krüpel-like factor 4 (KLF4) inactivation in chronic lymphocytic leukemia correlates with promoter DNA-methylation and can be reversed by inhibition of NOTCH signaling. <i>Haematologica</i> , 2016 , 101, e249-53	6.6	17
397	Efficacy of antineoplastic treatment is associated with the use of antibiotics that modulate intestinal microbiota. <i>Onc Immunology</i> , 2016 , 5, e1150399	7.2	55
396	DNA methylation dynamics during B cell maturation underlie a continuum of disease phenotypes in chronic lymphocytic leukemia. <i>Nature Genetics</i> , 2016 , 48, 253-64	36.3	193
395	Genomic Features: Impact on Pathogenesis and Treatment of Chronic Lymphocytic Leukemia. <i>Oncology Research and Treatment</i> , 2016 , 39, 34-40	2.8	3
394	Evaluation of geriatric assessment in patients with chronic lymphocytic leukemia: Results of the CLL9 trial of the German CLL study group. <i>Leukemia and Lymphoma</i> , 2016 , 57, 789-96	1.9	66
393	In Vivo modeling of Resistance to PI3K Inhibitor Treatment Using EµTCL1-Tg Tumor Transfer Model. <i>Blood</i> , 2016 , 128, 190-190	2.2	4
392	Outcome of Patients with Complex Karyotype in a Phase 3 Randomized Study of Idelalisib Plus Rituximab for Relapsed Chronic Lymphocytic Leukemia. <i>Blood</i> , 2016 , 128, 192-192	2.2	10
391	11q Deletion (del11q) Is Not a Prognostic Factor for Adverse Outcomes for Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) Treated with Ibrutinib: Pooled Data from 3 Randomized Phase 3 Studies. <i>Blood</i> , 2016 , 128, 2042-2042	2.2	6
390	Low Incidence of Tumor Lysis Syndromes (TLS) and Infusion Related Reactions (IRR) in the CLL2-Bag Trial Evaluating a Sequential Treatment of Bendamustine (B), Obinutuzumab (GA101, G) and Venetoclax (ABT-199, A) in Patients with Chronic Lymphocytic Leukemia (CLL): Interim Safety Results of the CLL2-Bag Trial. <i>Blood</i> , 2016 , 128, 2054-2054	2.2	4
389	Safety and Efficacy of Venetoclax and Obinutuzumab in Patients with Previously Untreated Chronic Lymphocytic Leukemia (CLL) and Coexisting Medical Conditions: Final Results of the Run-in Phase of the Randomized CLL14 Trial (BO25323). <i>Blood</i> , 2016 , 128, 2054-2054	2.2	8
388	Lenalidomide Maintenance after Front Line Therapy Substantially Prolongs Progression Free Survival in High Risk CLL: Interim Results of a Phase 3 Study (CLL M1 study of the German CLL Study Group). <i>Blood</i> , 2016 , 128, 229-229	2.2	10

387	Updated Analysis of Overall Survival in Randomized Phase III Study of Idelalisib in Combination with Bendamustine and Rituximab in Patients with Relapsed/Refractory CLL. <i>Blood</i> , 2016 , 128, 231-231	2.2	4
386	Whole-Exome Sequencing Revealed No Recurrent Mutations within the PI3K Pathway in Relapsed Chronic Lymphocytic Leukemia Patients Progressing Under Idelalisib Treatment. <i>Blood</i> , 2016 , 128, 2770-2770	2.2	16
385	CLL Exosome-Derived Y RNA hY4 Induces TLR7/8-Mediated Inflammation and PD-L1 Expression in Monocytes. <i>Blood</i> , 2016 , 128, 3217-3217	2.2	1
384	Gene Mutations and Treatment Outcome in the Context of Chlorambucil (Clb) without or with the Addition of Rituximab (R) or Obinutuzumab (GA-101, G) - Results of an Extensive Analysis of the Phase III Study CLL11 of the German CLL Study Group. <i>Blood</i> , 2016 , 128, 3227-3227	2.2	13
383	Pooled Multi-Trial Analysis of Venetoclax Efficacy in Patients with Relapsed or Refractory Chronic Lymphocytic Leukemia. <i>Blood</i> , 2016 , 128, 3230-3230	2.2	11
382	Reappraising Immunoglobulin Repertoire Restrictions in Chronic Lymphocytic Leukemia: Focus on Major Stereotyped Subsets and Closely Related Satellites. <i>Blood</i> , 2016 , 128, 4376-4376	2.2	1
381	Favorable Toxicity Profile and Long Term Outcome of Elderly, but Physically Fit CLL Patients (pts) Receiving First Line Bendamustine and Rituximab (BR) Frontline Chemoimmunotherapy in Comparison to Fludarabine, Cyclophosphamide, and Rituximab (FCR) in Advanced Chronic Lymphocytic Leukemia (CLL): Update Analysis of an International Randomized Study of the German CLL Study Group (GCLLSG) (CLL11 Study). <i>Blood</i> , 2016 , 128, 4387-4387	2.2	9
380	Bendamustine/Obinutuzumab in Patients with Relapsed/Refractory or Previously Untreated Chronic Lymphocytic Leukemia. <i>Blood</i> , 2016 , 128, 4393-4393	2.2	16
379	Safety Profile of Venetoclax Monotherapy in Patients with Chronic Lymphocytic Leukemia. <i>Blood</i> , 2016 , 128, 4395-4395	2.2	6
378	Evaluation of the International Prognostic Index for Chronic Lymphocytic Leukemia (CLL-IPI) in Elderly Patients with Comorbidities: Analysis of the CLL11 Study Population. <i>Blood</i> , 2016 , 128, 4401-4401	2.2	8
377	Effects on Survival and Neurocognitive Functions of Whole-Brain Radiotherapy (WBRT) and Autologous Stem Cell Transplantation (ASCT) as Consolidation Options after High-Dose Methotrexate-Based Chemoimmunotherapy in Patients with Newly Diagnosed Primary CNS Lymphoma (PCNSL): Results of the Second Randomization of the IFLS02 Trial. <i>Blood</i> , 2016 , 128, 511-511	2.2	13
376	CLL2-BIG - a Novel Treatment Regimen of Bendamustine Followed By GA101 and Ibrutinib Followed By Ibrutinib and GA101 Maintenance in Patients with Chronic Lymphocytic Leukemia (CLL): Results of a Phase II-Trial. <i>Blood</i> , 2016 , 128, 640-640	2.2	9
375	Long-Term Outcome of Allogeneic Hematopoietic Stem Cell Transplantation (HSCT) for Chronic Lymphocytic Leukemia (CLL): 10-Year Follow-up of the Gcllsg CLL3X Trial. <i>Blood</i> , 2016 , 128, 682-682	2.2	3
374	In Chronic Lymphocytic Leukemia CT7544-7545 Mutant NOTCH1 Maintains Transcription Factor Activity with Longer Lasting Effects Due to Slower Degradation. <i>Blood</i> , 2016 , 128, 971-971	2.2	
373	High-Dose Chemotherapy with Autologous Hematopoietic Stem Cell Support for Relapsed or Refractory Primary CNS Lymphoma - a Prospective Multicentre Trial By the German Cooperative PCNSL Study Group. <i>Blood</i> , 2016 , 128, 781-781	2.2	
372	Impact of Gender on Outcome after Chemoimmunotherapy with Fludarabine, Cyclophosphamide and Rituximab (FCR) or Bendamustine Plus Rituximab (BR) in Patients with Chronic Lymphocytic Leukemia (CLL): A Meta-Analysis of Three Phase II/III Studies of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2016 , 128, 1201-1201	2.2	0
371	Anti-Leukemia Activity of the Selective BCL-2 Inhibitor ABT-199 in Childhood B-Cell Precursor Acute Lymphoblastic Leukemia Is Characterized By MCL-1/BCL-2 Expression Serving As Biomarker for Treatment Response. <i>Blood</i> , 2016 , 128, 1081-1081	2.2	0
370	Assessment of Measures to Reduce Infusion-Related Reactions in Patients with Chronic Lymphocytic Leukemia Treated with Obinutuzumab in the GREEN Study. <i>Blood</i> , 2016 , 128, 5589-5589	2.2	

369	Telomere dysfunction and chromothripsis. <i>International Journal of Cancer</i> , 2016 , 138, 2905-14	7.5	34
368	Bendamustine and rituximab in combination with lenalidomide in patients with chronic lymphocytic leukemia. <i>European Journal of Haematology</i> , 2016 , 97, 253-60	3.8	18
367	Alterations of microRNA and microRNA-regulated messenger RNA expression in germinal center B-cell lymphomas determined by integrative sequencing analysis. <i>Haematologica</i> , 2016 , 101, 1380-1389	6.6	31
366	Venetoclax in relapsed or refractory chronic lymphocytic leukaemia with 17p deletion: a multicentre, open-label, phase 2 study. <i>Lancet Oncology, The</i> , 2016 , 17, 768-778	21.7	536
365	First-line chemoimmunotherapy with bendamustine and rituximab versus fludarabine, cyclophosphamide, and rituximab in patients with advanced chronic lymphocytic leukaemia (CLL10): an international, open-label, randomised, phase 3, non-inferiority trial. <i>Lancet Oncology, The</i> , 2016 , 17, 928-942	21.7	416
364	Prognostic Value of Ki-67 Index, Cytology, and Growth Pattern in Mantle-Cell Lymphoma: Results From Randomized Trials of the European Mantle Cell Lymphoma Network. <i>Journal of Clinical Oncology</i> , 2016 , 34, 1386-94	2.2	197
363	Chemoimmunotherapy with methotrexate, cytarabine, thiotepa, and rituximab (MATRix regimen) in patients with primary CNS lymphoma: results of the first randomisation of the International Extranodal Lymphoma Study Group-32 (IELSG32) phase 2 trial. <i>Lancet Haematology,the</i> , 2016 , 3, e217-27	14.6	288
362	Safety and efficacy of different lenalidomide starting doses in patients with relapsed or refractory chronic lymphocytic leukemia: results of an international multicenter double-blinded randomized phase II trial. <i>Leukemia and Lymphoma</i> , 2016 , 57, 1291-9	1.9	16
361	Long-term remissions after FCR chemoimmunotherapy in previously untreated patients with CLL: updated results of the CLL8 trial. <i>Blood</i> , 2016 , 127, 208-15	2.2	442
360	Complex karyotypes and KRAS and POT1 mutations impact outcome in CLL after chlorambucil-based chemotherapy or chemoimmunotherapy. <i>Blood</i> , 2016 , 128, 395-404	2.2	92
359	Postibrutinib outcomes in patients with mantle cell lymphoma. <i>Blood</i> , 2016 , 127, 1559-63	2.2	171
358	High-dose chemotherapy with autologous haemopoietic stem cell transplantation for newly diagnosed primary CNS lymphoma: a prospective, single-arm, phase 2 trial. <i>Lancet Haematology,the</i> , 2016 , 3, e388-97	14.6	93
357	Clinical Practice Recommendations for Use of Allogeneic Hematopoietic Cell Transplantation in Chronic Lymphocytic Leukemia on Behalf of the Guidelines Committee of the American Society for Blood and Marrow Transplantation. <i>Biology of Blood and Marrow Transplantation</i> , 2016 , 22, 2117-2125	4.7	70
356	Ofatumumab in poor-prognosis chronic lymphocytic leukemia: a phase IV, non-interventional, observational study from the European Research Initiative on Chronic Lymphocytic Leukemia. <i>Haematologica</i> , 2015 , 100, 511-6	6.6	34
355	Final results of EFC6663: a multicenter, international, phase 2 study of alvocidib for patients with fludarabine-refractory chronic lymphocytic leukemia. <i>Leukemia Research</i> , 2015 , 39, 495-500	2.7	41
354	Ofatumumab monotherapy in fludarabine-refractory chronic lymphocytic leukemia: final results from a pivotal study. <i>Haematologica</i> , 2015 , 100, e311-4	6.6	12
353	Mutations driving CLL and their evolution in progression and relapse. <i>Nature</i> , 2015 , 526, 525-30	50.4	658
352	Obinutuzumab (GA101) for the treatment of chronic lymphocytic leukemia and other B-cell non-hodgkin's lymphomas: a glycoengineered type II CD20 antibody. <i>Oncology Research and Treatment</i> , 2015 , 38, 185-92	2.8	36

351	The CLL12 trial protocol: a placebo-controlled double-blind Phase III study of ibrutinib in the treatment of early-stage chronic lymphocytic leukemia patients with risk of early disease progression. <i>Future Oncology</i> , 2015 , 11, 1895-903	3.6	27
350	MINCR is a MYC-induced lncRNA able to modulate MYC's transcriptional network in Burkitt lymphoma cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, E5261-70	11.5	75
349	Outcome of advanced chronic lymphocytic leukemia following different first-line and relapse therapies: a meta-analysis of five prospective trials by the German CLL Study Group (GCLLSG). <i>Haematologica</i> , 2015 , 100, 1451-9	6.6	26
348	Loss of cooperativity of secreted CD40L and increased dose-response to IL4 on CLL cell viability correlates with enhanced activation of NF- κ B and STAT6. <i>International Journal of Cancer</i> , 2015 , 136, 65-73	7.5	11
347	Prognostic markers and standard management of chronic lymphocytic leukemia. <i>Hematology American Society of Hematology Education Program</i> , 2015 , 2015, 368-77	3.1	24
346	Targeting inhibitor of apoptosis proteins by Smac mimetic elicits cell death in poor prognostic subgroups of chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , 2015 , 137, 2959-70	7.5	14
345	Management of chronic lymphocytic leukemia. <i>American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting</i> , 2015 , 164-75	7.1	20
344	Epigenetic silencing of miR-708 enhances NF- κ B signaling in chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , 2015 , 137, 1352-61	7.5	40
343	How best to manage patients with chronic lymphocytic leukemia with 17p deletion and/or TP53 mutation?. <i>Leukemia and Lymphoma</i> , 2015 , 56, 587-93	1.9	13
342	Progressive Epigenetic Programming during B Cell Maturation Is Reflected in a Continuum of Epigenetic Disease Phenotypes in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2015 , 126, 2436-2436	2.2	1
341	Ibrutinib in Early Stage CLL: Preliminary Safety Results of a Placebo-Controlled Phase III Study. <i>Blood</i> , 2015 , 126, 2934-2934	2.2	5
340	Quantitative Clonal Dynamics Define Mechanisms of CLL Evolution in Response to Combination Chemotherapy. <i>Blood</i> , 2015 , 126, 362-362	2.2	3
339	CLL2-BIG - a Novel Treatment Regimen of Bendamustine Followed By GA101 and Ibrutinib Followed By Ibrutinib and GA101 Maintenance in Patients with Chronic Lymphocytic Leukemia (CLL): Interim Results of a Phase II-Trial. <i>Blood</i> , 2015 , 126, 4151-4151	2.2	7
338	Prognostic Impact and Risk Factors of Reducing Prescribed Doses of Fludarabine, Cyclophosphamide and Rituximab (FCR) during Frontline Treatment of Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2015 , 126, 4156-4156	2.2	12
337	Safety and Efficacy of Obinutuzumab Plus Bendamustine in Previously Untreated Patients with Chronic Lymphocytic Leukemia: Subgroup Analysis of the Green Study. <i>Blood</i> , 2015 , 126, 493-493	2.2	11
336	Results of the Safety Run-in Phase of CLL14 (BO25323): A Prospective, Open-Label, Multicenter Randomized Phase III Trial to Compare the Efficacy and Safety of Obinutuzumab and Venetoclax (GDC-0199/ABT-199) with Obinutuzumab and Chlorambucil in Patients with Previously Untreated CLL and Coexisting Medical Conditions. <i>Blood</i> , 2015 , 126, 496-496	2.2	14
335	Lenalidomide Exhibits Activity in Mantle Cell Lymphoma through Increased NK Cell Mediated Cytotoxicity. <i>Blood</i> , 2015 , 126, 821-821	2.2	1
334	Updated Safety and Preliminary Efficacy Data from a Phase 1b Study Combining Venetoclax (GDC-0199, ABT-199) with Bendamustine/Rituximab in Patients with Relapsed/Refractory or Previously Untreated Chronic Lymphocytic Leukemia. <i>Blood</i> , 2015 , 126, 829-829	2.2	5

333	Outcome of Ibrutinib Treatment by Baseline Genetic Features in Patients with Relapsed or Refractory CLL/SLL with del17p in the Resonate-17 Study. <i>Blood</i> , 2015 , 126, 833-833	2.2	4
332	Targeting Mutant p53 in Pediatric Acute Lymphoblastic Leukemia. <i>Blood</i> , 2015 , 126, 903-903	2.2	1
331	E μ -TCL1mTerc -/- Mouse Model for Telomere Dysfunction in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2015 , 126, 1724-1724	2.2	
330	Genomic Disruption of the Histone Methyltransferase SETD2 in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2015 , 126, 365-365	2.2	
329	Microenvironmental Stromal Cells Rescue CLL Cells from Apoptosis Via Hypoxia That Can be Targeted Therapeutically. <i>Blood</i> , 2015 , 126, 4140-4140	2.2	
328	CLL with Mutated IGHV4-34 Antigen Receptors Is Clinically Heterogeneous: Antigen Receptor Stereotypy Makes the Difference. <i>Blood</i> , 2015 , 126, 5263-5263	2.2	
327	Idelalisib Treatment Is Associated with Improved Cytopenias in Patients with Relapsed/Refractory iNHL and CLL. <i>Blood</i> , 2015 , 126, 1747-1747	2.2	
326	Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. <i>New England Journal of Medicine</i> , 2014 , 370, 997-1007	59.2	1303
325	Obinutuzumab plus chlorambucil in patients with CLL and coexisting conditions. <i>New England Journal of Medicine</i> , 2014 , 370, 1101-10	59.2	1048
324	Development of a comprehensive prognostic index for patients with chronic lymphocytic leukemia. <i>Blood</i> , 2014 , 124, 49-62	2.2	202
323	Resistance mechanisms for the Bruton's tyrosine kinase inhibitor ibrutinib. <i>New England Journal of Medicine</i> , 2014 , 370, 2286-94	59.2	800
322	Confirmation of the mantle-cell lymphoma International Prognostic Index in randomized trials of the European Mantle-Cell Lymphoma Network. <i>Journal of Clinical Oncology</i> , 2014 , 32, 1338-46	2.2	112
321	Evolution of DNA methylation is linked to genetic aberrations in chronic lymphocytic leukemia. <i>Cancer Discovery</i> , 2014 , 4, 348-61	24.4	115
320	Interactions between comorbidity and treatment of chronic lymphocytic leukemia: results of German Chronic Lymphocytic Leukemia Study Group trials. <i>Haematologica</i> , 2014 , 99, 1095-100	6.6	74
319	MDM2 promotor polymorphism and disease characteristics in chronic lymphocytic leukemia: results of an individual patient data-based meta-analysis. <i>Haematologica</i> , 2014 , 99, 1285-91	6.6	1
318	Gene mutations and treatment outcome in chronic lymphocytic leukemia: results from the CLL8 trial. <i>Blood</i> , 2014 , 123, 3247-54	2.2	352
317	PTK2 expression and immunochemotherapy outcome in chronic lymphocytic leukemia. <i>Blood</i> , 2014 , 124, 420-5	2.2	11
316	Managing high-risk CLL during transition to a new treatment era: stem cell transplantation or novel agents?. <i>Blood</i> , 2014 , 124, 3841-9	2.2	158

315	Poor efficacy and tolerability of R-CHOP in relapsed/refractory chronic lymphocytic leukemia and Richter transformation. <i>American Journal of Hematology</i> , 2014 , 89, E239-43	7.1	55
314	Assessment of TP53 functionality in chronic lymphocytic leukaemia by different assays; an ERIC-wide approach. <i>British Journal of Haematology</i> , 2014 , 167, 565-9	4.5	7
313	Frontline Chemoimmunotherapy with Fludarabine (F), Cyclophosphamide (C), and Rituximab (R) (FCR) Shows Superior Efficacy in Comparison to Bendamustine (B) and Rituximab (BR) in Previously Untreated and Physically Fit Patients (pts) with Advanced Chronic Lymphocytic Leukemia (CLL): Subclonal Analysis of an International, Randomized Study of the German CLL Study Group (GCLLSG) CLL10 Study. <i>Blood</i> , 2014 , 124, 1911-1919	2.2	40
312	Subclonal Driver Mutations Predict Shorter Progression Free Survival in Chronic Lymphocytic Leukemia Following First-Line Chemo(immuno)Therapy: Results from the CLL8 Trial. <i>Blood</i> , 2014 , 124, 1938-1938	2.2	3
311	Alemtuzumab Combined with Dexamethasone, Followed By Alemtuzumab Maintenance or Allo-SCT in Ultra High-risk CLL: Final Results from the CLL20 Phase II Study. <i>Blood</i> , 2014 , 124, 1991-1991	2.2	10
310	Gene Mutations and Treatment Outcome in CLL Patients Treated with Chlorambucil (Chl) or Ofatumumab-Chl (O-Chl): Results from the Phase III Study COMPLEMENT1 (OMB110911). <i>Blood</i> , 2014 , 124, 1992-1992	2.2	2
309	Value of Minimal Residual Disease (MRD) Negative Status at Response Evaluation in Chronic Lymphocytic Leukemia (CLL): Combined Analysis of Two Phase III Studies of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2014 , 124, 23-23	2.2	2
308	Clinical Activity of Abemaciclib (LY2835219), a Cell Cycle Inhibitor Selective for CDK4 and CDK6, in Patients with Relapsed or Refractory Mantle Cell Lymphoma. <i>Blood</i> , 2014 , 124, 3067-3067	2.2	13
307	Efficacy and Safety of Ibrutinib in Patients with Relapsed or Refractory Chronic Lymphocytic Leukemia or Small Lymphocytic Leukemia with 17p Deletion: Results from the Phase II RESONATE17 Trial. <i>Blood</i> , 2014 , 124, 327-327	2.2	29
306	Second Interim Analysis of a Phase 3 Study of Idelalisib (ZYDELIG®) Plus Rituximab (R) for Relapsed Chronic Lymphocytic Leukemia (CLL): Efficacy Analysis in Patient Subpopulations with Del(17p) and Other Adverse Prognostic Factors. <i>Blood</i> , 2014 , 124, 330-330	2.2	54
305	Salvage Therapy with Obinutuzumab (GA101) Plus Chlorambucil (Clb) after Treatment Failure of Clb Alone in Patients with Chronic Lymphocytic Leukemia (CLL) and Comorbidities: Results of the CLL11 Study. <i>Blood</i> , 2014 , 124, 3327-3327	2.2	5
304	Preliminary Results of a Phase 1b Study (GO28440) Combining GDC-0199 (ABT-199) with Bendamustine/Rituximab in Patients with Relapsed/Refractory or Previously Untreated Chronic Lymphocytic Leukemia. <i>Blood</i> , 2014 , 124, 3337-3337	2.2	4
303	Preliminary Safety Results from the Phase IIIb GREEN Study of Obinutuzumab (GA101) Alone or in Combination with Chemotherapy for Previously Untreated or Relapsed/Refractory Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2014 , 124, 3345-3345	2.2	8
302	Single-Agent Ibrutinib Demonstrates Safety and Durability of Response at 2 Years Follow-up in Patients with Relapsed or Refractory Mantle Cell Lymphoma: Updated Results of an International, Multicenter, Open-Label Phase 2 Study. <i>Blood</i> , 2014 , 124, 4453-4453	2.2	11
301	Good Tolerance of Lenalidomide Maintenance Therapy in Patients with High Risk Profile Chronic Lymphocytic Leukemia (CLL) after Frontline Chemoimmunotherapy: Preliminary Safety Overview of the CLLM1 Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2014 , 124, 4699-4699	2.2	2
300	Advances in treating chronic lymphocytic leukemia. <i>F1000prime Reports</i> , 2014 , 6, 65		12
299	Identification of the Antigenic Targets of the Major CLL-Derived Stereotyped BCRs and First Demonstration of Sub-Subsets. <i>Blood</i> , 2014 , 124, 298-298	2.2	
298	Charting Unique Signatures of Somatic Hypermutation Amongst Chronic Lymphocytic Leukemia Patients Expressing IGHV4-34 Clonotypic B Cell Receptors. <i>Blood</i> , 2014 , 124, 1969-1969	2.2	

297	Telomere Length and Outcome of Allogeneic Stem Cell Transplantation for Poor Risk Chronic Lymphocytic Leukemia: Results from the GCLLSG CLL3X Trial. <i>Blood</i> , 2014 , 124, 1951-1951	2.2	
296	Chronic Lymphocytic Leukemia-Derived Extracellular Vesicles Contain a Distinctive Proteome, As Well As Specific Micro RNAs and Y RNAs. <i>Blood</i> , 2014 , 124, 1968-1968	2.2	1
295	NFATc1 Is Transcriptionally Activated in Chronic Lymphocytic Leukemia (CLL) By Promotor DNA-Hypomethylation Which Correlates with in-Vitro Vulnerability to Calcineurin Inhibitors. <i>Blood</i> , 2014 , 124, 1941-1941	2.2	
294	High-Resolution Genomic Copy Number Analysis on Sequential Samples from the CLL8 Trial: Relation Between Clonal Evolution and Defects in DNA Damage Response?. <i>Blood</i> , 2014 , 124, 1964-1964 ^{2,2}	2.2	
293	High Dose-Chemotherapy Followed By Autologous Peripheral Blood Stem Cell Transplantation for Patients with Refractory or Recurrent Primary Central Nervous System Lymphoma Results of a Multicenter Study By the Germany Collaborative PCNSL Study Group. <i>Blood</i> , 2014 , 124, 2527-2527	2.2	
292	Allogeneic HCT in Patients with 17p-CLL: Results of a Non-Interventional Study of the EBMT & Eric. <i>Blood</i> , 2014 , 124, 1224-1224	2.2	
291	The Flavonoid Wogonin Reduces CLL Cell Survival in Vitro and Leukemia Development in Eµ-TCL1 Mice By Targeting Aberrant TNF Receptor Signaling. <i>Blood</i> , 2014 , 124, 1966-1966	2.2	
290	High Resolution Genomic Profiling of Primary Ultra High Risk and Refractory Chronic Lymphocytic Leukemia: Results from the CLL2O Trial. <i>Blood</i> , 2014 , 124, 3288-3288	2.2	
289	Genomic Mechanisms of 17p / TP53 Loss in Primary Ultra High-risk and Refractory Chronic Lymphocytic Leukemia: Results from the CLL2O Trial. <i>Blood</i> , 2014 , 124, 2184-2184	2.2	
288	Induction Treatment with Alemtuzumab Combined with Polychemotherapy Containing Fludarabine, Cyclophosphamide and Mitoxantrone (FCM) Followed By Alemtuzumab Maintenance in Patients with T-Cell Prolymphocytic Leukaemia (T-PLL): First Analysis of a Prospective Multicenter Phase-II Trial (T-PLL2) Conducted By the German CLL Study Group (GCLLSG). <i>Blood</i> , 2014 , 124, 1000-1000	2.2	
287	MicroRNAs in Hematologic Malignancies 2014 , 67-95		
286	17p deletion in chronic lymphocytic leukemia: risk stratification and therapeutic approach. <i>Hematology/Oncology Clinics of North America</i> , 2013 , 27, 289-301	3.1	10
285	Targeting BTK with ibrutinib in relapsed or refractory mantle-cell lymphoma. <i>New England Journal of Medicine</i> , 2013 , 369, 507-16	59.2	1139
284	NOTCH1, SF3B1, and TP53 mutations in fludarabine-refractory CLL patients treated with alemtuzumab: results from the CLL2H trial of the GCLLSG. <i>Blood</i> , 2013 , 122, 1266-70	2.2	57
283	ESMO Guidelines consensus conference on malignant lymphoma 2011 part 1: diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL) and chronic lymphocytic leukemia (CLL). <i>Annals of Oncology</i> , 2013 , 24, 561-76	10.3	162
282	Missing the notch in NOTCH1. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1579-80	1.9	
281	Inhibitor kinazy Brutona u chorych z nawrotowym lub opornym na leczenie chłoniakiem z komórek płaszczą Wyniki międzynarodowego, wieloośrodkowego, badania II fazy z ibrutynibem (PCI-32765) EHA Encore. <i>Acta Haematologica Polonica</i> , 2013 , 44, 314-318	0.4	1
280	The gene expression signature associated with TP53 mutation/deletion in chronic lymphocytic leukaemia is dominated by the under-expression of TP53 and other genes on chromosome 17p. <i>British Journal of Haematology</i> , 2013 , 160, 53-62	4.5	7

279	Lenalidomide reduces survival of chronic lymphocytic leukemia cells in primary cocultures by altering the myeloid microenvironment. <i>Blood</i> , 2013 , 121, 2503-11	2.2	37
278	The VEGF receptor, neuropilin-1, represents a promising novel target for chronic lymphocytic leukemia patients. <i>International Journal of Cancer</i> , 2013 , 133, 1489-96	7.5	37
277	Epigenetic upregulation of lncRNAs at 13q14.3 in leukemia is linked to the In Cis downregulation of a gene cluster that targets NF-κB. <i>PLoS Genetics</i> , 2013 , 9, e1003373	6	108
276	Overview of available p53 function tests in relation to TP53 and ATM gene alterations and chemoresistance in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1849-53	1.9	14
275	Sequential chemoimmunotherapy of fludarabine, mitoxantrone, and cyclophosphamide induction followed by alemtuzumab consolidation is effective in T-cell prolymphocytic leukemia. <i>Cancer</i> , 2013 , 119, 2258-67	6.4	44
274	Extracellular vesicles in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1826-30	1.9	12
273	Second-line therapies of patients initially treated with fludarabine and cyclophosphamide or fludarabine, cyclophosphamide and rituximab for chronic lymphocytic leukemia within the CLL8 protocol of the German CLL Study Group. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1821-2	1.9	22
272	Chronische lymphatische Leukämie - Standardtherapien und vielversprechende neue Behandlungsansätze. <i>Kliniker</i> , 2013 , 42, 350-354	0	
271	TCL1A and ATM are co-expressed in chronic lymphocytic leukemia cells without deletion of 11q. <i>Haematologica</i> , 2013 , 98, 269-73	6.6	4
270	BRAF mutations in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2013 , 54, 1177-82	1.9	36
269	Autoantigenic targets of B-cell receptors derived from chronic lymphocytic leukemias bind to and induce proliferation of leukemic cells. <i>Blood</i> , 2013 , 121, 4708-17	2.2	33
268	Telomere length in mantle cell lymphoma. <i>Blood</i> , 2013 , 121, 1184-7	2.2	15
267	TP53, SF3B1, and NOTCH1 mutations and outcome of allotransplantation for chronic lymphocytic leukemia: six-year follow-up of the GCLLSG CLL3X trial. <i>Blood</i> , 2013 , 121, 3284-8	2.2	79
266	Germline allele-specific expression of DAPK1 in chronic lymphocytic leukemia. <i>PLoS ONE</i> , 2013 , 8, e55261.7	3.7	18
265	Progranulin is a novel independent predictor of disease progression and overall survival in chronic lymphocytic leukemia. <i>PLoS ONE</i> , 2013 , 8, e72107	3.7	22
264	TP53 Mutation or Deletion and Efficacy with Single-Agent Lenalidomide in Relapsed or Refractory Chronic Lymphocytic Leukemia (CLL) (CC-5013-CLL-009 Study). <i>Blood</i> , 2013 , 122, 1638-1638	2.2	3
263	A Multicenter, Phase IV Observational Study Of Ofatumumab In Chronic Lymphocytic Leukemia (CLL): A European Research Initiative On CLL (ERIC) Study. <i>Blood</i> , 2013 , 122, 1645-1645	2.2	2
262	Role Of High-Dose Cytarabine and Total Body Irradiation Conditioning before Autologous Stem Cell Transplantation In Mantle Cell Lymphoma - A Comparison Of Nordic MCL2, HOVON 45, and European MCL Younger Trials. <i>Blood</i> , 2013 , 122, 3367-3367	2.2	4

261	Overall Survival In Early Stage Chronic Lymphocytic Leukemia Patients With Treatment Indication Due To Disease Progression: Follow-Up Data Of The CLL1 Trial Of The German CLL Study Group (GCLLSG). <i>Blood</i> , 2013 , 122, 4127-4127	2.2	4
260	Early Versus Deferred Treatment With Combined Fludarabine, Cyclophosphamide and Rituximab (FCR) Improves Event-Free Survival In Patients With High-Risk Binet Stage A Chronic Lymphocytic Leukemia [First Results Of a Randomized German-French Cooperative Phase III Trial. <i>Blood</i> , 2013 , 122, 531-531	2.2	26
259	Chemoimmunotherapy With Fludarabine (F), Cyclophosphamide (C), and Rituximab (R) (FCR) Versus Bendamustine and Rituximab (BR) In Previously Untreated and Physically Fit Patients (pts) With Advanced Chronic Lymphocytic Leukemia (CLL): Results Of a Planned Interim Analysis Of The CLL10 Trial. An International, Randomized Study Of The German CLL Study Group (GCLLSG). <i>Blood</i> , 2013 , 122, 526-526	2.2	33
258	NOTCH1 Mutation and Treatment Outcome In CLL Patients Treated With Chlorambucil (Chl) Or Ofatumumab-Chl (O-Chl): Results From The Phase III Study Complement 1 (OMB110911). <i>Blood</i> , 2013 , 122, 527-527	2.2	7
257	Head-To-Head Comparison Of Obinutuzumab (GA101) Plus Chlorambucil (Clb) Versus Rituximab Plus Clb In Patients With Chronic Lymphocytic Leukemia (CLL) and Co-Existing Medical Conditions (Comorbidities): Final Stage 2 Results Of The CLL11 Trial. <i>Blood</i> , 2013 , 122, 6-6	2.2	12
256	Telomere Length and Treatment Outcome In Chronic Lymphocytic Leukemia: Results From The CLL8 Trial. <i>Blood</i> , 2013 , 122, 671-671	2.2	3
255	Obinutuzumab (GA101) plus chlorambucil (Clb) or rituximab (R) plus Clb versus Clb alone in patients with chronic lymphocytic leukemia (CLL) and preexisting medical conditions (comorbidities): Final stage 1 results of the CLL11 (BO21004) phase III trial.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 7004-7004	2.2	18
254	Use of tumor genomic profiling to reveal mechanisms of resistance to the BTK inhibitor ibrutinib in chronic lymphocytic leukemia (CLL).. <i>Journal of Clinical Oncology</i> , 2013 , 31, 7014-7014	2.2	16
253	A comprehensive prognostic index for patients with CLL.. <i>Journal of Clinical Oncology</i> , 2013 , 31, 7015-7015	2.2	1
252	Novel Gene Mutations In Chronic Lymphocytic Leukemia: Prevalence and Clinical Implications In A Series Of 3185 Cases - Initial Results From The European Research Initiative On CLL. <i>Blood</i> , 2013 , 122, 1614-1614	2.2	
251	Microenvironmental Factors and The Role Of Tumor Necrosis Factor Receptor Type 1 (TNFR-1) In Chronic Lymphocytic Leukemia. <i>Blood</i> , 2013 , 122, 4149-4149	2.2	
250	Heterogeneity and Evolution Of DNA Methylation In Chronic Lymphocytic Leukemia. <i>Blood</i> , 2013 , 122, 1626-1626	2.2	
249	ADARB1 Is Involved In a Reduced Maturation Of The miR15a/Mir-16-1 Family In Chronic Lymphocytic Leukemia. <i>Blood</i> , 2013 , 122, 1252-1252	2.2	
248	Prognostic Assessment In Patients With Chronic Lymphocytic Leukemia (CLL) In Clinical Practice: A European Research Initiative On CLL (ERIC) Survey. <i>Blood</i> , 2013 , 122, 4156-4156	2.2	1
247	Early autologous stem cell transplantation for chronic lymphocytic leukemia: long-term follow-up of the German CLL Study Group CLL3 trial. <i>Blood</i> , 2012 , 119, 4851-9	2.2	26
246	CLL and deletion 13q14: merely the miRs?. <i>Blood</i> , 2012 , 119, 2974-5	2.2	6
245	Risk categories and refractory CLL in the era of chemoimmunotherapy. <i>Blood</i> , 2012 , 119, 4101-7	2.2	95
244	Synergy between PI3K signaling and MYC in Burkitt lymphomagenesis. <i>Cancer Cell</i> , 2012 , 22, 167-79	24.3	212

243	Final results of a multicenter phase 1 study of lenalidomide in patients with relapsed or refractory chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2012 , 53, 417-23	1.9	51
242	Cellular origin and pathophysiology of chronic lymphocytic leukemia. <i>Journal of Experimental Medicine</i> , 2012 , 209, 2183-98	16.6	178
241	High-resolution genomic profiling of chronic lymphocytic leukemia reveals new recurrent genomic alterations. <i>Blood</i> , 2012 , 120, 4783-94	2.2	156
240	Genetics in chronic lymphocytic leukemia 2012 , 16-24		
239	Minimal residual disease quantification is an independent predictor of progression-free and overall survival in chronic lymphocytic leukemia: a multivariate analysis from the randomized GCLLSG CLL8 trial. <i>Journal of Clinical Oncology</i> , 2012 , 30, 980-8	2.2	334
238	miRNA-130a targets ATG2B and DICER1 to inhibit autophagy and trigger killing of chronic lymphocytic leukemia cells. <i>Cancer Research</i> , 2012 , 72, 1763-72	10.1	161
237	Bendamustine in combination with rituximab for previously untreated patients with chronic lymphocytic leukemia: a multicenter phase II trial of the German Chronic Lymphocytic Leukemia Study Group. <i>Journal of Clinical Oncology</i> , 2012 , 30, 3209-16	2.2	332
236	Quantitative DNA methylation analysis identifies a single CpG dinucleotide important for ZAP-70 expression and predictive of prognosis in chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , 2012 , 30, 2483-91	2.2	100
235	Autologous and allogeneic stem-cell transplantation for transformed chronic lymphocytic leukemia (Richter's syndrome): A retrospective analysis from the chronic lymphocytic leukemia subcommittee of the chronic leukemia working party and lymphoma working party of the European Group for Blood and Marrow Transplantation. <i>Journal of Clinical Oncology</i> , 2012 , 30, 2211-7	2.2	83
234	Assessment of p53 Functionality in Chronic Lymphocytic Leukemia by Different Assays; An Eric-Wide Approach.. <i>Blood</i> , 2012 , 120, 2872-2872	2.2	1
233	Sequential High Dose Immuno-Chemotherapy Followed by Autologous Peripheral Blood Stem Cell Transplantation for Patients with Untreated Primary Central Nervous System Lymphoma - a Multicentre Study by the Collaborative PCNSL Study Group Freiburg. <i>Blood</i> , 2012 , 120, 302-302	2.2	14
232	Updated Interim Results of the Safety and Efficacy of Different Lenalidomide Starting Dose Regimens in Patients with Relapsed or Refractory (rel/ref) Chronic Lymphocytic Leukemia (CLL) (CC-5013-CLL-009 Study). <i>Blood</i> , 2012 , 120, 3925-3925	2.2	2
231	Gene Mutations and Treatment Outcome in Chronic Lymphocytic Leukemia: Results From the CLL8 Trial. <i>Blood</i> , 2012 , 120, 433-433	2.2	6
230	Extended Follow up of the CLL8 Protocol, a Randomized Phase-III Trial of the German CLL Study Group (GCLLSG) Comparing Fludarabine and Cyclophosphamide (FC) to FC Plus Rituximab (FCR) for Previously Untreated Patients with Chronic Lymphocytic Leukemia (CLL): Results On Survival, Response, Toxicity, Quality of Life, and Minimal Residual Disease. <i>Journal of Clinical Oncology</i> , 2012 , 30, 2211-7	2.2	14
229	Alemtuzumab Plus Oral Dexamethasone, Followed by Alemtuzumab Maintenance or Allogeneic Transplantation in Ultra High-Risk CLL: Updated Results From a Phase II Study of the Gcllsg and fcgcll/MW. <i>Blood</i> , 2012 , 120, 716-716	2.2	1
228	SF3B1, NOTCH1 and TP53 Mutations Do Not Affect the Outcome of Allogeneic Hematopoietic Stem Cell Transplantation (HSCT) for Chronic Lymphocytic Leukemia (CLL): 6-Year Follow-up of the Gcllsg CLL3X Trial. <i>Blood</i> , 2012 , 120, 966-966	2.2	1
227	Lenalidomide Reduces Survival of Chronic Lymphocytic Leukemia Cells in Primary Co-Cultures by Altering the Myeloid Microenvironment. <i>Blood</i> , 2012 , 120, 3894-3894	2.2	
226	Autoantigenic Targets of B-Cell Receptors (BCR) Derived From Chronic Lymphocytic Leukemias Bind to and Induce Proliferation of Leukemic Cells.. <i>Blood</i> , 2012 , 120, 2884-2884	2.2	

225	TP53 Mutation Is an Independent Predictor of Poor Survival in Untreated Patients with CD20+ Aggressive B-Cell Lymphoma: Analysis within the Ricover-60 Trial. <i>Blood</i> , 2012 , 120, 546-546	2.2	
224	Telomere Length in Mantle Cell Lymphoma.. <i>Blood</i> , 2012 , 120, 2509-2509	2.2	
223	IL4 and CD40L Prevent Apoptosis of Chronic Lymphocytic Leukemia Cells Via Intracellular pSTAT6 and NFkB Signaling and Not Via Receptor Kinetics. <i>Blood</i> , 2012 , 120, 3893-3893	2.2	
222	Microrna Expression in Fludarabine-Refractory CLL Implicates Independent Mechanisms of Resistance and Is Associated with Response and Progression Free Survival After Alemtuzumab Treatment: Results From the CLL2H Trial.. <i>Blood</i> , 2012 , 120, 2874-2874	2.2	
221	The Use of Chemoimmunotherapy Improves the Outcome of Patients with Chronic Lymphocytic Leukaemia (CLL) ▯ Metaanalysis of Five Trials of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2012 , 120, 3936-3936	2.2	
220	DNA damage-induced transcriptional program in CLL: biological and diagnostic implications for functional p53 testing. <i>Blood</i> , 2011 , 117, 1622-32	2.2	33
219	Immune modulatory agents in hematopoietic malignancies. <i>Cancer Treatment Reviews</i> , 2011 , 37 Suppl 1, S2-7	14.4	1
218	Bendamustine combined with rituximab in patients with relapsed and/or refractory chronic lymphocytic leukemia: a multicenter phase II trial of the German Chronic Lymphocytic Leukemia Study Group. <i>Journal of Clinical Oncology</i> , 2011 , 29, 3559-66	2.2	315
217	Chemoimmunotherapy with O-FC in previously untreated patients with chronic lymphocytic leukemia. <i>Blood</i> , 2011 , 117, 6450-8	2.2	107
216	Genetics of chronic lymphocytic leukemia. <i>Clinics in Laboratory Medicine</i> , 2011 , 31, 649-58, ix	2.1	6
215	P53 and microRNAs in chronic lymphocytic leukemia. <i>Journal of Nucleic Acids Investigation</i> , 2011 , 2, 8		1
214	Limited clinical relevance of imaging techniques in the follow-up of patients with advanced chronic lymphocytic leukemia: results of a meta-analysis. <i>Blood</i> , 2011 , 117, 1817-21	2.2	32
213	Inflammatory cytokines and signaling pathways are associated with survival of primary chronic lymphocytic leukemia cells in vitro: a dominant role of CCL2. <i>Haematologica</i> , 2011 , 96, 408-16	6.6	63
212	Nurse-like cells show deregulated expression of genes involved in immunocompetence. <i>British Journal of Haematology</i> , 2011 , 154, 349-56	4.5	26
211	Clinical efficacy of immunochemotherapy with fludarabine, epirubicin and rituximab in the treatment for chronic lymphocytic leukaemia and prolymphocytic leukaemia. <i>European Journal of Haematology</i> , 2011 , 87, 426-33	3.8	20
210	A novel Fc-engineered monoclonal antibody to CD37 with enhanced ADCC and high proapoptotic activity for treatment of B-cell malignancies. <i>Blood</i> , 2011 , 118, 4159-68	2.2	85
209	Mutation analysis of the TNFAIP3 (A20) tumor suppressor gene in CLL. <i>International Journal of Cancer</i> , 2011 , 128, 1747-50	7.5	15
208	Importance of genetics in chronic lymphocytic leukemia. <i>Blood Reviews</i> , 2011 , 25, 131-7	11.1	66

207	Novel targeted treatment strategies for refractory chronic lymphocytic leukaemia. <i>Therapeutic Advances in Hematology</i> , 2011 , 2, 249-65	5.7	3
206	Treating chronic lymphocytic leukemia with thalidomide and lenalidomide. <i>Expert Opinion on Pharmacotherapy</i> , 2011 , 12, 2857-64	4	13
205	MicroRNA-130a Targets ATG2B, AGO4 and DICER1, Inhibits Autophagy and Induces Cell Death in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2011 , 118, 1768-1768	2.2	1
204	Minimal Residual Disease (MRD) Re-Growth Kinetics Are An Independent Predictor for Progression Free Survival (PFS) in Chronic Lymphocytic Leukemia (CLL) and Are Related to Biologically Defined CLL-Subgroups Results From the CLL8 Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2011 , 118, 4577-4577	2.2	4
203	Correlations Between Ofatumumab Exposure and Treatment Outcomes for Patients with Chronic Lymphocytic Leukemia (CLL) Treated with Frontline Ofatumumab, Fludarabine, and Cyclophosphamide Chemoimmunotherapy. <i>Blood</i> , 2011 , 118, 1793-1793	2.2	1
202	Alemtuzumab Plus Oral Dexamethasone, Followed by Alemtuzumab Maintenance or Allogeneic Transplantation in Ultra High-Risk CLL: Interim Analysis of a Phase II Study of the GCLLSG and fcgcll/MW. <i>Blood</i> , 2011 , 118, 2854-2854	2.2	2
201	Interim Results for the Safety and Efficacy of Different Lenalidomide Starting Dose Regimens in Subjects with Relapsed or Refractory Chronic Lymphocytic Leukemia (CC-5013-CLL-009 Study). <i>Blood</i> , 2011 , 118, 2859-2859	2.2	1
200	Cyclophosphamide, Adriamycin, Vincristine and Prednisone Plus Rituximab (CHOP-R) in Fludarabine (F) Refractory Chronic Lymphocytic Leukemia (CLL) or CLL with Autoimmune Cytopenia (AIC) or Richter's Transformation (RT): Final Analysis of a Phase II Study of the German CLL Study Group. <i>Blood</i> , 2011 , 118, 2860-2860	2.2	2
199	Prediction of Poor Outcome in CLL Patients Treated with FCR (Fludarabine, Cyclophosphamide, Rituximab) in the CLL8 Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2011 , 118, 977-977	2.2	2
198	Uncovering Thalidomide Mechanism of Action: Neuropilin 1 Represents the Target of Antiangiogenic and Immunomodulatory Effect in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2011 , 118, 4608-4608	2.2	2
197	The Toll-Like Receptor-Like Molecule CD180 and Soluble CD14 Transmit Survival Signals in B-Cell Chronic Lymphocytic Leukemia Cells Presumably by Acting As Co-Receptors,. <i>Blood</i> , 2011 , 118, 3883-3883	2.2	2
196	Prognostic Impact of Biallelic Versus Monoallelic Deletion 13q and the Proportion of Cells with Deletion 13q within the CLL8 Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2011 , 118, 2841-2841	2.2	2
195	Reduced c-FOS Is Associated with Poor Prognosis and Clinical Course in Chronic Lymphocytic Leukemia. <i>Blood</i> , 2011 , 118, 2433-2433	2.2	
194	Role of Microenvironment-Associated Chemokines and Cytokines for Binet Stage A CLL Patients Included in a Prospective Trial (CLL1 trial) of the German CLL Study Group (GCLLSG): sll2Ralpha Is An Independent Predictor of Progression-Free Survival (PFS),. <i>Blood</i> , 2011 , 118, 3869-3869	2.2	
193	SNP-Arrays Provide New Insights Into the Pathogenesis of Richter Syndrome (RS). <i>Blood</i> , 2011 , 118, 263-263		
192	Proposal of A Prognostic Score for Previously Untreated Patients with Chronic Lymphocytic Leukemia Based on An Overall Survival Analysis of Three German CLL Study Group Phase III Trials. <i>Blood</i> , 2011 , 118, 2831-2831	2.2	
191	Lenalidomide treatment of chronic lymphocytic leukaemia patients reduces regulatory T cells and induces Th17 T helper cells. <i>British Journal of Haematology</i> , 2010 , 148, 948-50	4.5	50
190	SYK carries no activating point mutations in patients with chronic lymphocytic leukaemia (CLL). <i>British Journal of Haematology</i> , 2010 , 150, 633-6	4.5	9

189	From pathogenesis to treatment of chronic lymphocytic leukaemia. <i>Nature Reviews Cancer</i> , 2010 , 10, 37-50	31.3	438
188	Ofatumumab as single-agent CD20 immunotherapy in fludarabine-refractory chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , 2010 , 28, 1749-55	2.2	483
187	Immunoglobulin heavy chain variable gene usage and (super)-antigen drive in chronic lymphocytic leukemia. <i>Clinical Cancer Research</i> , 2010 , 16, 373-5	12.9	3
186	Understanding and managing ultra high-risk chronic lymphocytic leukemia. <i>Hematology American Society of Hematology Education Program</i> , 2010 , 2010, 481-8	3.1	96
185	More (on) prognostic factors in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2010 , 51, 5-6	1.9	1
184	Chronic lymphocytic leukemia: new concepts for future therapy. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2010 , 10, 369-78	2	9
183	Moving from prognostic to predictive factors in chronic lymphocytic leukaemia (CLL). <i>Best Practice and Research in Clinical Haematology</i> , 2010 , 23, 71-84	4.2	51
182	TP53 mutation and survival in chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , 2010 , 28, 4473-82	2.2	430
181	Allogeneic stem cell transplantation provides durable disease control in poor-risk chronic lymphocytic leukemia: long-term clinical and MRD results of the German CLL Study Group CLL3X trial. <i>Blood</i> , 2010 , 116, 2438-47	2.2	240
180	Protein expression analysis of chronic lymphocytic leukemia defines the effect of genetic aberrations and uncovers a correlation of CDK4, P27 and P53 with hierarchical risk. <i>Haematologica</i> , 2010 , 95, 1880-8	6.6	4
179	Gene expression factors as predictors of genetic risk and survival in chronic lymphocytic leukemia. <i>Haematologica</i> , 2010 , 95, 102-9	6.6	24
178	Spleen tyrosine kinase inhibition prevents chemokine- and integrin-mediated stromal protective effects in chronic lymphocytic leukemia. <i>Blood</i> , 2010 , 115, 4497-506	2.2	152
177	Soluble CD14 is a novel monocyte-derived survival factor for chronic lymphocytic leukemia cells, which is induced by CLL cells in vitro and present at abnormally high levels in vivo. <i>Blood</i> , 2010 , 116, 4223-30	2.2	56
176	Highlights der 51. Jahrestagung der American Society of Hematology (ASH) 2009. <i>Onkopipeline</i> , 2010 , 3, 52-61		
175	High-throughput detection of nuclear factor-kappaB activity using a sensitive oligo-based chemiluminescent enzyme-linked immunosorbent assay. <i>International Journal of Cancer</i> , 2010 , 127, 404-11	7.5	21
174	CD5+ B cells from individuals with systemic lupus erythematosus express granzyme B. <i>European Journal of Immunology</i> , 2010 , 40, 2060-9	6.1	40
173	Alternating Courses of 3x CHOP and 3x DHAP Plus Rituximab Followed by a High Dose ARA-C Containing Myeloablative Regimen and Autologous Stem Cell Transplantation (ASCT) Is Superior to 6 Courses CHOP Plus Rituximab Followed by Myeloablative Radiochemotherapy and ASCT In Mantle Cell Lymphoma: Results of the MCL Younger Trial of the European Mantle Cell Lymphoma	2.2	21
172	Final Results of the Phase I Study of Lenalidomide In Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia (CLL-001 Study). <i>Blood</i> , 2010 , 116, 1376-1376	2.2	1

171	Allogeneic Stem Cell Transplantation Can Overcome the Adverse Prognostic Impact of TP53 Mutation In Chronic Lymphocytic Leukemia (CLL): Results From the GCLLSG CLL3x Trial. <i>Blood</i> , 2010 , 116, 2357-2357	2.2	1
170	Genetics of Patients with F-Refractory CLL or Early Relapse After FC or FCR: Results From the CLL8 Trial of the GCLLSG. <i>Blood</i> , 2010 , 116, 2427-2427	2.2	9
169	High-Resolution SNP-Array Profiling of Chronic Lymphocytic Leukemia. <i>Blood</i> , 2010 , 116, 50-50	2.2	1
168	Interim Analysis of EFC6663, a Multicenter Phase 2 Study of Alvocidib (flavopiridol), Demonstrates Clinical Responses Among Patients with Fludarabine Refractory CLL. <i>Blood</i> , 2010 , 116, 58-58	2.2	5
167	Serum Factors Predict Therapeutic Outcome In Patients with Chronic Lymphocytic Leukemia Treated In the CLL8 Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2010 , 116, 918-918	2.2	2
166	Subcutaneous Alemtuzumab Combined with Oral Dexamethasone, Followed by Alemtuzumab Maintenance or Allo-SCT In CLL with 17p- or Refractory to Fludarabine [Interim Analysis of the CLL20 Trial of the GCLLSG and FCGCLL/MW. <i>Blood</i> , 2010 , 116, 920-920	2.2	12
165	Final Analysis From the International Trial of Single-Agent Ofatumumab In Patients with Fludarabine-Refractory Chronic Lymphocytic Leukemia. <i>Blood</i> , 2010 , 116, 921-921	2.2	6
164	Refractory chronic lymphocytic leukemia - new therapeutic strategies. <i>Oncotarget</i> , 2010 , 1, 472-482	3.3	17
163	Refractory chronic lymphocytic leukemia--new therapeutic strategies. <i>Oncotarget</i> , 2010 , 1, 472-82	3.3	11
162	IGHV-Mutation Status, IGHV-Gene Usage and Chromosomal Aberrations In CLL: Pooled Analysis within First-Line Clinical Trials of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2010 , 116, 3609-3609	2.2	
161	Exceptional In Vitro Activity of CD37 Antibodies In CLL. <i>Blood</i> , 2010 , 116, 2460-2460	2.2	
160	Induction Therapy with Idarubicin and Etoposide Combined with Sequential or Concurrent Azacitidine In Patients with High-Risk Acute Myeloid Leukemia: Pilot-Phase of the AMLSG 12-09 Study. <i>Blood</i> , 2010 , 116, 2184-2184	2.2	
159	Maturation of the miR15a/miR16-1 Family Is Impaired In Chronic Lymphocytic Leukaemia. <i>Blood</i> , 2010 , 116, 53-53	2.2	
158	Chemoimmunotherapy in chronic lymphocytic leukemia. <i>Clinical Advances in Hematology and Oncology</i> , 2010 , 8, 867-8	0.6	
157	BCL2-938C>A polymorphism and disease progression in chronic lymphocytic leukemia. <i>Leukemia and Lymphoma</i> , 2009 , 50, 1837-42	1.9	12
156	miR-34a as part of the resistance network in chronic lymphocytic leukemia. <i>Blood</i> , 2009 , 113, 3801-8	2.2	229
155	Fatal <i>Ophiostoma piceae</i> infection in a patient with acute lymphoblastic leukaemia. <i>Journal of Medical Microbiology</i> , 2009 , 58, 381-385	3.2	12
154	A novel paradigm to trigger apoptosis in chronic lymphocytic leukemia. <i>Cancer Research</i> , 2009 , 69, 8977-86.1	3.6	51

153	Danish CLL2-Study revisited: FISH on a cohort with a 20-yr follow-up confirms the validity of the hierarchical model of genomic aberrations in chronic lymphocytic leukaemia. <i>European Journal of Haematology</i> , 2009 , 83, 156-8	3.8	5
152	Chronic lymphocytic leukemia and 13q14: miRs and more. <i>Leukemia and Lymphoma</i> , 2009 , 50, 502-5	1.9	35
151	Detailed analysis of p53 pathway defects in fludarabine-refractory chronic lymphocytic leukemia (CLL): dissecting the contribution of 17p deletion, TP53 mutation, p53-p21 dysfunction, and miR34a in a prospective clinical trial. <i>Blood</i> , 2009 , 114, 2589-97	2.2	232
150	Treatment resistance in chronic lymphocytic leukemia: the role of the p53 pathway. <i>Leukemia and Lymphoma</i> , 2009 , 50, 510-3	1.9	32
149	Therapy with the FCR regimen does not overcome chronic lymphocytic leukemia biology: aberrant p53 expression predicts response and survival. <i>Leukemia and Lymphoma</i> , 2009 , 50, 1559-61	1.9	2
148	Subcutaneous alemtuzumab in fludarabine-refractory chronic lymphocytic leukemia: clinical results and prognostic marker analyses from the CLL2H study of the German Chronic Lymphocytic Leukemia Study Group. <i>Journal of Clinical Oncology</i> , 2009 , 27, 3994-4001	2.2	230
147	TP53 Mutations and Outcome After Fludarabine and Cyclophosphamide (FC) or FC Plus Rituximab (FCR) in the CLL8 Trial of the GCLLSG.. <i>Blood</i> , 2009 , 114, 1267-1267	2.2	6
146	Bendamustine Combined with Rituximab (BR) in First-Line Therapy of Advanced CLL: A Multicenter Phase II Trial of the German CLL Study Group (GCLLSG).. <i>Blood</i> , 2009 , 114, 205-205	2.2	22
145	Ofatumumab Combined with Fludarabine and Cyclophosphamide (O-FC) Shows High Activity in Patients with Previously Untreated Chronic Lymphocytic Leukemia (CLL): Results From a Randomized, Multicenter, International, Two-Dose, Parallel Group, Phase II Trial.. <i>Blood</i> , 2009 , 114, 207-207	2.2	14
144	Chemoimmuno-Therapy with Fludarabine, Cyclophosphamide and Alemtuzumab (FC-Cam) in Patients with Relapsed or Genetic High-Risk CLL: Final Analysis of the CLL2L Trial of the German CLL Study Group.. <i>Blood</i> , 2009 , 114, 209-209	2.2	4
143	Identification of RHAMM-Derived CD8+ Restricted, Heteroclitical, Cryptic Epitope R9Y as a Promising Target for Immunotherapy of Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2009 , 114, 3034-3034	2.2	1
142	Correlation Between Serum Ofatumumab Concentrations, Baseline Patient Characteristics and Clinical Outcomes in Patients with Fludarabine-Refractory Chronic Lymphocytic Leukemia (CLL) Treated with Single-Agent Ofatumumab.. <i>Blood</i> , 2009 , 114, 3433-3433	2.2	4
141	First-Line Treatment with Fludarabine (F), Cyclophosphamide (C), and Rituximab (R) (FCR) Improves Overall Survival (OS) in Previously Untreated Patients (pts) with Advanced Chronic Lymphocytic Leukemia (CLL): Results of a Randomized Phase III Trial On Behalf of An International Group of Investigators and the German CLL Study Group. <i>Blood</i> , 2009 , 114, 535-535	2.2	31
140	Early Autologous Stem Cell Transplantation (autoSCT) May Overcome the Adverse Impact of Del 11q- in Poor-Risk Chronic Lymphocytic Leukemia (CLL): Results From the GCLLSG CLL3 Trial.. <i>Blood</i> , 2009 , 114, 879-879	2.2	1
139	Comprehensive Protein Expression Analysis of Chronic Lymphocytic Leukemia Uncovers a Correlation of CDK4, P27 and P53 with Hierarchical Risk and Suggests a Dichotomy of Apoptosis and Proliferation.. <i>Blood</i> , 2009 , 114, 1255-1255	2.2	
138	A Novel Paradigm to Trigger Apoptosis in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2009 , 114, 731-731	2.2	
137	Identification and Characterization of Peptide Ligands for Stereotyped Subset and Non-Subset B-Cell Receptors of Patients with M- and U-CLL.. <i>Blood</i> , 2009 , 114, 4369-4369	2.2	
136	In Vitro Activity of the Type II Anti-CD20 Antibody GA101 in Refractory, Genetic High-Risk CLL.. <i>Blood</i> , 2009 , 114, 2379-2379	2.2	1

135	Detection of NFB Activity in Primary CLL Cells Using a Sensitive Oligo-Based Chemiluminescent ELISA.. <i>Blood</i> , 2009 , 114, 4709-4709	2.2	
134	In-Vitro Effect of the Small Molecule Thrombopoietin (TPO) Receptor Agonist Eltrombopag in Chronic Lymphocytic Leukemia (CLL).. <i>Blood</i> , 2009 , 114, 4408-4408	2.2	
133	Neuropilin-1 [Novel, Promising Target for Chronic Lymphocytic Leukemia patients.. <i>Blood</i> , 2009 , 114, 4392-4392	2.2	0
132	Delineation of distinct tumour profiles in mantle cell lymphoma by detailed cytogenetic, interphase genetic and morphological analysis. <i>British Journal of Haematology</i> , 2008 , 142, 538-50	4.5	37
131	First demonstration of leukemia imaging with the proliferation marker 18F-fluorodeoxythymidine. <i>Journal of Nuclear Medicine</i> , 2008 , 49, 1756-62	8.9	55
130	Molecular diagnostics in chronic lymphocytic leukemia - pathogenetic and clinical implications. <i>Leukemia and Lymphoma</i> , 2008 , 49, 864-73	1.9	36
129	Monoallelic TP53 inactivation is associated with poor prognosis in chronic lymphocytic leukemia: results from a detailed genetic characterization with long-term follow-up. <i>Blood</i> , 2008 , 112, 3322-9	2.2	322
128	Allogeneic hematopoietic stem-cell transplantation for chronic lymphocytic leukemia with 17p deletion: a retrospective European Group for Blood and Marrow Transplantation analysis. <i>Journal of Clinical Oncology</i> , 2008 , 26, 5094-100	2.2	139
127	The MDM2 -309 T/G promoter single nucleotide polymorphism does not alter disease characteristics in chronic lymphocytic leukemia. <i>Haematologica</i> , 2008 , 93, 1111-3	6.6	31
126	Chronic lymphocytic leukemia and treatment resistance in cancer: the role of the p53 pathway. <i>Cell Cycle</i> , 2008 , 7, 3810-4	4.7	66
125	Short telomeres are associated with genetic complexity, high-risk genomic aberrations, and short survival in chronic lymphocytic leukemia. <i>Blood</i> , 2008 , 111, 2246-52	2.2	117
124	Response: Or both?. <i>Blood</i> , 2008 , 111, 5756-5757	2.2	1
123	Genomic aberrations in mantle cell lymphoma detected by interphase fluorescence in situ hybridization. Incidence and clinicopathological correlations. <i>Haematologica</i> , 2008 , 93, 680-7	6.6	37
122	Autoimmune conditions and chronic infections in chronic lymphocytic leukemia patients at diagnosis are associated with unmutated IgVH genes. <i>Haematologica</i> , 2008 , 93, 1912-6	6.6	20
121	Identification of Genes Associated with Resistance and Response in Vivo to Therapy with Rituximab, Fludarabine and Cyclophosphamide in Patients with Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2008 , 112, 1622-1622	2.2	2
120	Biologic and Clinical Markers for Outcome after Fludarabine (F) or F Plus Cyclophosphamide (FC) - Comprehensive Analysis of the CLL4 Trial of the GCLLSG.. <i>Blood</i> , 2008 , 112, 2089-2089	2.2	4
119	Preliminary Results of a Phase 1/2, Multi-Center, Open-Label Study (CLL- 001) Investigating a Stepwise Dose-Escalation Schedule of Lenalidomide in Relapsed or Refractory Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2008 , 112, 2104-2104	2.2	4
118	Standardized MRD Flow and ASO IGH RQ-PCR for MRD Quantification in CLL Patients after Rituximab-Containing Immunochemotherapy [Comparative Analysis in 574 Samples from the Randomized GCLLSG CLL8 Trial. <i>Blood</i> , 2008 , 112, 3139-3139	2.2	1

117	Induction of Apoptosis in CLL by Peptides Binding the B-Cell Antigen Receptor in Vitro. <i>Blood</i> , 2008 , 112, 3151-3151	2.2	1
116	Peptide Vaccination Induces Dynamic Changes in CD4+ and CD8+ T Cell Subsets: Report on the First Peptide Vaccination Trial in Patients with Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2008 , 112, 3159-3159	2.2	2
115	Effective Targeting of the PI3-K Pathway in CLL with NVP-BEZ235, a Novel Orally Available Dual PI3K/mTOR Inhibitor. <i>Blood</i> , 2008 , 112, 3166-3166	2.2	1
114	Chemoimmuno-Therapy with Fludarabine, Cyclophosphamide and Alemtuzumab in Patients with Relapsed/Refractory CLL: Interim Analysis of the CLL2L Trial of the German CLL Study Group. <i>Blood</i> , 2008 , 112, 3170-3170	2.2	2
113	Immunochemotherapy with Fludarabine (F), Cyclophosphamide (C), and Rituximab (R) (FCR) Versus Fludarabine and Cyclophosphamide (FC) Improves Response Rates and Progression-Free Survival (PFS) of Previously Untreated Patients (pts) with Advanced Chronic Lymphocytic Leukemia (CLL). <i>Blood</i> , 2008 , 112, 325-325	2.2	78
112	Quantitative MRD Assessments Predict Progression Free Survival in CLL Patients Treated with Fludarabine and Cyclophosphamide with or without Rituximab: A Prospective Analysis in 471 Patients from the Randomized GCLLSG CLL8 Trial. <i>Blood</i> , 2008 , 112, 326-326	2.2	12
111	Ofatumumab (HuMax-CD20), a Novel CD20 Monoclonal Antibody, Is An Active Treatment for Patients with CLL Refractory to Both Fludarabine and Alemtuzumab or Bulky Fludarabine-Refractory Disease: Results from the Planned Interim Analysis of An International Pivotal Trial. <i>Blood</i> , 2008 , 112, 328-328	2.2	21
110	Subcutaneous Alemtuzumab (Campath) in Fludarabine-Refractory CLL: Final Results of the CLL2H Trial of the GCLLSG and Comprehensive Analysis of Prognostic Markers. <i>Blood</i> , 2008 , 112, 329-329	2.2	2
109	Bendamustine in Combination with Rituximab (BR) for Patients with Relapsed Chronic Lymphocytic Leukemia (CLL): A Multicentre Phase II Trial of the German CLL Study Group (GCLLSG). <i>Blood</i> , 2008 , 112, 330-330	2.2	29
108	Prognostic Factors for Outcome of Nonmyeloablative Allogeneic Stem Cell Transplantation (NST) in Poor-Risk Chronic Lymphocytic Leukemia (CLL): Final Results from a Prospective Multicenter Trial (GCLLSG CLL3X study). <i>Blood</i> , 2008 , 112, 565-565	2.2	4
107	Genomic Aberrations, VH Mutation Status and Outcome after Fludarabine and Cyclophosphamide (FC) or FC Plus Rituximab (FCR) in the CLL8 Trial. <i>Blood</i> , 2008 , 112, 781-781	2.2	16
106	Differential Diagnosis, Staging and Prognostic Factors 2008 , 103-119		2
105	Mir-34a as Part of the Chemotherapy Resistance Network in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2008 , 112, 1209-1209	2.2	1
104	The Response to DNA Damage in CLL Cells Is Partly Determined by the Type of TP53 Mutation and Genomic Aberrations. <i>Blood</i> , 2008 , 112, 3119-3119	2.2	
103	Molecular and Immunological Effects of Thalidomide in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2008 , 112, 2092-2092	2.2	
102	Epimutation of the Tumor Suppressor Mechanism in 13q14.3 Involves Monoallelic Expression, Non-Coding RNA Genes and Deregulation of NFkB Signalling. <i>Blood</i> , 2008 , 112, 783-783	2.2	
101	17p Deletion in CLL: Detailed Analysis of TP53 Mutations, Alternative Mechanisms of p53 Inactivation, Clone Size and Clonal Evolution. <i>Blood</i> , 2008 , 112, 782-782	2.2	1
100	In-Vitro Treatment with the AKT-Inhibitor GSK 690693 Induces Cell Death in Lymphoma Cell Lines and in Primary CLL Cells and Is Followed by Caspase-3 Activation and Cytochrome C Release.. <i>Blood</i> , 2008 , 112, 1589-1589	2.2	

99	TP53 Abnormalities in Chronic Lymphocytic Leukemia Exhibit a Disease Specific Profile: Meta-Analysis of 270 Mutations.. <i>Blood</i> , 2008 , 112, 2077-2077	2.2	
98	Quantitative Gene Expression Analysis of Surrogate Markers for Genetic Risk Groups and Survival in CLL. <i>Blood</i> , 2008 , 112, 4170-4170	2.2	
97	Advances in the use of alemtuzumab in CLL. <i>Clinical Advances in Hematology and Oncology</i> , 2008 , 6, 23-40.6		1
96	Update on Genomic Profiling in Chronic Lymphocytic Leukemia. <i>Clinical Leukemia</i> , 2007 , 1, 217-222		
95	Genetics and risk-stratified approach to therapy in chronic lymphocytic leukemia. <i>Best Practice and Research in Clinical Haematology</i> , 2007 , 20, 439-53	4.2	58
94	High rate of centrosome aberrations and correlation with proliferative activity in patients with untreated B-cell chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , 2007 , 121, 978-83	7.5	11
93	Overexpression of the paternally expressed gene 10 (PEG10) from the imprinted locus on chromosome 7q21 in high-risk B-cell chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , 2007 , 121, 1984-93	7.5	50
92	Proposals for standardized protocols for cytogenetic analyses of acute leukemias, chronic lymphocytic leukemia, chronic myeloid leukemia, chronic myeloproliferative disorders, and myelodysplastic syndromes. <i>Genes Chromosomes and Cancer</i> , 2007 , 46, 494-9	5	61
91	Reduced-intensity conditioning followed by T-cell depleted allogeneic stem cell transplantation for patients with chronic myeloid leukaemia and minimal residual disease at the time of transplant: high risk of molecular relapse. <i>British Journal of Haematology</i> , 2007 , 136, 127-30	4.5	5
90	Quantitative gene expression deregulation in mantle-cell lymphoma: correlation with clinical and biologic factors. <i>Journal of Clinical Oncology</i> , 2007 , 25, 2770-7	2.2	48
89	Clonal evolution in chronic lymphocytic leukemia: acquisition of high-risk genomic aberrations associated with unmutated VH, resistance to therapy, and short survival. <i>Haematologica</i> , 2007 , 92, 1242-5	6.6	179
88	p53 Inactivation in CLL: Pattern of 110 TP53 Mutations.. <i>Blood</i> , 2007 , 110, 2064-2064	2.2	3
87	Bendamustine in Combination with Rituximab (BR) for Patients with Relapsed Chronic Lymphocytic Leukemia (CLL): A Multicentre Phase II Trial of the German CLL Study Group (GCLLSG).. <i>Blood</i> , 2007 , 110, 3106-3106	2.2	10
86	Subcutaneous Alemtuzumab (MabCampath) in Fludarabine-Refractory CLL (CLL2H Trial of the GCLLSG).. <i>Blood</i> , 2007 , 110, 3120-3120	2.2	10
85	Thalidomide Alone and in Combination with Fludarabine Exerts Distinct Molecular and Antileukemic Effects in B-Cell Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2007 , 110, 3124-3124	2.2	1
84	Allogeneic Hematopoietic Cell Transplantation for Chronic Lymphocytic Leukemia (CLL) with 17p Deletion: A Retrospective EBMT Analysis.. <i>Blood</i> , 2007 , 110, 47-47	2.2	1
83	Prospective Evaluation of Prognostic Parameters in Early Stage Chronic Lymphocytic Leukemia (CLL): Results of the CLL1-Protocol of the German CLL Study Group (GCLLSG).. <i>Blood</i> , 2007 , 110, 625-625	2.2	10
82	Prognostic Assessment of Three Single Nucleotide Polymorphisms (BCL2B38>A, MTHFR 677C>T, GNAS1 T393C) in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2007 , 110, 2080-2080	2.2	

81	Gene Expression Signature of B-Cell Chronic Lymphocytic Leukemia with Trisomy 12.. <i>Blood</i> , 2007 , 110, 2079-2079	2.2	
80	In-Vitro Treatment of Lymphoma Cell Lines with BAY 43-9006 (Sorafenib) Is Followed by Typical Features of Apoptosis and Down-Regulation of MCL-1 Pointing to an Efficacy of Sorafenib in the Treatment of Lymphoma.. <i>Blood</i> , 2007 , 110, 1392-1392	2.2	
79	The Receptor for Hyaluronic Acid Mediated Motility (RHAMM): Characterization as an Immunotherapeutical Target in Chronic Lymphocytic Leukemia (CLL) and First Results of RHAMM-Derived Peptide Vaccination Trial.. <i>Blood</i> , 2007 , 110, 2051-2051	2.2	
78	BCRs of Mutated and Unmutated CLL Bind Peptides with Distinct Sequence Features: Evidence from Phage Display Libraries.. <i>Blood</i> , 2007 , 110, 1118-1118	2.2	
77	Uncovering the Epigenetic Pathomechanism in 13q14.. <i>Blood</i> , 2007 , 110, 487-487	2.2	
76	Autoimmune or Chronic Infectious Disease in B-CLL at Diagnosis: Association with Unmutated VH Gene Status and Unfavorable Cytogenetics.. <i>Blood</i> , 2007 , 110, 3092-3092	2.2	
75	Risk stratification in chronic lymphocytic leukemia. <i>Seminars in Oncology</i> , 2006 , 33, 186-94	5.5	62
74	Ala228 variant of trail receptor 1 affecting the ligand binding site is associated with chronic lymphocytic leukemia, mantle cell lymphoma, prostate cancer, head and neck squamous cell carcinoma and bladder cancer. <i>International Journal of Cancer</i> , 2006 , 118, 1831-5	7.5	37
73	Allelic silencing at the tumor-suppressor locus 13q14.3 suggests an epigenetic tumor-suppressor mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 7741-6	11.5	47
72	Molecular imaging of proliferation in malignant lymphoma. <i>Cancer Research</i> , 2006 , 66, 11055-61	10.1	158
71	Additional genetic high-risk features such as 11q deletion, 17p deletion, and V3-21 usage characterize discordance of ZAP-70 and VH mutation status in chronic lymphocytic leukemia. <i>Journal of Clinical Oncology</i> , 2006 , 24, 969-75	2.2	157
70	Antagonizing inactivated tumor suppressor genes and activated oncogenes by a versatile transgenesis system: application in mantle cell lymphoma. <i>FASEB Journal</i> , 2006 , 20, 1188-90	0.9	21
69	Fludarabine plus cyclophosphamide versus fludarabine alone in first-line therapy of younger patients with chronic lymphocytic leukemia. <i>Blood</i> , 2006 , 107, 885-91	2.2	459
68	Strikingly homologous immunoglobulin gene rearrangements and poor outcome in VH3-21-using chronic lymphocytic leukemia patients independent of geographic origin and mutational status. <i>Blood</i> , 2006 , 107, 2889-94	2.2	149
67	Distinct gene expression patterns in chronic lymphocytic leukemia defined by usage of specific VH genes. <i>Blood</i> , 2006 , 107, 2090-3	2.2	46
66	Genetics, gene expression, and targeted therapies in chronic lymphocytic leukemia. <i>Current Drug Targets</i> , 2006 , 7, 1313-27	3	7
65	Clonal Evolution in Chronic Lymphocytic Leukemia: Acquisition of High-Risk Genomic Aberrations Associated with Unmutated VH, Resistance to Therapy, and Short Survival.. <i>Blood</i> , 2006 , 108, 296-296	2.2	1
64	Occurrence of Chromosomal Translocations as Independent Prognostic Factor in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2006 , 108, 2084-2084	2.2	

63	RHAMM/CD168 Is a Novel Leukemia Associated Antigen with Prognostic Value for Patients with B-Cell Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2006 , 108, 2773-2773	2.2	
62	Influence of MDM2 Single Nucleotide Polymorphism SNP309 on Disease Onset and Course in CLL.. <i>Blood</i> , 2006 , 108, 4938-4938	2.2	
61	Array-CGH Based Genomic Profiling in B-Cell Chronic Lymphocytic Leukemia Reveals Specific Correlations of Genomic Imbalances and Prognostic Subgroups and Underlines the Consistency of Chromosomal Aberration Patterns within This Disease.. <i>Blood</i> , 2006 , 108, 2083-2083	2.2	
60	Bone marrow transplantation nephropathy after an intensified conditioning regimen with radioimmunotherapy and allogeneic stem cell transplantation. <i>Journal of Nuclear Medicine</i> , 2006 , 47, 278-86	8.9	16
59	188Re or 90Y-labelled anti-CD66 antibody as part of a dose-reduced conditioning regimen for patients with acute leukaemia or myelodysplastic syndrome over the age of 55: results of a phase III study. <i>British Journal of Haematology</i> , 2005 , 130, 604-13	4.5	84
58	Duplication of chromosome arms 9q and 11q: evidence for a novel, 14q32 translocation-independent pathogenetic pathway in multiple myeloma. <i>Genes Chromosomes and Cancer</i> , 2005 , 42, 78-81	5	4
57	Translocation t(X;11)(q13;q23) in B-cell chronic lymphocytic leukemia disrupts two novel genes. <i>Genes Chromosomes and Cancer</i> , 2005 , 42, 128-43	5	29
56	Evidence for distinct pathomechanisms in genetic subgroups of chronic lymphocytic leukemia revealed by quantitative expression analysis of cell cycle, activation, and apoptosis-associated genes. <i>Journal of Clinical Oncology</i> , 2005 , 23, 3780-92	2.2	60
55	In-Vitro Treatment of Chronic Lymphocytic Leukemia (CLL) Cells with Fludarabine, Etoposide and Alemtuzumab (Campath-1H) Reveals Different Rates and Mechanisms of Cell Death.. <i>Blood</i> , 2005 , 106, 2112-2112	2.2	1
54	Fludarabine Versus Fludarabine Plus Epirubicin in the Treatment of Chronic Lymphocytic Leukemia - Final Results of a German Randomized Phase-III Study.. <i>Blood</i> , 2005 , 106, 2123-2123	2.2	3
53	Graft Versus Host Disease-Like Syndrome after Autologous Stem Cell Transplantation in CLL Patients Treated with a TBI/Cy/Alemtuzumab (CAMPATH-1H) High-Dose Regimen.. <i>Blood</i> , 2005 , 106, 2928-2928	2.2	1
52	17p Deletion Predicts for Inferior Overall Survival after Fludarabine - Based First Line Therapy in Chronic Lymphocytic Leukemia: First Analysis of Genetics in the CLL4 Trial of the GCLLSG.. <i>Blood</i> , 2005 , 106, 715-715	2.2	36
51	The Paternally Expressed Gene 10 (PEG10) on Chromosome 7q21 Is Overexpressed and Imprinted in High-Risk B-CLL.. <i>Blood</i> , 2005 , 106, 1221-1221	2.2	
50	Short Telomeres Are Associated with Genetic Instability and the Occurrence of High Risk Genomic Aberrations in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2005 , 106, 1178-1178	2.2	
49	Strikingly Homologous Immunoglobulin Gene Rearrangements and Poor Outcome in VH3-21-Utilizing Chronic Lymphocytic Leukemia Independent of Geographical Origin and Mutational Status.. <i>Blood</i> , 2005 , 106, 175-175	2.2	
48	MRD Kinetics Can Predict the Time to Relapse after Autologous Stem Cell Transplantation (SCT) in Chronic Lymphocytic Leukemia (CLL).. <i>Blood</i> , 2005 , 106, 714-714	2.2	1
47	ZAP-70 methylation and expression status in chronic lymphocytic leukemia. <i>Haematologica</i> , 2005 , 90, 1012	6.6	
46	Microarray gene expression profiling of B-cell chronic lymphocytic leukemia subgroups defined by genomic aberrations and VH mutation status. <i>Journal of Clinical Oncology</i> , 2004 , 22, 3937-49	2.2	177

45	Automated array-based genomic profiling in chronic lymphocytic leukemia: development of a clinical tool and discovery of recurrent genomic alterations. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 1039-44	11.5	206
44	Molecular genetics and its clinical relevance. <i>Hematology/Oncology Clinics of North America</i> , 2004 , 18, 827-48, viii	3.1	16
43	The prognostic impact of autologous stem cell transplantation in patients with chronic lymphocytic leukemia: a risk-matched analysis based on the VH gene mutational status. <i>Blood</i> , 2004 , 103, 2850-8	2.2	93
42	Graft-versus-leukemia activity may overcome therapeutic resistance of chronic lymphocytic leukemia with unmutated immunoglobulin variable heavy-chain gene status: implications of minimal residual disease measurement with quantitative PCR. <i>Blood</i> , 2004 , 104, 2600-2	2.2	139
41	Early Autologous Stem Cell Transplantation (SCT) in Genetically Poor-Risk Chronic Lymphocytic Leukemia Is Feasible and Effective: Results from a Prospective Multicenter Study (GCLLSG CLL3 Protocol).. <i>Blood</i> , 2004 , 104, 146-146	2.2	7
40	Minimal Residual Disease (MRD) Kinetics after Autologous Stem Cell Transplantation for Chronic Lymphocytic Leukemia (CLL) Correlate to IgH VH Mutational Status and Predict Post Transplant Outcome: Real Time ASO-IgH PCR Results of the GCLLSG CLL3 Protocol.. <i>Blood</i> , 2004 , 104, 147-147	2.2	5
39	ZAP-70 Expression, VH-Mutation Status, Genomic Aberrations and Prognosis in Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2004 , 104, 1920-1920	2.2	1
38	Consolidation with Alemtuzumab in First Remission Induces Pronounced MRD Reduction and Clinical Remissions - Update on a Randomized Phase III Trial of the German CLL Study Group (GCLLSG).. <i>Blood</i> , 2004 , 104, 2506-2506	2.2	3
37	Subcutaneous Campath-1H (Alemtuzumab) in Fludarabine-Refractory CLL: Interim Analysis of the CLL2h Study of the German CLL Study Group (GCLLSG).. <i>Blood</i> , 2004 , 104, 478-478	2.2	24
36	Protein Expression Patterns of Candidate Genes Involved in Apoptosis and Cell Cycle Control in Genetic Subgroups of Chronic Lymphocytic Leukemia.. <i>Blood</i> , 2004 , 104, 2796-2796	2.2	
35	Prognostic Relevance of Lipoprotein Lipase (LPL) Expression in B-CLL.. <i>Blood</i> , 2004 , 104, 177-177	2.2	
34	Differential Expression Patterns of Apoptosis and Cell Cycle Proteins after Staurosporine Treatment in EHEB (B-CLL) and JURKAT (T-ALL) Cell Lines.. <i>Blood</i> , 2004 , 104, 4304-4304	2.2	
33	Correlation of Quantitative Gene Expression by RQ-PCR with Clinical and Biologic Factors in Mantle Cell Lymphoma.. <i>Blood</i> , 2004 , 104, 696-696	2.2	1
32	High ZAP-70 and Differential Expression of B-Cell Receptor Related Genes in Chronic Lymphocytic Leukemia with V3-21 Gene Usage.. <i>Blood</i> , 2004 , 104, 773-773	2.2	
31	Protein Expression Analysis of Chromosome 12 Candidate Genes in Chronic Lymphocytic Leukemia (B-CLL).. <i>Blood</i> , 2004 , 104, 4797-4797	2.2	
30	Genetic analysis of de novo CD5+ diffuse large B-cell lymphomas suggests an origin from a somatically mutated CD5+ progenitor B cell. <i>Blood</i> , 2003 , 101, 699-702	2.2	39
29	Unmutated immunoglobulin variable heavy-chain gene status remains an adverse prognostic factor after autologous stem cell transplantation for chronic lymphocytic leukemia. <i>Blood</i> , 2003 , 101, 2049-53	2.2	102
28	VH mutation status and VDJ rearrangement structure in mantle cell lymphoma: correlation with genomic aberrations, clinical characteristics, and outcome. <i>Blood</i> , 2003 , 102, 3003-9	2.2	122

27	Value of comparative genomic hybridization and fluorescence in situ hybridization for molecular diagnostics in multiple myeloma. <i>British Journal of Haematology</i> , 2003 , 122, 193-201	4.5	30
26	Campath-1H-induced complete remission of chronic lymphocytic leukemia despite p53 gene mutation and resistance to chemotherapy. <i>New England Journal of Medicine</i> , 2002 , 347, 452-3	59.2	177
25	Down-regulation of candidate tumor suppressor genes within chromosome band 13q14.3 is independent of the DNA methylation pattern in B-cell chronic lymphocytic leukemia. <i>Blood</i> , 2002 , 99, 4116-21	2.2	86
24	Pathogenic complexity of gastric B-cell lymphoma. <i>Blood</i> , 2002 , 100, 1095-6; author reply 1096-7	2.2	5
23	Genetic imbalances in progressed B-cell chronic lymphocytic leukemia and transformed large-cell lymphoma (Richter's syndrome). <i>American Journal of Pathology</i> , 2002 , 161, 957-68	5.8	75
22	Evidence for distinct pathomechanisms in B-cell chronic lymphocytic leukemia and mantle cell lymphoma by quantitative expression analysis of cell cycle and apoptosis-associated genes. <i>Blood</i> , 2002 , 99, 4554-61	2.2	117
21	V H mutation status, CD38 expression level, genomic aberrations, and survival in chronic lymphocytic leukemia. <i>Blood</i> , 2002 , 100, 1410-1416	2.2	633
20	V(H) mutation status, CD38 expression level, genomic aberrations, and survival in chronic lymphocytic leukemia. <i>Blood</i> , 2002 , 100, 1410-6	2.2	155
19	CDNA microarray gene expression analysis of B-cell chronic lymphocytic leukemia proposes potential new prognostic markers involved in lymphocyte trafficking. <i>International Journal of Cancer</i> , 2001 , 91, 474-80	7.5	93
18	Molecular-cytogenetic comparison of mucosa-associated marginal zone B-cell lymphoma and large B-cell lymphoma arising in the gastro-intestinal tract. <i>Genes Chromosomes and Cancer</i> , 2001 , 31, 316-25	5	73
17	B-cell neoplasia associated gene with multiple splicing (BCMS): the candidate B-CLL gene on 13q14 comprises more than 560 kb covering all critical regions. <i>Human Molecular Genetics</i> , 2001 , 10, 1275-85	5.6	70
16	t(11;14)-positive mantle cell lymphomas exhibit complex karyotypes and share similarities with B-cell chronic lymphocytic leukemia 2000 , 27, 285-294		112
15	BCMSUN, a candidate gene for B-cell chronic lymphocytic leukemia and mantle-cell lymphoma, has an independently expressed homolog on 1p22-p31, BCMSUN-like. <i>International Journal of Cancer</i> , 2000 , 88, 692-7	7.5	15
14	Genomic aberrations and survival in chronic lymphocytic leukemia. <i>New England Journal of Medicine</i> , 2000 , 343, 1910-6	59.2	2573
13	Molecular Characterization of 11q Deletions Points to a Pathogenic Role of the ATM Gene in Mantle Cell Lymphoma. <i>Blood</i> , 1999 , 94, 3262-3264	2.2	106
12	Somatic ATM Mutations Indicate a Pathogenic Role of ATM in B-Cell Chronic Lymphocytic Leukemia. <i>Blood</i> , 1999 , 94, 748-753	2.2	224
11	Chromosome aberrations in B-cell chronic lymphocytic leukemia: reassessment based on molecular cytogenetic analysis. <i>Journal of Molecular Medicine</i> , 1999 , 77, 266-81	5.5	174
10	Exclusive detection of the t(11;18)(q21;q21) in extranodal marginal zone B cell lymphomas (MZBL) of MALT type in contrast to other MZBL and extranodal large B cell lymphomas. <i>American Journal of Pathology</i> , 1999 , 155, 1817-21	5.8	112

9	Expression of costimulatory molecules in low-grade mucosa-associated lymphoid tissue-type lymphomas in vivo. <i>American Journal of Pathology</i> , 1999 , 155, 2019-27	5.8	40
8	Expressed sequences as candidates for a novel tumor suppressor gene at band 13q14 in B-cell chronic lymphocytic leukemia and mantle cell lymphoma. <i>Oncogene</i> , 1998 , 16, 1891-7	9.2	130
7	11q Deletions Identify a New Subset of B-Cell Chronic Lymphocytic Leukemia Characterized by Extensive Nodal Involvement and Inferior Prognosis. <i>Blood</i> , 1997 , 89, 2516-2522	2.2	328
6	Biallelic mutations in the ATM gene in T-prolymphocytic leukemia. <i>Nature Medicine</i> , 1997 , 3, 1155-9	50.5	217
5	CDKN2 gene deletion is not found in chronic lymphoid leukaemias of B- and T-cell origin but is frequent in acute lymphoblastic leukaemia. <i>British Journal of Haematology</i> , 1995 , 91, 865-70	4.5	19
4	Molecular cytogenetic analysis of RB-1 deletions in chronic B-cell leukemias. <i>Leukemia and Lymphoma</i> , 1994 , 16, 97-103	1.9	23
3	Reciprocal translocation t(12;13)(p13;q14) in acute nonlymphoblastic leukemia: report and cytogenetic analysis of two cases. <i>Cancer Genetics and Cytogenetics</i> , 1994 , 77, 106-10		13
2	Chronic Lymphocytic Leukemia: Pathophysiology, Diagnosis and Manifestations, and Prognostic Markers 297-307		
1	NOTCH1 drives immune-escape mechanisms in B cell malignancies		1