
## Hana Å ubrtovÃ; SalmonovÃ;

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5669228/publications.pdf

Version: 2024-02-01



| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Research Note: The effects of genotype, sex, and feeding regime on performance, carcasses characteristic, and microbiota in chickens. Poultry Science, 2021, 100, 760-764.   | 3.4 | 18        |
| 2  | In Vitro Selective Antibacterial and Antiproliferative Effects of Ethanolic Extracts from Cambodian<br>and Philippine Plants Used in Folk Medicine for Diarrhea Treatment. Frontiers in Pharmacology, 2021,<br>12, 746808.   | 3.5 | 3         |
| 3  | In Vitro Selective Growth-Inhibitory Activities of Phytochemicals, Synthetic Phytochemical Analogs,<br>and Antibiotics against Diarrheagenic/Probiotic Bacteria and Cancer/Normal Intestinal Cells.<br>Pharmaceuticals, 2020, 13, 233.                                     | 3.8 | 16        |
| 4  | Pathogenic profile and cytotoxic activity of Aeromonas spp. isolated from Pectinatella magnifica and surrounding water in the South Bohemian aquaculture region. Journal of Fish Diseases, 2020, 43, 1213-1227.  | 1.9 | 0         |
| 5  | Colonization of Germ-Free Piglets with Mucinolytic and Non-Mucinolytic Bifidobacterium boum<br>Strains Isolated from the Intestine of Wild Boar and Their Interference with Salmonella Typhimurium.<br>Microorganisms, 2020, 8, 2002.                                      | 3.6 | 7         |
| 6  | Effect of probiotic Clostridium butyricum CBM 588 on microbiota and growth performance of broiler chickens. Czech Journal of Animal Science, 2019, 64, 387-394.  | 1.3 | 5         |
| 7  | High Mobility Group Box 1 and TLR4 Signaling Pathway in Gnotobiotic Piglets Colonized/Infected with<br>L. amylovorus, L. mucosae, E. coli Nissle 1917 and S. Typhimurium. International Journal of Molecular<br>Sciences, 2019, 20, 6294.                                  | 4.1 | 13        |
| 8  | Melanoma-related changes in skin microbiome. Folia Microbiologica, 2019, 64, 435-442.  | 2.3 | 54        |
| 9  | Cultivable bacteria from Pectinatella magnifica and the surrounding water in South Bohemia indicate<br>potential new Gammaproteobacterial, Betaproteobacterial and Firmicutes taxa. FEMS Microbiology<br>Letters, 2018, 365, .   | 1.8 | 4         |
| 10 | POLLEN CAN - TESTING OF BEE POLLEN FERMENTATION IN MODEL CONDITIONS. Journal of Microbiology, Biotechnology and Food Sciences, 2018, 8, 805-811.   | 0.8 | 7         |
| 11 | Analysis of Cutaneous Microbiota of Piglets with Hereditary Melanoma. Scientia Agriculturae<br>Bohemica, 2018, 49, 285-290.  | 0.3 | 1         |
| 12 | Selection of prebiotic oligosaccharides suitable for synbiotic use in calves. Animal Feed Science and Technology, 2017, 229, 73-78.  | 2.2 | 5         |
| 13 | Methods of Studying Diversity of Bacterial Comunities: A Review. Scientia Agriculturae Bohemica, 2017,<br>48, 154-165.   | 0.3 | 9         |
| 14 | Bifidobacterium apri sp. nov., a thermophilic actinobacterium isolated from the digestive tract of wild<br>pigs (Sus scrofa). International Journal of Systematic and Evolutionary Microbiology, 2017, 67,<br>2349-2356.   | 1.7 | 21        |
| 15 | Galliscardovia ingluviei gen. nov., sp. nov., a thermophilic bacterium of the family Bifidobacteriaceae<br>isolated from the crop of a laying hen (Gallus gallus f. domestica). International Journal of<br>Systematic and Evolutionary Microbiology, 2017, 67, 2403-2411. | 1.7 | 14        |
| 16 | Alloscardovia venturai sp. nov., a fructose 6-phosphate phosphoketolase-positive species isolated<br>from the oral cavity of a guinea-pig (Cavia aperea f. porcellus). International Journal of Systematic and<br>Evolutionary Microbiology, 2017, 67, 2842-2847.          | 1.7 | 9         |
| 17 | Lactobacillus caviae sp. nov., an obligately heterofermentative bacterium isolated from the oral cavity<br>of a guinea pig (Cavia aperea f. porcellus). International Journal of Systematic and Evolutionary<br>Microbiology, 2017, 67, 2903-2909.                         | 1.7 | 10        |
| 18 | Assessment of Chemical Impact of Invasive Bryozoan Pectinatella magnifica on the Environment:<br>Cytotoxicity and Antimicrobial Activity of P. magnifica Extracts. Molecules, 2016, 21, 1476.  | 3.8 | 4         |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Anticlostridial agent 8-hydroxyquinoline improves the isolation of faecal bifidobacteria on modified<br>Wilkins-Chalgren agar with mupirocin. Letters in Applied Microbiology, 2016, 62, 330-335. | 2.2 | 7         |
| 20 | Identification of microbiota associated with Pectinatella magnifica in South Bohemia. Biologia<br>(Poland), 2015, 70, 365-371.  | 1.5 | 2         |
| 21 | A new medium containing mupirocin, acetic acid, and norfloxacin for the selective cultivation of bifidobacteria. Anaerobe, 2015, 34, 27-33.   | 2.1 | 23        |