

Zhong Xin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5669368/publications.pdf>

Version: 2024-02-01

34
papers

626
citations

566801

15
h-index

610482

24
g-index

34
all docs

34
docs citations

34
times ranked

826
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | The value of SPECT/CT imaging of lacrimal glands as a means of assessing the activity of Gravesâ€™ orbitopathy. <i>Endocrine Connections</i> , 2022, 11, . | 0.8 | 4 |
| 2 | Increasing obstructive sleep apnea risk is associated with hearing impairment in middle-aged Chinese menâ€™A cross-sectional study. <i>PLoS ONE</i> , 2022, 17, e0268412. | 1.1 | 3 |
| 3 | A High TSH Level is Associated with Diabetic Macular Edema: A Cross-Sectional Study of Patients with Type 2 Diabetes Mellitus. <i>Endocrine Connections</i> , 2022, , . | 0.8 | 2 |
| 4 | Nifuroxazide improves insulin secretion and attenuates high glucose-induced inflammation and apoptosis in INS-1Îšcells. <i>European Journal of Pharmacology</i> , 2021, 899, 174042. | 1.7 | 13 |
| 5 | Early Onset Age Increased the Risk of Diabetic Retinopathy in Type 2 Diabetes Patients with Duration of 10â€™20 Years and HbA1C â‰¥7%: A Hospital-Based Case-Control Study. <i>International Journal of Endocrinology</i> , 2021, 2021, 1-6. | 0.6 | 7 |
| 6 | Effect of hemoglobin content on cerebral oxygen saturation during surgery for scoliosis in pediatric patients. <i>BMC Anesthesiology</i> , 2021, 21, 165. | 0.7 | 6 |
| 7 | Incidence of Type 1 Diabetes May Be Underestimated in the Chinese Population: Evidence From 21.7 Million People Between 2007 and 2017. <i>Diabetes Care</i> , 2021, 44, 2503-2509. | 4.3 | 23 |
| 8 | Relationship between Central Obesity and the incidence of Cognitive Impairment and Dementia from Cohort Studies Involving 5,060,687 Participants. <i>Neuroscience and Biobehavioral Reviews</i> , 2021, 130, 301-313. | 2.9 | 43 |
| 9 | Comparative assessment of gut microbial composition and function in patients with Gravesâ€™ disease and Gravesâ€™ orbitopathy. <i>Journal of Endocrinological Investigation</i> , 2021, 44, 297-310. | 1.8 | 24 |
| 10 | Clinical and genetic analysis of idiopathic normophosphatemic tumoral calcinosis in 19 patients. <i>Journal of Endocrinological Investigation</i> , 2020, 43, 173-183. | 1.8 | 4 |
| 11 | Angiotensinâ€™(1â€™7), the product of ACE2 ameliorates NAFLD by acting through its receptor Mas to regulate hepatic mitochondrial function and glycolipid metabolism. <i>FASEB Journal</i> , 2020, 34, 16291-16306. | 0.2 | 19 |
| 12 | Association of tidal volume during mechanical ventilation with postoperative pulmonary complications in pediatric patients undergoing major scoliosis surgery. <i>Paediatric Anaesthesia</i> , 2020, 30, 806-813. | 0.6 | 0 |
| 13 | Identifying the insulin resistance index in nondiabetic Chinese subjects. <i>Medicine (United States)</i> , 2020, 99, e19023. | 0.4 | 12 |
| 14 | Preliminary evidence of the association between DNAm and orbital volumetry in GO. <i>Endocrine Connections</i> , 2020, 9, 617-626. | 0.8 | 2 |
| 15 | Alterations in the intestinal microbiota of patients with severe and active Gravesâ€™ orbitopathy: a cross-sectional study. <i>Journal of Endocrinological Investigation</i> , 2019, 42, 967-978. | 1.8 | 41 |
| 16 | Vitamin D deficiency is associated with risk of developing peripheral arterial disease in type 2 diabetic patients. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 145. | 0.7 | 14 |
| 17 | Identifying and Validating Genes with DNA Methylation Data in the Context of Biological Network for Chinese Patients with Gravesâ€™ Orbitopathy. <i>International Journal of Endocrinology</i> , 2019, 2019, 1-9. | 0.6 | 8 |
| 18 | A Pathway Analysis Based on Genome-Wide DNA Methylation of Chinese Patients with Gravesâ€™ Orbitopathy. <i>BioMed Research International</i> , 2019, 2019, 1-10. | 0.9 | 10 |

| # | ARTICLE | IF | CITATIONS |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | The Potential Link between Gut Microbiota and Serum TRAb in Chinese Patients with Severe and Active Graves' Orbitopathy. <i>International Journal of Endocrinology</i> , 2019, 2019, 1-12. | 0.6 | 16 |
| 20 | Angiotensin-converting enzyme 2 regulates endoplasmic reticulum stress and mitochondrial function to preserve skeletal muscle lipid metabolism. <i>Lipids in Health and Disease</i> , 2019, 18, 207. | 1.2 | 33 |
| 21 | Angiotensin-converting enzyme 2 inhibits endoplasmic reticulum stress-associated pathway to preserve nonalcoholic fatty liver disease. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3123. | 1.7 | 24 |
| 22 | A genome-wide DNA methylation analysis in peripheral blood from patients identifies risk loci associated with Graves' orbitopathy. <i>Journal of Endocrinological Investigation</i> , 2018, 41, 719-727. | 1.8 | 14 |
| 23 | Are low levels of serum bicarbonate associated with risk of progressing to impaired fasting glucose/diabetes? A single-centre prospective cohort study in Beijing, China. <i>BMJ Open</i> , 2018, 8, e019145. | 0.8 | 3 |
| 24 | Prevalence and associated factors of diabetic retinopathy in Beijing, China: a cross-sectional study. <i>BMJ Open</i> , 2017, 7, e015473. | 0.8 | 45 |
| 25 | Reanalysis and External Validation of a Decision Tree Model for Detecting Unrecognized Diabetes in Rural Chinese Individuals. <i>International Journal of Endocrinology</i> , 2017, 2017, 1-6. | 0.6 | 5 |
| 26 | The Status of Maculopathy in Diabetes and Prediabetes Patients in a Population-Based Study Detected by Optical Coherence Tomography: The 2011 Health Examination Survey in Beijing. <i>BioMed Research International</i> , 2017, 2017, 1-7. | 0.9 | 1 |
| 27 | Angiotensin-converting enzyme 2/angiotensin-(1-7)/Mas axis activates Akt signaling to ameliorate hepatic steatosis. <i>Scientific Reports</i> , 2016, 6, 21592. | 1.6 | 63 |
| 28 | Using Hashimoto thyroiditis as gold standard to determine the upper limit value of thyroid stimulating hormone in a Chinese cohort. <i>BMC Endocrine Disorders</i> , 2016, 16, 57. | 0.9 | 3 |
| 29 | High Prevalence of Diabetic Retinopathy in Diabetic Patients Concomitant with Metabolic Syndrome. <i>PLoS ONE</i> , 2016, 11, e0145293. | 1.1 | 16 |
| 30 | High Prevalence of Lower Extremity Peripheral Artery Disease in Type 2 Diabetes Patients with Proliferative Diabetic Retinopathy. <i>PLoS ONE</i> , 2015, 10, e0122022. | 1.1 | 26 |
| 31 | The ACE2/Ang-(1-7)/Mas axis can inhibit hepatic insulin resistance. <i>Molecular and Cellular Endocrinology</i> , 2014, 393, 30-38. | 1.6 | 70 |
| 32 | Identifying obesity indicators which best correlate with type 2 diabetes in a Chinese population. <i>BMC Public Health</i> , 2012, 12, 732. | 1.2 | 24 |
| 33 | Evaluation for Fasting and 2-hour Glucose and HbA1c for Diagnosing Diabetes Based on Prevalence of Retinopathy in a Chinese Population. <i>PLoS ONE</i> , 2012, 7, e40610. | 1.1 | 26 |
| 34 | A simple tool detected diabetes and prediabetes in rural Chinese. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 1030-1035. | 2.4 | 22 |