

Felicita Pedata

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5669940/felicita-pedata-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

121
papers

5,414
citations

43
h-index

69
g-index

131
ext. papers

5,895
ext. citations

5.3
avg, IF

5.21
L-index

#	Paper	IF	Citations
121	Role of Carbonic Anhydrase in Cerebral Ischemia and Carbonic Anhydrase Inhibitors as Putative Protective Agents. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
120	A Adenosine Receptors and Sphingosine 1-Phosphate Signaling Cross-Talk in Oligodendroglioneogenesis. <i>Frontiers in Neuroscience</i> , 2021 , 15, 677988	5.1	
119	Uncovering the Mechanisms of Adenosine Receptor-Mediated Pain Control: Focus on the A Receptor Subtype. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	4
118	Acetylcholine modulates K and Na currents in human basal forebrain cholinergic neuroblasts through an autocrine/paracrine mechanism. <i>Journal of Neurochemistry</i> , 2021 , 157, 1182-1195	6	2
117	Protective effects of carbonic anhydrase inhibition in brain ischaemia and models. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2021 , 36, 964-976	5.6	4
116	A Adenosine Receptors: When Outsiders May Become an Attractive Target to Treat Brain Ischemia or Demyelination. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	10
115	Adenosine: A Relevant Role in Hypoxia. <i>Journal of Caffeine and Adenosine Research</i> , 2020 , 10, 2-3	1.6	
114	Adenosine A receptors inhibit K currents and cell differentiation in cultured oligodendrocyte precursor cells and modulate sphingosine-1-phosphate signaling pathway. <i>Biochemical Pharmacology</i> , 2020 , 177, 113956	6	12
113	Multicentre translational Trial of Remote Ischaemic Conditioning in Acute Ischaemic Stroke (TRICS): protocol of multicentre, parallel group, randomised, preclinical trial in female and male rat and mouse from the Italian Stroke Organization (ISO) Basic Science network.. <i>BMJ Open Science</i> , 2020 , 1, e000043	4.6	5
112	Acute visceral pain relief mediated by A3AR agonists in rats: involvement of N-type voltage-gated calcium channels. <i>Pain</i> , 2020 , 161, 2179-2190	8	11
111	New Insight into the Role of Adenosine in Demyelination, Stroke and Neuropathic Pain. <i>Frontiers in Pharmacology</i> , 2020 , 11, 625662	5.6	3
110	Protective Effect of Adenosine A Receptor Agonist, BAY60-6583, Against Transient Focal Brain Ischemia in Rat. <i>Frontiers in Pharmacology</i> , 2020 , 11, 588757	5.6	5
109	Functional characterization of a novel adenosine A receptor agonist on short-term plasticity and synaptic inhibition during oxygen and glucose deprivation in the rat CA1 hippocampus. <i>Brain Research Bulletin</i> , 2019 , 151, 174-180	3.9	11
108	Adenosine A3 receptor activation inhibits pronociceptive N-type Ca ²⁺ currents and cell excitability in dorsal root ganglion neurons. <i>Pain</i> , 2019 , 160, 1103-1118	8	24
107	The Selective Antagonism of Adenosine A Receptors Reduces the Synaptic Failure and Neuronal Death Induced by Oxygen and Glucose Deprivation in Rat CA1 Hippocampus. <i>Frontiers in Pharmacology</i> , 2018 , 9, 399	5.6	23
106	Adenosine and Oxygen/Glucose Deprivation in the Brain 2018 , 351-378		
105	A Selective Histamine H Receptor Antagonist, JNJ7777120, Is Protective in a Rat Model of Transient Cerebral Ischemia. <i>Frontiers in Pharmacology</i> , 2018 , 9, 1231	5.6	6

104	The neuron-astrocyte-microglia triad in CA3 after chronic cerebral hypoperfusion in the rat: Protective effect of dipyridamole. <i>Experimental Gerontology</i> , 2017 , 96, 46-62	4.5	22
103	Imidazo[1,2-a]pyrazin-8-amine core for the design of new adenosine receptor antagonists: Structural exploration to target the A and A subtypes. <i>European Journal of Medicinal Chemistry</i> , 2017 , 125, 611-628	6.8	15
102	Adenosine and Oxygen/Glucose Deprivation in the Brain 2017 , 151-173		
101	Equilibrative nucleoside transporter ENT1 as a biomarker of Huntington disease. <i>Neurobiology of Disease</i> , 2016 , 96, 47-53	7.5	15
100	Purinergic signalling in brain ischemia. <i>Neuropharmacology</i> , 2016 , 104, 105-30	5.5	101
99	Time-course of protection by the selective A2A receptor antagonist SCH58261 after transient focal cerebral ischemia. <i>Neurological Sciences</i> , 2015 , 36, 1441-8	3.5	19
98	Role of adenosine in oligodendrocyte precursor maturation. <i>Frontiers in Cellular Neuroscience</i> , 2015 , 9, 155	6.1	28
97	Adenosine receptors in cerebral ischemia. <i>International Review of Neurobiology</i> , 2014 , 119, 309-48	4.4	39
96	Low doses of the selective adenosine A2A receptor agonist CGS21680 are protective in a rat model of transient cerebral ischemia. <i>Brain Research</i> , 2014 , 1551, 59-72	3.7	42
95	The selective antagonism of P2X7 and P2Y1 receptors prevents synaptic failure and affects cell proliferation induced by oxygen and glucose deprivation in rat dentate gyrus. <i>PLoS ONE</i> , 2014 , 9, e115273	3.7	12
94	The neuron-astrocyte-microglia triad in a rat model of chronic cerebral hypoperfusion: protective effect of dipyridamole. <i>Frontiers in Aging Neuroscience</i> , 2014 , 6, 322	5.3	34
93	Adenosine A2A receptors modulate acute injury and neuroinflammation in brain ischemia. <i>Mediators of Inflammation</i> , 2014 , 2014, 805198	4.3	39
92	Amyloid- β oligomer synaptotoxicity is mimicked by oligomers of the model protein HypF-N. <i>Neurobiology of Aging</i> , 2013 , 34, 2100-9	5.6	26
91	Effects of oxygen and glucose deprivation on synaptic transmission in rat dentate gyrus: role of A2A adenosine receptors. <i>Neuropharmacology</i> , 2013 , 67, 511-20	5.5	24
90	Adenosine A _{2A} receptors inhibit delayed rectifier potassium currents and cell differentiation in primary purified oligodendrocyte cultures. <i>Neuropharmacology</i> , 2013 , 73, 301-10	5.5	41
89	UDP-glucose enhances outward K(+) currents necessary for cell differentiation and stimulates cell migration by activating the GPR17 receptor in oligodendrocyte precursors. <i>Glia</i> , 2013 , 61, 1155-71	9	45
88	Adenosine is present in rat brain synaptic vesicles. <i>NeuroReport</i> , 2013 , 24, 982-7	1.7	17
87	Adenosine and Stroke 2013 , 273-306		1

86	3-Hydroxy-1H-quinazoline-2,4-dione derivatives as new antagonists at ionotropic glutamate receptors: molecular modeling and pharmacological studies. <i>European Journal of Medicinal Chemistry</i> , 2012 , 54, 470-82	6.8	28
85	Ecto-ATPase inhibition: ATP and adenosine release under physiological and ischemic in vivo conditions in the rat striatum. <i>Experimental Neurology</i> , 2012 , 233, 193-204	5.7	73
84	P2Y1 receptor modulation of Ca ²⁺ -activated K ⁺ currents in medium-sized neurons from neonatal rat striatal slices. <i>Journal of Neurophysiology</i> , 2012 , 107, 1009-21	3.2	18
83	P2 receptor antagonists prevent synaptic failure and extracellular signal-regulated kinase 1/2 activation induced by oxygen and glucose deprivation in rat CA1 hippocampus in vitro. <i>European Journal of Neuroscience</i> , 2011 , 33, 2203-15	3.5	18
82	CGS 21680, an agonist of the adenosine (A2A) receptor, decreases acute lung inflammation. <i>European Journal of Pharmacology</i> , 2011 , 668, 305-16	5.3	39
81	Selective adenosine A2A receptor agonists and antagonists protect against spinal cord injury through peripheral and central effects. <i>Journal of Neuroinflammation</i> , 2011 , 8, 31	10.1	47
80	Cognitive impairment with vascular impairment and degeneration. <i>Current Neurovascular Research</i> , 2011 , 8, 342-50	1.8	31
79	CGS 21680, an agonist of the adenosine (A2A) receptor, reduces progression of murine type II collagen-induced arthritis. <i>Journal of Rheumatology</i> , 2011 , 38, 2119-29	4.1	53
78	Effect of intravenous administration of dipyridamole in a rat model of chronic cerebral ischemia. <i>Annals of the New York Academy of Sciences</i> , 2010 , 1207, 89-96	6.5	23
77	Adenosine A3 Receptor Signaling in the Central Nervous System 2010 , 165-188		4
76	Adenosine A2A receptor-selective stimulation reduces signaling pathways involved in the development of intestine ischemia and reperfusion injury. <i>Shock</i> , 2010 , 33, 541-51	3.4	19
75	Functional characterization of two isoforms of the P2Y-like receptor GPR17: [³⁵ S]GTPγS binding and electrophysiological studies in 1321N1 cells. <i>American Journal of Physiology - Cell Physiology</i> , 2009 , 297, C1028-40	5.4	44
74	The adenosine A2A receptor antagonist ZM241385 enhances neuronal survival after oxygen-glucose deprivation in rat CA1 hippocampal slices. <i>British Journal of Pharmacology</i> , 2009 , 157, 818-30	8.6	43
73	Pyrido[2,3-e]-1,2,4-triazolo[4,3-a]pyrazin-1-one as a new scaffold to develop potent and selective human A3 adenosine receptor antagonists. Synthesis, pharmacological evaluation, and ligand-receptor modeling studies. <i>Journal of Medicinal Chemistry</i> , 2009 , 52, 2407-19	8.3	33
72	Selective adenosine A2a receptor antagonism reduces JNK activation in oligodendrocytes after cerebral ischaemia. <i>Brain</i> , 2009 , 132, 1480-95	11.2	73
71	The selective adenosine A2A receptor agonist CGS 21680 reduces JNK MAPK activation in oligodendrocytes in injured spinal cord. <i>Shock</i> , 2009 , 32, 578-85	3.4	40
70	Inducible nitric oxide synthase appears and is co-expressed with the neuronal isoform in interneurons of the rat hippocampus after transient ischemia induced by middle cerebral artery occlusion. <i>Experimental Neurology</i> , 2008 , 211, 433-40	5.7	16
69	Modulation of ischemic brain injury and neuroinflammation by adenosine A2A receptors. <i>Current Pharmaceutical Design</i> , 2008 , 14, 1490-9	3.3	66

68	Regulation of A(2A) adenosine receptor expression and functioning following permanent focal ischemia in rat brain. <i>Journal of Neurochemistry</i> , 2008 , 104, 479-90	6	35
67	Synthesis, ligand-receptor modeling studies and pharmacological evaluation of novel 4-modified-2-aryl-1,2,4-triazolo[4,3-a]quinoxalin-1-one derivatives as potent and selective human A3 adenosine receptor antagonists. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 6086-102	3.4	36
66	ATP modulates cell proliferation and elicits two different electrophysiological responses in human mesenchymal stem cells. <i>Stem Cells</i> , 2007 , 25, 1840-9	5.8	73
65	Differential glutamate-dependent and glutamate-independent adenosine A1 receptor-mediated modulation of dopamine release in different striatal compartments. <i>Journal of Neurochemistry</i> , 2007 , 101, 355-63	6	89
64	Role of adenosine A3 receptors on CA1 hippocampal neurotransmission during oxygen-glucose deprivation episodes of different duration. <i>Biochemical Pharmacology</i> , 2007 , 74, 768-79	6	52
63	Relationships between neurons expressing neuronal nitric oxide synthase, degree of microglia activation and animal survival. A study in the rat cortex after transient ischemia. <i>Brain Research</i> , 2007 , 1132, 218-27	3.7	17
62	Role of P2 purinergic receptors in synaptic transmission under normoxic and ischaemic conditions in the CA1 region of rat hippocampal slices. <i>Purinergic Signalling</i> , 2007 , 3, 203-19	3.8	30
61	The role of ATP and adenosine in the brain under normoxic and ischemic conditions. <i>Purinergic Signalling</i> , 2007 , 3, 299-310	3.8	43
60	Adenosine A2A receptors and brain injury: broad spectrum of neuroprotection, multifaceted actions and "fine tuning" modulation. <i>Progress in Neurobiology</i> , 2007 , 83, 310-31	10.9	205
59	New 2-arylpyrazolo[3,4-c]quinoline derivatives as potent and selective human A3 adenosine receptor antagonists. Synthesis, pharmacological evaluation, and ligand-receptor modeling studies. <i>Journal of Medicinal Chemistry</i> , 2007 , 50, 4061-74	8.3	55
58	The selective A2A receptor antagonist SCH 58261 protects from neurological deficit, brain damage and activation of p38 MAPK in rat focal cerebral ischemia. <i>Brain Research</i> , 2006 , 1073-1074, 470-80	3.7	65
57	A3 adenosine receptor antagonists delay irreversible synaptic failure caused by oxygen and glucose deprivation in the rat CA1 hippocampus in vitro. <i>British Journal of Pharmacology</i> , 2006 , 147, 524-32	8.6	60
56	P2X7 receptor modulation on microglial cells and reduction of brain infarct caused by middle cerebral artery occlusion in rat. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2006 , 26, 974-82	7.3	120
55	ATP extracellular concentrations are increased in the rat striatum during in vivo ischemia. <i>Neurochemistry International</i> , 2005 , 47, 442-8	4.4	201
54	Expression of neuronal and inducible nitric oxide synthase in neuronal and glial cells after transient occlusion of the middle cerebral artery. <i>Neuroscience</i> , 2005 , 136, 1015-26	3.9	14
53	The protective effect of adenosine A2A receptor antagonism in cerebral ischemia. <i>Neurological Research</i> , 2005 , 27, 169-74	2.7	49
52	The Controversial Role of Adenosine A2A Receptor Antagonists as Neuro-protective Agents. <i>Current Medicinal Chemistry - Central Nervous System Agents</i> , 2004 , 4, 35-45		5
51	Extracellular levels of amino acids and choline in human high grade gliomas: an intraoperative microdialysis study. <i>Neurochemical Research</i> , 2004 , 29, 325-34	4.6	46

50	Adenosine and glutamate extracellular concentrations and mitogen-activated protein kinases in the striatum of Huntington transgenic mice. Selective antagonism of adenosine A2A receptors reduces transmitter outflow. <i>Neurobiology of Disease</i> , 2004 , 17, 77-88	7.5	63
49	The selective A2A receptor antagonist SCH 58261 reduces striatal transmitter outflow, turning behavior and ischemic brain damage induced by permanent focal ischemia in the rat. <i>Brain Research</i> , 2003 , 959, 243-50	3.7	105
48	Adenosine A(2A) antagonism increases striatal glutamate outflow in the quinolinic acid rat model of Huntington's disease. <i>Brain Research</i> , 2003 , 979, 225-9	3.7	21
47	Adenosine A2A receptor antagonism increases striatal glutamate outflow in dopamine-denervated rats. <i>European Journal of Pharmacology</i> , 2003 , 464, 33-8	5.3	42
46	Brief, repeated, oxygen-glucose deprivation episodes protect neurotransmission from a longer ischemic episode in the in vitro hippocampus: role of adenosine receptors. <i>British Journal of Pharmacology</i> , 2003 , 140, 305-14	8.6	82
45	Adenosine extracellular levels in human brain gliomas: an intraoperative microdialysis study. <i>Neuroscience Letters</i> , 2003 , 346, 93-6	3.3	28
44	Modification of adenosine extracellular levels and adenosine A(2A) receptor mRNA by dopamine denervation. <i>European Journal of Pharmacology</i> , 2002 , 446, 75-82	5.3	63
43	Interactions among adenosine deaminase, adenosine A(1) receptors and dopamine D(1) receptors in stably cotransfected fibroblast cells and neurons. <i>Neuroscience</i> , 2002 , 113, 709-19	3.9	51
42	Adenosine in the central nervous system: release mechanisms and extracellular concentrations. <i>Journal of Neurochemistry</i> , 2001 , 79, 463-84	6	561
41	Adenosine extracellular brain concentrations and role of A2A receptors in ischemia. <i>Annals of the New York Academy of Sciences</i> , 2001 , 939, 74-84	6.5	96
40	Changes in hippocampal adenosine efflux, ATP levels, and synaptic transmission induced by increased temperature. <i>Synapse</i> , 2001 , 41, 58-64	2.4	18
39	Striatal A2A adenosine receptor antagonism differentially modifies striatal glutamate outflow in vivo in young and aged rats. <i>NeuroReport</i> , 2000 , 11, 2591-5	1.7	69
38	Extracellular adenosine concentrations during in vitro ischaemia in rat hippocampal slices. <i>British Journal of Pharmacology</i> , 1999 , 127, 729-39	8.6	105
37	Effect of A2A adenosine receptor stimulation and antagonism on synaptic depression induced by in vitro ischaemia in rat hippocampal slices. <i>British Journal of Pharmacology</i> , 1999 , 128, 1035-44	8.6	49
36	Adenosine and memory storage: effect of A(1) and A(2) receptor antagonists. <i>Psychopharmacology</i> , 1999 , 146, 214-9	4.7	80
35	Effect of N-methyl-D-aspartate on motor activity and in vivo adenosine striatal outflow in the rat. <i>European Journal of Pharmacology</i> , 1999 , 385, 15-9	5.3	30
34	Striatal outflow of adenosine, excitatory amino acids, gamma-aminobutyric acid, and taurine in awake freely moving rats after middle cerebral artery occlusion: correlations with neurological deficit and histopathological damage. <i>Stroke</i> , 1999 , 30, 2448-54; discussion 2455	6.7	128
33	Effect of adenosine A2A receptor stimulation on GABA release from the striatum of young and aged rats in vivo. <i>NeuroReport</i> , 1999 , 10, 3933-7	1.7	33

32	Striatal A2A adenosine receptors differentially regulate spontaneous and K ⁺ -evoked glutamate release in vivo in young and aged rats. <i>NeuroReport</i> , 1999 , 10, 687-91	1.7	67
31	Temporal correlation between adenosine outflow and synaptic potential inhibition in rat hippocampal slices during ischemia-like conditions. <i>Brain Research</i> , 1998 , 794, 325-8	3.7	42
30	In vivo amino acid release from the striatum of aging rats: adenosine modulation. <i>Neurobiology of Aging</i> , 1997 , 18, 243-50	5.6	35
29	The contribution of different types of calcium channels to electrically-evoked adenosine release from rat hippocampal slices. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1997 , 355, 250-5	3.4	7
28	A2 adenosine receptors: their presence and neuromodulatory role in the central nervous system. <i>General Pharmacology</i> , 1996 , 27, 925-33		74
27	Regulation of extracellular adenosine levels in the striatum of aging rats. <i>Brain Research</i> , 1995 , 684, 103-6	3.7	46
26	The source of brain adenosine outflow during ischemia and electrical stimulation. <i>Neurochemistry International</i> , 1995 , 27, 239-44	4.4	34
25	In vivo regulation of extracellular adenosine levels in the cerebral cortex by NMDA and muscarinic receptors. <i>European Journal of Pharmacology</i> , 1994 , 254, 277-82	5.3	34
24	Effect of idebenone on adenosine outflow and adenine nucleotide level in hippocampal slices under ischemia-like conditions. <i>European Journal of Pharmacology</i> , 1993 , 249, 65-70	5.3	3
23	Effect of K ⁺ depolarization, tetrodotoxin, and NMDA receptor inhibition on extracellular adenosine levels in rat striatum. <i>European Journal of Pharmacology</i> , 1993 , 234, 61-5	5.3	57
22	Adenosine outflow from hippocampal slices evoked by ischemic-like conditions: Effect of the excitatory amino acid antagonists. <i>Drug Development Research</i> , 1993 , 28, 395-398	5.1	4
21	Investigations into the adenosine outflow from hippocampal slices evoked by ischemia-like conditions. <i>Journal of Neurochemistry</i> , 1993 , 61, 284-9	6	99
20	Endogenous adenosine release from hippocampal slices: excitatory amino acid agonists stimulate release, antagonists reduce the electrically-evoked release. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1991 , 344, 538-43	3.4	29
19	Principal aspects of the regulation of acetylcholine release in the brain. <i>Progress in Brain Research</i> , 1990 , 84, 273-8	2.9	2
18	Regional differences in the electrically stimulated release of endogenous and radioactive adenosine and purine derivatives from rat brain slices. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1990 , 342, 447-53	3.4	29
17	Cholinergic and noradrenergic denervations decrease labelled purine release from electrically stimulated rat cortical slices. <i>Neuroscience</i> , 1989 , 32, 629-36	3.9	11
16	Muscarinic modulation of purine release from electrically stimulated rat cortical slices. <i>Journal of Neurochemistry</i> , 1988 , 50, 1074-9	6	12
15	Purinergetic modulation of cortical acetylcholine release is decreased in aging rats. <i>Experimental Gerontology</i> , 1988 , 23, 175-81	4.5	22

14	Effect of adenosine, adenosine derivatives, and caffeine on acetylcholine release from brain synaptosomes: interaction with muscarinic autoregulatory mechanisms. <i>Journal of Neurochemistry</i> , 1986 , 46, 1593-8	6	51
13	Chronic caffeine treatment reduces caffeine but not adenosine effects on cortical acetylcholine release. <i>British Journal of Pharmacology</i> , 1986 , 88, 671-6	8.6	16
12	Are the neurochemical and behavioral changes induced by lesions of the nucleus basalis in the rat a model of Alzheimer's disease?. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1986 , 10, 541-51	5.5	27
11	Phosphatidylserine increases acetylcholine release from cortical slices in aged rats. <i>Neurobiology of Aging</i> , 1985 , 6, 337-9	5.6	63
10	Biphasic effect of methylxanthines on acetylcholine release from electrically-stimulated brain slices. <i>British Journal of Pharmacology</i> , 1984 , 83, 69-73	8.6	25
9	A1 and A2 adenosine receptors modulate acetylcholine release from brain slices. <i>European Journal of Pharmacology</i> , 1984 , 97, 341-2	5.3	87
8	Acetylcholine release from rat cortical slices during postnatal development and aging. <i>Neurobiology of Aging</i> , 1983 , 4, 31-5	5.6	104
7	Effect of adenosine, adenosine triphosphate, adenosine deaminase, dipyridamole and aminophylline on acetylcholine release from electrically-stimulated brain slices. <i>Neuropharmacology</i> , 1983 , 22, 609-14	5.5	59
6	Changes in high affinity choline uptake in rat cortex following lesions of the magnocellular forebrain nuclei. <i>Brain Research</i> , 1982 , 233, 359-67	3.7	96
5	Changes in synaptosomal high affinity choline uptake following electrical stimulation of guinea-pig cortical slices: effect of atropine and physostigmine. <i>British Journal of Pharmacology</i> , 1981 , 74, 525-31	8.6	43
4	Choline high-affinity uptake and metabolism and choline acetyltransferase activity in the striatum of rats chronically treated with neuroleptics. <i>Journal of Neurochemistry</i> , 1980 , 35, 606-11	6	23
3	Acetylcholine input from the cerebral cortex, choline uptake and muscarinic receptors in morphine-dependent, freely-moving rats. <i>Neuropharmacology</i> , 1980 , 19, 597-605	5.5	23
2	Effect of choline, phosphorylcholine and dimethylaminoethanol on brain acetylcholine level in the rat. <i>Pharmacological Research Communications</i> , 1977 , 9, 755-61		37
1	Changes in regional brain acetylcholine levels during drug-induced convulsions. <i>European Journal of Pharmacology</i> , 1976 , 40, 329-35	5.3	8