

Suvi T Vaara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5674737/publications.pdf>

Version: 2024-02-01

47
papers

2,562
citations

361045

20
h-index

223531

46
g-index

48
all docs

48
docs citations

48
times ranked

3117
citing authors

#	ARTICLE	IF	CITATIONS
1	Causes of death for intensive care survivors with and without acute kidney injury in 5-year follow-up. <i>Acta Anaesthesiologica Scandinavica</i> , 2021, 65, 507-514.	0.7	0
2	Early prolonged neutrophil activation in critically ill patients with sepsis. <i>Innate Immunity</i> , 2021, 27, 192-200.	1.1	7
3	Fluid balance-adjusted creatinine in diagnosing acute kidney injury in the critically ill. <i>Acta Anaesthesiologica Scandinavica</i> , 2021, 65, 1079-1086.	0.7	4
4	Restrictive fluid management versus usual care in acute kidney injury (REVERSE-AKI): a pilot randomized controlled feasibility trial. <i>Intensive Care Medicine</i> , 2021, 47, 665-673.	3.9	33
5	Fluid management in patients with acute kidney injury – A post-hoc analysis of the FINNAKI study. <i>Journal of Critical Care</i> , 2021, 64, 205-210.	1.0	3
6	Clinical examination findings as predictors of acute kidney injury in critically ill patients. <i>Acta Anaesthesiologica Scandinavica</i> , 2020, 64, 69-74.	0.7	7
7	Burden of acute kidney injury and 90-day mortality in critically ill patients. <i>BMC Nephrology</i> , 2020, 21, 1.	0.8	86
8	Intravenous Contrast computed tomography versus native computed tomography in patients with acute Abdomen and impaired Renal function (INCARO): a multicentre, open-label, randomised controlled trial - study protocol. <i>BMJ Open</i> , 2020, 10, e037928.	0.8	4
9	Protocol and statistical analysis plan for the REstricted fluid therapy VERsus Standard treatment in Acute Kidney Injury – REVERSE-AKI randomized controlled pilot trial. <i>Acta Anaesthesiologica Scandinavica</i> , 2020, 64, 831-838.	0.7	6
10	Point-of-care creatinine measurements to predict acute kidney injury. <i>Acta Anaesthesiologica Scandinavica</i> , 2020, 64, 766-773.	0.7	5
11	Urinary cell cycle arrest biomarkers and chitinase 3-like protein 1 (CHI3L1) to detect acute kidney injury in the critically ill: a post hoc laboratory analysis on the FINNAKI cohort. <i>Critical Care</i> , 2020, 24, 144.	2.5	16
12	Noninterventional follow-up vs fluid bolus in RESPONSE to oliguria – The RESPONSE trial protocol and statistical analysis plan. <i>Acta Anaesthesiologica Scandinavica</i> , 2020, 64, 1210-1217.	0.7	3
13	Controversies in acute kidney injury: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Conference. <i>Kidney International</i> , 2020, 98, 294-309.	2.6	254
14	Different applications of the KDIGO criteria for AKI lead to different incidences in critically ill patients: a post hoc analysis from the prospective observational SICS-II study. <i>Critical Care</i> , 2020, 24, 164.	2.5	35
15	Costs and Cost-Utility of Critical Care and Subsequent Health Care: A Multicenter Prospective Study*. <i>Critical Care Medicine</i> , 2020, 48, e345-e355.	0.4	9
16	Two subphenotypes of septic acute kidney injury are associated with different 90-day mortality and renal recovery. <i>Critical Care</i> , 2020, 24, 150.	2.5	54
17	Urine NGAL as a biomarker for septic AKI: a critical appraisal of clinical utility – data from the observational FINNAKI study. <i>Annals of Intensive Care</i> , 2020, 10, 51.	2.2	27
18	Cytokine and lipid metabolome effects of low-dose acetylsalicylic acid in critically ill patients with systemic inflammation: a pilot, feasibility, multicentre, randomised, placebo-controlled trial. <i>Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine</i> , 2020, 22, 227-236.	0.0	2

#	ARTICLE	IF	CITATIONS
19	Neutrophil activation in septic acute kidney injury: A post hoc analysis of the FINNAKI study. <i>Acta Anaesthesiologica Scandinavica</i> , 2019, 63, 1390-1397.	0.7	11
20	Heme oxygenase-1 repeat polymorphism in septic acute kidney injury. <i>PLoS ONE</i> , 2019, 14, e0217291.	1.1	16
21	Common Inflammation-Related Candidate Gene Variants and Acute Kidney Injury in 2647 Critically Ill Finnish Patients. <i>Journal of Clinical Medicine</i> , 2019, 8, 342.	1.0	5
22	Association of endothelial and glycocalyx injury biomarkers with fluid administration, development of acute kidney injury, and 90-day mortality: data from the FINNAKI observational study. <i>Annals of Intensive Care</i> , 2019, 9, 103.	2.2	36
23	The predictive value of NT-proBNP and hs-TnT for risk of death in cardiac surgical patients. <i>Clinical Biochemistry</i> , 2018, 53, 65-71.	0.8	14
24	One- and three-year outcomes in patients treated with intermittent hemodialysis for acute kidney injury: prospective observational multicenter post hoc FINNAKI study. <i>Acta Anaesthesiologica Scandinavica</i> , 2018, 62, 1452-1459.	0.7	2
25	Fluid accumulation in acute kidney injury: More evidence toward harm and current challenges. <i>Acta Anaesthesiologica Scandinavica</i> , 2018, 62, 739-741.	0.7	0
26	How to improve the care of patients with acute kidney injury. <i>Intensive Care Medicine</i> , 2017, 43, 727-729.	3.9	7
27	Acute kidney injury in sepsis. <i>Intensive Care Medicine</i> , 2017, 43, 816-828.	3.9	490
28	Genetic variants in SERPINA4 and SERPINA5, but not BCL2 and SIK3 are associated with acute kidney injury in critically ill patients with septic shock. <i>Critical Care</i> , 2017, 21, 47.	2.5	21
29	Acute Kidney Injury After Cardiac Surgery by Complete KDIGO Criteria Predicts Increased Mortality. <i>Journal of Cardiothoracic and Vascular Anesthesia</i> , 2017, 31, 827-836.	0.6	44
30	Heparin-binding protein (HBP) improves prediction of sepsis-related acute kidney injury. <i>Annals of Intensive Care</i> , 2017, 7, 105.	2.2	34
31	Assessment of plasma endostatin to predict acute kidney injury in critically ill patients. <i>Acta Anaesthesiologica Scandinavica</i> , 2017, 61, 1286-1295.	0.7	14
32	Urinary Biomarkers Indicative of Apoptosis and Acute Kidney Injury in the Critically Ill. <i>PLoS ONE</i> , 2016, 11, e0149956.	1.1	20
33	Three-year mortality in 30-day survivors of critical care with acute kidney injury: data from the prospective observational FINNAKI study. <i>Annals of Intensive Care</i> , 2016, 6, 118.	2.2	10
34	Association of plasma chloride values with acute kidney injury in the critically ill – a prospective observational study. <i>Acta Anaesthesiologica Scandinavica</i> , 2016, 60, 790-799.	0.7	50
35	Plasma anti-FXa level as a surrogate marker of the adequacy of thromboprophylaxis in critically ill patients: A systematic review. <i>Thrombosis Research</i> , 2016, 139, 10-16.	0.8	15
36	Association of oliguria with the development of acute kidney injury in the critically ill. <i>Kidney International</i> , 2016, 89, 200-208.	2.6	52

#	ARTICLE	IF	CITATIONS
37	Soluble CD73 in Critically Ill Septic Patients â€” Data from the Prospective FINNAKI Study. PLoS ONE, 2016, 11, e0164420.	1.1	7
38	Predicting one-year mortality of critically ill patients with early acute kidney injury: data from the prospective multicenter FINNAKI study. Critical Care, 2015, 19, 125.	2.5	21
39	The Attributable Mortality of Acute Kidney Injury. Critical Care Medicine, 2014, 42, 878-885.	0.4	60
40	Timing of RRT Based on the Presence of Conventional Indications. Clinical Journal of the American Society of Nephrology: CJASN, 2014, 9, 1577-1585.	2.2	75
41	Incidence, risk factors and 90-day mortality of patients with acute kidney injury in Finnish intensive care units: the FINNAKI study. Intensive Care Medicine, 2013, 39, 420-428.	3.9	348
42	Hemodynamic variables and progression of acute kidney injury in critically ill patients with severe sepsis: data from the prospective observational FINNAKI study. Critical Care, 2013, 17, R295.	2.5	124
43	Acute kidney injury in patients with severe sepsis in Finnish intensive care units. Acta Anaesthesiologica Scandinavica, 2013, 57, 863-872.	0.7	102
44	Six-month survival and quality of life of intensive care patients with acute kidney injury. Critical Care, 2013, 17, R250.	2.5	42
45	Association of ICU size and annual case volume of renal replacement therapy patients with mortality. Acta Anaesthesiologica Scandinavica, 2012, 56, 1175-1182.	0.7	18
46	Population-based incidence, mortality and quality of life in critically ill patients treated with renal replacement therapy: a nationwide retrospective cohort study in Finnish intensive care units. Critical Care, 2012, 16, R13.	2.5	60
47	Fluid overload is associated with an increased risk for 90-day mortality in critically ill patients with renal replacement therapy: data from the prospective FINNAKI study. Critical Care, 2012, 16, R197.	2.5	308