

Dong Luo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5681489/publications.pdf>

Version: 2024-02-01

12
papers

311
citations

1163117

8
h-index

1199594

12
g-index

12
all docs

12
docs citations

12
times ranked

381
citing authors

#	ARTICLE	IF	CITATIONS
1	Geoclimatic factors influence the population genetic connectivity of <i>Incarvillea arguta</i> (Bignoniaceae) in the Himalaya–Hengduan Mountains biodiversity hotspot. <i>Journal of Systematics and Evolution</i> , 2021, 59, 151-168.	3.1	28
2	Molecular phylogeny, biogeography and character evolution of the montane genus <i>Incarvillea</i> Juss. (Bignoniaceae). <i>Plant Diversity</i> , 2021, 43, 1-14.	3.7	7
3	Estimating climate-induced “Nowhere to go” range shifts of the Himalayan <i>Incarvillea</i> Juss. using multi-model median ensemble species distribution models. <i>Ecological Indicators</i> , 2021, 121, 107127.	6.3	28
4	Biogeographical divides delineated by the three-step landforms of China and the East China Sea: Insights from the phylogeography of <i>Kerria japonica</i> . <i>Journal of Biogeography</i> , 2021, 48, 372-385.	3.0	18
5	<i>Saussurea talungensis</i> (Asteraceae), a new species from Humla, Nepal Himalayas. <i>PhytoKeys</i> , 2021, 176, 55-66.	1.0	1
6	Evolutionary radiations of cushion plants on the Qinghai–Tibet Plateau: Insights from molecular phylogenetic analysis of two subgenera of <i>Arenaria</i> and <i>Thylacospermum</i> (Caryophyllaceae). <i>Taxon</i> , 2019, 68, 1003-1020.	0.7	5
7	Geological and Climatic Factors Affect the Population Genetic Connectivity in <i>Mirabilis himalaica</i> (Nyctaginaceae): Insight From Phylogeography and Dispersal Corridors in the Himalaya-Hengduan Biodiversity Hotspot. <i>Frontiers in Plant Science</i> , 2019, 10, 1721.	3.6	21
8	Phylogeography of rare fern <i>Polystichum glaciale</i> endemic to the subnival zone of the Sino-Himalaya. <i>Plant Systematics and Evolution</i> , 2018, 304, 485-499.	0.9	11
9	Phylogeography of <i>Parasyncalathium souliei</i> (Asteraceae) and Its Potential Application in Delimiting Phylogeoregions in the Qinghai-Tibet Plateau (QTP)-Hengduan Mountains (HDM) Hotspot. <i>Frontiers in Genetics</i> , 2018, 9, 171.	2.3	16
10	The “Ward Line” Mekong–Salween Divide™ is an important floristic boundary between the eastern Himalaya and Hengduan Mountains: evidence from the phylogeographical structure of subnival herbs <i>Marmoritis complanatum</i> (Lamiaceae). <i>Botanical Journal of the Linnean Society</i> , 2017, 185, 482-496.	1.6	61
11	Evolutionary history of the subnival flora of the Himalaya–Hengduan Mountains: first insights from comparative phylogeography of four perennial herbs. <i>Journal of Biogeography</i> , 2016, 43, 31-43.	3.0	112
12	A karyological study of six species of <i>Silene</i> L. (Caryophyllaceae) from the Hengduan Mountains, SW China. <i>Caryologia</i> , 2011, 64, 10-13.	0.3	3