Hiroki Hayashi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5682060/publications.pdf

Version: 2024-02-01

| | | 1163117 | 1125743 |
|----------|----------------|--------------|----------------|
| 19 | 176 | 8 | 13 |
| papers | citations | h-index | g-index |
| | | | |
| | | | |
| | | | |
| 20 | 20 | 20 | 250 |
| all docs | docs citations | times ranked | citing authors |
| | | | |

| # | Article | IF | CITATIONS |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|-----------|
| 1 | Prescription rate of erythropoietin-stimulating agents is low for patients with renal impairment under non-nephrology care in a tertiary-level academic medical center in Japan. Clinical and Experimental Nephrology, 2022, , 1. | 1.6 | O |
| 2 | A nationwide analysis of renal and patient outcomes for adults with lupus nephritis in Japan. Clinical and Experimental Nephrology, 2022, 26, 898-908. | 1.6 | 3 |
| 3 | Vitamin K2 supplementation and the progression of abdominal aortic calcification in dialysis patients , 2021, 7, 136-138. | | 1 |
| 4 | Suppression of inflammation during cellâ€free concentrated ascites reinfusion therapy using a blood purification device. Therapeutic Apheresis and Dialysis, 2020, 24, 511-515. | 0.9 | 1 |
| 5 | Comparison of the 2018 and 2003 International Society of Nephrology/Renal Pathology Society classification in terms of renal prognosis in patients of lupus nephritis: a retrospective cohort study. Arthritis Research and Therapy, 2020, 22, 260. | 3. 5 | 14 |
| 6 | Long-term changes in renal function after treatment initiation and the importance of early diagnosis in maintaining renal function among IgG4-related tubulointerstitial nephritis patients in Japan. Arthritis Research and Therapy, 2020, 22, 261. | 3.5 | 8 |
| 7 | Incidence of remission and relapse of proteinuria, end-stage kidney disease, mortality, and major outcomes in primary nephrotic syndrome: the Japan Nephrotic Syndrome Cohort Study (JNSCS). Clinical and Experimental Nephrology, 2020, 24, 526-540. | 1.6 | 33 |
| 8 | Uric acid distribution volume calculated by kinetic modeling and extracellular volume predicted by bioimpedance method. International Journal of Artificial Organs, 2020, 43, 701-709. | 1.4 | 3 |
| 9 | Better remission rates in elderly Japanese patients with primary membranous nephropathy in nationwide real-world practice: The Japan Nephrotic Syndrome Cohort Study (JNSCS). Clinical and Experimental Nephrology, 2020, 24, 893-909. | 1.6 | 6 |
| 10 | Prediction model for cardiovascular events or all-cause mortality in incident dialysis patients. PLoS ONE, 2019, 14, e0221352. | 2.5 | 7 |
| 11 | Regional variations in immunosuppressive therapy in patients with primary nephrotic syndrome: the Japan nephrotic syndrome cohort study. Clinical and Experimental Nephrology, 2018, 22, 1266-1280. | 1.6 | 21 |
| 12 | Ratio of blood urea nitrogen to serum creatinine at initiation of dialysis is associated with mortality: a multicenter prospective cohort study. Clinical and Experimental Nephrology, 2018, 22, 353-364. | 1.6 | 16 |
| 13 | Aortic stenosis is a risk factor for all-cause mortality in patients on dialysis: a multicenter prospective cohort analysis. BMC Nephrology, 2018, 19, 80. | 1.8 | 4 |
| 14 | Impact of high mortality in incident dialysis patients due to hypertensive nephrosclerosis: a multicenter prospective cohort study in Aichi, Japan. Clinical and Experimental Nephrology, 2018, 22, 1360-1370. | 1.6 | 5 |
| 15 | Efficacy of urinary midkine as a biomarker in patients with acute kidney injury. Clinical and Experimental Nephrology, 2017, 21, 597-607. | 1.6 | 13 |
| 16 | Tubulointerstitial fibrosis in patients with IgG4-related kidney disease: pathological findings on repeat renal biopsy. Rheumatology International, 2015, 35, 1093-1101. | 3.0 | 13 |
| 17 | Efficacy and safety of febuxostat in elderly female patients. Clinical Interventions in Aging, 2014, 9, 1489. | 2.9 | 7 |
| 18 | CD147/basigin reflects renal dysfunction in patients with acute kidney injury. Clinical and Experimental Nephrology, 2014, 18, 746-754. | 1.6 | 11 |

| # | Article | IF | CITATIONS |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Mycophenolate mofetil versus intravenous cyclophosphamide for induction treatment of proliferative lupus nephritis in a Japanese population: a retrospective study. Modern Rheumatology, 2013, 23, 89-96. | 1.8 | 10 |