Anne Lübbeke

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5700646/publications.pdf

Version: 2024-02-01

236925 223800 2,199 59 25 46 citations h-index g-index papers 60 60 60 2541 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Patient-reported outcome measures in arthroplasty registries. Monthly Notices of the Royal Astronomical Society: Letters, 2016, 87, 9-23.	3.3	202
2	Differences in outcomes of obese women and men undergoing primary total hip arthroplasty. Arthritis and Rheumatism, 2007, 57, 327-334.	6.7	133
3	Patient-reported outcome measures in arthroplasty registries. Monthly Notices of the Royal Astronomical Society: Letters, 2016, 87, 3-8.	3.3	133
4	Orthopaedic registries with patient-reported outcome measures. EFORT Open Reviews, 2019, 4, 357-367.	4.1	116
5	Risk factors for post-traumatic osteoarthritis of the ankle: an eighteen year follow-up study. International Orthopaedics, 2012, 36, 1403-1410.	1.9	111
6	Low incidence of haematogenous seeding to total hip and knee prostheses in patients with remote infections. Journal of Infection, 2009, 59, 337-345.	3.3	98
7	International variation in shoulder arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2017, 88, 592-599.	3.3	97
8	Primary and revision hip arthroplasty: 5-year outcomes and influence of age and comorbidity. Journal of Rheumatology, 2007, 34, 394-400.	2.0	93
9	Outcomes of obese and nonobese patients undergoing revision total hip arthroplasty. Arthritis and Rheumatism, 2008, 59, 738-745.	6.7	86
10	Adipokines correlate with pain in lower limb osteoarthritis: different associations in hip and knee. International Orthopaedics, 2014, 38, 2577-2583.	1.9	73
11	Results of Surgical Repair of Abductor Avulsion After Primary Total Hip Arthroplasty. Journal of Arthroplasty, 2008, 23, 694-698.	3.1	69
12	Clinical Outcomes and Development of Symptomatic Osteoarthritis 2 to 24 Years After Surgical Treatment of Tarsometatarsal Joint Complex Injuries. Journal of Bone and Joint Surgery - Series A, 2016, 98, 713-720.	3.0	66
13	Body mass and weight thresholds for increased prosthetic joint infection rates after primary total joint arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2016, 87, 132-138.	3.3	63
14	Presence of IL-17 in synovial fluid identifies a potential inflammatory osteoarthritic phenotype. PLoS ONE, 2017, 12, e0175109.	2.5	61
15	Influence of preoperative patient education on the risk of dislocation after primary total hip arthroplasty. Arthritis and Rheumatism, 2009, 61, 552-558.	6.7	53
16	Activity and impact on antibiotic use and costs of a dedicated infectious diseases consultant on a septic orthopaedic unit. Journal of Infection, 2009, 58, 205-212.	3.3	51
17	Do synovial leptin levels correlate with pain in end stage arthritis?. International Orthopaedics, 2013, 37, 2071-2079.	1.9	51
18	Is There an Association Between Smoking Status and Prosthetic Joint Infection After Primary Total Joint Arthroplasty?. Journal of Arthroplasty, 2018, 33, 2218-2224.	3.1	45

#	Article	IF	CITATIONS
19	Physical Activity Before and After Primary Total Hip Arthroplasty: A Registryâ€Based Study. Arthritis Care and Research, 2014, 66, 277-284.	3.4	37
20	Cup Size and Risk of Dislocation After Primary Total Hip Arthroplasty. Journal of Arthroplasty, 2011, 26, 1305-1309.	3.1	34
21	Clinical and radiographic predictors of acute compartment syndrome in the treatment of tibial plateau fractures: a retrospective cohort study. BMC Musculoskeletal Disorders, 2017, 18, 307.	1.9	31
22	Influence of Obesity on Femoral Osteolysis Five and Ten Years Following Total Hip Arthroplasty. Journal of Bone and Joint Surgery - Series A, 2010, 92, 1964-1972.	3.0	30
23	Effectiveness of the Surgical Safety Checklist in a High Standard Care Environment. Medical Care, 2013, 51, 425-429.	2.4	30
24	BMI and Severity of Clinical and Radiographic Signs of Hip Osteoarthritis. Obesity, 2009, 17, 1414-1419.	3.0	27
25	The role of national registries in improving patient safety for hip and knee replacements. BMC Musculoskeletal Disorders, 2017, 18, 414.	1.9	27
26	Influence of body mass index on revision rates after primary total knee arthroplasty. International Orthopaedics, 2016, 40, 723-729.	1.9	26
27	Influence of Body Mass Index on Sagittal Knee Range of Motion and Gait Speed Recovery 1-Year After Total Knee Arthroplasty. Journal of Arthroplasty, 2017, 32, 2404-2410.	3.1	26
28	Third-generation pure alumina and alumina matrix composites in total hip arthroplasty. EFORT Open Reviews, 2018, 3, 7-14.	4.1	24
29	MoM total hip replacements in Europe: a NORE report. EFORT Open Reviews, 2019, 4, 423-429.	4.1	24
30	ASA class is associated with early revision and reoperation after total hip arthroplasty: an analysis of the Geneva and Swedish Hip Arthroplasty Registries. Monthly Notices of the Royal Astronomical Society: Letters, 2019, 90, 324-330.	3.3	23
31	Important patient characteristics differ prior to total knee arthroplasty and total hip arthroplasty between Switzerland and the United States. BMC Musculoskeletal Disorders, 2017, 18, 14.	1.9	22
32	Low risk despite high endemicity of methicillin-resistantStaphylococcus aureusinfections following elective total joint arthroplasty: A 12-year experience. Annals of Medicine, 2012, 44, 360-368.	3.8	21
33	Revision Total Hip Arthroplasty in Patients 80 Years or Older. Journal of Arthroplasty, 2012, 27, 1041-1046.	3.1	20
34	Improvement in mental health following total hip arthroplasty: the role of pain and function. BMC Musculoskeletal Disorders, 2019, 20, 307.	1.9	18
35	Ten-year results with the Morscher press-fit cup: an uncemented, non-modular, porous-coated cup inserted without screws. International Orthopaedics, 2011, 35, 957-963.	1.9	16
36	Strong association between smoking and the risk of revision in a cohort study of patients with metalâ€onâ€metal total hip arthroplasty. Journal of Orthopaedic Research, 2014, 32, 762-768.	2.3	16

#	Article	IF	CITATIONS
37	Registry stakeholders. EFORT Open Reviews, 2019, 4, 330-336.	4.1	15
38	Clinical and radiographic predictors of acute compartment syndrome in the treatment of tibial shaft fractures: a retrospective cohort study. BMC Musculoskeletal Disorders, 2020, 21, 25.	1.9	14
39	Associations between gait analysis parameters and patient satisfaction one year following primary total knee arthroplasty. Gait and Posture, 2020, 80, 44-48.	1.4	12
40	Three-Year Rates of Reoperation and Revision Following Mobile Versus Fixed-Bearing Total Ankle Arthroplasty. Journal of Bone and Joint Surgery - Series A, 2021, 103, 2080-2088.	3.0	12
41	Statins may reduce femoral osteolysis in patients with total Hip arthroplasty. Journal of Orthopaedic Research, 2013, 31, 814-820.	2.3	11
42	Research methodology for orthopaedic surgeons, with a focus on outcome. EFORT Open Reviews, 2018, 3, 160-167.	4.1	11
43	Development of a model predicting non-satisfaction 1 year after primary total knee replacement in the UK and transportation to Switzerland. Scientific Reports, 2018, 8, 3380.	3.3	10
44	Operatively treated ankle fractures in Switzerland, 2002–2012: epidemiology and associations between baseline characteristics and fracture types. BMC Musculoskeletal Disorders, 2021, 22, 266.	1.9	10
45	Walking Speed and Maximal Knee Flexion During Gait After Total Knee Arthroplasty: Minimal Clinically Important Improvement Is Not Determinable; Patient Acceptable Symptom State Is Potentially Useful. Journal of Arthroplasty, 2020, 35, 2865-2871.e2.	3.1	7
46	International variation in distribution of ASA class in patients undergoing total hip arthroplasty and its influence on mortality: data from an international consortium of arthroplasty registries. Monthly Notices of the Royal Astronomical Society: Letters, 2021, 92, 304-310.	3.3	7
47	The effect of BMI on long-term outcomes after operatively treated ankle fractures: a studyÂwith up to 16 years of follow-up. BMC Musculoskeletal Disorders, 2022, 23, 317.	1.9	6
48	Short stem total hip arthroplasty with the direct anterior approach demonstrates suboptimal fixation. International Orthopaedics, 2021, 45, 575-583.	1.9	5
49	Obesity in total hip arthroplasty—does it really matter?. Monthly Notices of the Royal Astronomical Society: Letters, 2012, 83, 99-100.	3.3	4
50	History of previous surgery is associated with higher risk of revision after primary total knee arthroplasty: a cohort study from the Geneva Arthroplasty Registry. Monthly Notices of the Royal Astronomical Society: Letters, 2021, 92, 709-715.	3.3	4
51	1-year trajectories of patients undergoing primary total hip arthroplasty: Patient reported outcomes and resource needs according to education level. BMC Musculoskeletal Disorders, 2022, 23, 84.	1.9	4
52	Factors related to outcome of early and delayed prosthetic joint infections. Journal of Infection, 2016, 72, 255-257.	3.3	3
53	Development and psychometric performance of the French language version of the Manchester-Oxford Foot Questionnaire (MOXFQ). Foot and Ankle Surgery, 2020, 26, 902-906.	1.7	3
54	Acetabular Peri-Prosthetic Fractures—A Narrative Review. Medicina (Lithuania), 2022, 58, 630.	2.0	3

Anne LÃ1⁄4bbeke

#	Article	IF	CITATIONS
55	Gait Analysis 1 Year after Primary TKA: No Difference between Gap Balancing and Measured Resection Technique. Journal of Knee Surgery, 2019, 34, 898-905.	1.6	2
56	Feasibility and sustainability of working in different types of jobs after total hip arthroplasty: analysis of longitudinal data from two cohorts. Occupational and Environmental Medicine, 2022, 79, 486-493.	2.8	1
57	Reply to the comments of Kancherla et al. to the article: Prospective randomised study comparing screw versus helical blade in the treatment of low-energy trochanteric fractures. International Orthopaedics, 2012, 36, 1111-1111.	1.9	0
58	Uncemented science at its best!. Monthly Notices of the Royal Astronomical Society: Letters, 2020, 91, 228-229.	3.3	0
59	The case for an academic discipline of medical device science. EFORT Open Reviews, 2021, 6, 160-163.	4.1	0