

Hongke Zhou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5712079/publications.pdf>

Version: 2024-02-01

18
papers

782
citations

623734

14
h-index

839539

18
g-index

18
all docs

18
docs citations

18
times ranked

690
citing authors

#	ARTICLE	IF	CITATIONS
1	Excellent and reversible mechanofluorochromism in donor-acceptor π -systems based on bisarylic methanone derivatives. <i>Dyes and Pigments</i> , 2022, 198, 109983.	3.7	7
2	Phenothiazine and diphenylsulfone-based donor-acceptor π -systems exhibiting remarkable mechanofluorochromism. <i>Dyes and Pigments</i> , 2021, 184, 108868.	3.7	21
3	D π -A type luminophores with a twisted molecular conformation constructed by phenoxazine and diphenylsulfone showing high contrast mechanofluorochromism. <i>New Journal of Chemistry</i> , 2020, 44, 17882-17890.	2.8	16
4	Aggregation-induced enhanced emission-type cruciform luminophore constructed by carbazole exhibiting mechanical force-induced luminescent enhancement and chromism. <i>RSC Advances</i> , 2020, 10, 12025-12034.	3.6	12
5	Mechanochromic luminescence of AIEE-active tetraphenylethene-containing cruciform luminophores. <i>Dyes and Pigments</i> , 2019, 171, 107739.	3.7	24
6	Reversible solid-state mechanochromic luminescence originated from aggregation-induced enhanced emission-active Donor-Acceptor cruciform luminophores containing triphenylamine. <i>Dyes and Pigments</i> , 2019, 171, 107689.	3.7	22
7	Tetraphenylethene-containing cruciform luminophores with aggregation-induced emission and mechanoresponsive behavior. <i>Dyes and Pigments</i> , 2019, 170, 107606.	3.7	9
8	Twisted Donor-Acceptor Cruciform Luminophores Possessing Substituent-Dependent Properties of Aggregation-Induced Emission and Mechanofluorochromism. <i>Journal of Physical Chemistry C</i> , 2018, 122, 2297-2306.	3.1	98
9	Tetraphenylethene modified β^2 -ketoiminate boron complexes bearing aggregation-induced emission and mechanofluorochromism. <i>RSC Advances</i> , 2017, 7, 1348-1356.	3.6	35
10	Tetraphenylethene-based β^2 -diketonate boron complex: Efficient aggregation-induced emission and high contrast mechanofluorochromism. <i>Dyes and Pigments</i> , 2017, 139, 157-165.	3.7	76
11	Triphenylamine functionalized β^2 -Ketoiminate boron complex exhibiting aggregation-induced emission and mechanofluorochromism. <i>Dyes and Pigments</i> , 2017, 137, 200-207.	3.7	79
12	Effects of cyano groups on the properties of thiazole-based β^2 -ketoiminate boron complexes: aggregation-induced emission and mechanofluorochromism. <i>RSC Advances</i> , 2016, 6, 69560-69568.	3.6	43
13	Scaffold-like 3D networks fabricated via the organogelation of β^2 -diketone-boron for fluorescent sensing organic amine vapors. <i>Science Bulletin</i> , 2012, 57, 4264-4271.	1.7	17
14	Solvent-dependent photophysical and anion responsive properties of one glutamide gelator. <i>Soft Matter</i> , 2011, 7, 8296.	2.7	49
15	Low-dimensional nanostructures fabricated from bis(dioxaborine)carbazole derivatives as fluorescent chemosensors for detecting organic amine vapors. <i>Journal of Materials Chemistry</i> , 2011, 21, 8756.	6.7	124
16	New dendritic gelator bearing carbazole in each branching unit: selected response to fluoride ion in gel phase. <i>Organic and Biomolecular Chemistry</i> , 2011, 9, 1523.	2.8	70
17	Luminescent Organic 1D Nanomaterials Based on Bis(β^2 -diketone)carbazole Derivatives. <i>Chemistry - A European Journal</i> , 2011, 17, 1660-1669.	3.3	75
18	Subporphyrins with Monodisperse Oligocarbazole Arms. <i>European Journal of Organic Chemistry</i> , 2009, 2009, 53-60.	2.4	5