

Yvette I Sheline

List of Publications by Citations

Source: <https://exaly.com/author-pdf/5712230/yvette-i-sheline-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

54
papers

7,586
citations

32
h-index

71
g-index

71
ext. papers

9,171
ext. citations

7.3
avg, IF

6.01
L-index

#	Paper	IF	Citations
54	The default mode network and self-referential processes in depression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 1942-7	11.5	981
53	Untreated depression and hippocampal volume loss. <i>American Journal of Psychiatry</i> , 2003 , 160, 1516-8	11.9	954
52	Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 11020-5	11.5	838
51	Neuroimaging studies of mood disorder effects on the brain. <i>Biological Psychiatry</i> , 2003 , 54, 338-52	7.9	491
50	Amyloid plaques disrupt resting state default mode network connectivity in cognitively normal elderly. <i>Biological Psychiatry</i> , 2010 , 67, 584-7	7.9	421
49	Amygdala core nuclei volumes are decreased in recurrent major depression. <i>NeuroReport</i> , 1998 , 9, 2023-8	7.9	387
48	Altered emotional interference processing in affective and cognitive-control brain circuitry in major depression. <i>Biological Psychiatry</i> , 2008 , 63, 377-84	7.9	374
47	APOE4 allele disrupts resting state fMRI connectivity in the absence of amyloid plaques or decreased CSF A β 2. <i>Journal of Neuroscience</i> , 2010 , 30, 17035-40	6.6	340
46	Resting state functional connectivity in preclinical Alzheimer's disease. <i>Biological Psychiatry</i> , 2013 , 74, 340-7	7.9	302
45	Cognitive function in late life depression: relationships to depression severity, cerebrovascular risk factors and processing speed. <i>Biological Psychiatry</i> , 2006 , 60, 58-65	7.9	288
44	Harmonization of cortical thickness measurements across scanners and sites. <i>NeuroImage</i> , 2018 , 167, 104-120	7.9	286
43	Support for the vascular depression hypothesis in late-life depression: results of a 2-site, prospective, antidepressant treatment trial. <i>Archives of General Psychiatry</i> , 2010 , 67, 277-85		233
42	Regional white matter hyperintensity burden in automated segmentation distinguishes late-life depressed subjects from comparison subjects matched for vascular risk factors. <i>American Journal of Psychiatry</i> , 2008 , 165, 524-32	11.9	157
41	Common and Dissociable Dysfunction of the Reward System in Bipolar and Unipolar Depression. <i>Neuropsychopharmacology</i> , 2015 , 40, 2258-68	8.7	149
40	Statistical harmonization corrects site effects in functional connectivity measurements from multi-site fMRI data. <i>Human Brain Mapping</i> , 2018 , 39, 4213-4227	5.9	117
39	An antidepressant decreases CSF A β production in healthy individuals and in transgenic AD mice. <i>Science Translational Medicine</i> , 2014 , 6, 236re4	17.5	111
38	Depression and coronary heart disease: a review for cardiologists. <i>Clinical Cardiology</i> , 1997 , 20, 196-200	3.3	110

37	Greater loss of 5-HT(2A) receptors in midlife than in late life. <i>American Journal of Psychiatry</i> , 2002 , 159, 430-5	11.9	104
36	Efficacy and Safety of Low-field Synchronized Transcranial Magnetic Stimulation (sTMS) for Treatment of Major Depression. <i>Brain Stimulation</i> , 2015 , 8, 787-94	5.1	89
35	Depression and the hippocampus: cause or effect?. <i>Biological Psychiatry</i> , 2011 , 70, 308-9	7.9	85
34	Childhood trauma history is linked to abnormal brain connectivity in major depression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 8582-8590	11.5	66
33	Treatment course with antidepressant therapy in late-life depression. <i>American Journal of Psychiatry</i> , 2012 , 169, 1185-93	11.9	65
32	Decreased hippocampal 5-HT(2A) receptor binding in older depressed patients using [18F]altanserin positron emission tomography. <i>Neuropsychopharmacology</i> , 2004 , 29, 2235-41	8.7	63
31	Meta-Analysis of the Antidepressant Effects of Acute Sleep Deprivation. <i>Journal of Clinical Psychiatry</i> , 2017 , 78, e1020-e1034	4.6	59
30	Cognitive behavioral therapy increases amygdala connectivity with the cognitive control network in both MDD and PTSD. <i>NeuroImage: Clinical</i> , 2017 , 14, 464-470	5.3	54
29	Parsing the Hippocampus in Depression: Chronic Stress, Hippocampal Volume, and Major Depressive Disorder. <i>Biological Psychiatry</i> , 2019 , 85, 436-438	7.9	53
28	A Trial of Sertraline or Cognitive Behavior Therapy for Depression in Epilepsy. <i>Annals of Neurology</i> , 2019 , 86, 552-560	9.4	45
27	Redundant Gs-coupled serotonin receptors regulate amyloid- β metabolism in vivo. <i>Molecular Neurodegeneration</i> , 2016 , 11, 45	19	43
26	Imaging biomarkers associated with cognitive decline: a review. <i>Biological Psychiatry</i> , 2015 , 77, 685-92	7.9	39
25	Effects of traumatic brain injury and posttraumatic stress disorder on Alzheimer's disease in veterans, using the Alzheimer's Disease Neuroimaging Initiative. <i>Alzheimer's and Dementia</i> , 2014 , 10, S226-35	1.2	39
24	Amyloid burden in cognitively normal elderly is associated with preferential hippocampal subfield volume loss. <i>Journal of Alzheimer's Disease</i> , 2015 , 45, 27-33	4.3	33
23	Longitudinal ComBat: A method for harmonizing longitudinal multi-scanner imaging data. <i>NeuroImage</i> , 2020 , 220, 117129	7.9	32
22	F-Flortaucipir PET/MRI Correlations in Nonamnesic and Amnesic Variants of Alzheimer Disease. <i>Journal of Nuclear Medicine</i> , 2018 , 59, 299-306	8.9	32
21	Cognitive Behavioral Therapy Is Associated With Enhanced Cognitive Control Network Activity in Major Depression and Posttraumatic Stress Disorder. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2018 , 3, 311-319	3.4	23
20	Comparison of brain structural variables, neuropsychological factors, and treatment outcome in early-onset versus late-onset late-life depression. <i>American Journal of Geriatric Psychiatry</i> , 2014 , 22, 1039-46	6.5	22

19	Network changes associated with transdiagnostic depressive symptom improvement following cognitive behavioral therapy in MDD and PTSD. <i>Molecular Psychiatry</i> , 2018 , 23, 2314-2323	15.1	20
18	Antidepressant response to aripiprazole augmentation associated with enhanced FDOPA utilization in striatum: a preliminary PET study. <i>Psychiatry Research - Neuroimaging</i> , 2014 , 221, 231-9	2.9	17
17	Severe hippocampal atrophy is not associated with depression in temporal lobe epilepsy. <i>Epilepsy and Behavior</i> , 2014 , 34, 9-14	3.2	11
16	Effect of escitalopram dose and treatment duration on CSF A β levels in healthy older adults: A controlled clinical trial. <i>Neurology</i> , 2020 , 95, e2658-e2665	6.5	11
15	Effect of escitalopram on A β levels and plaque load in an Alzheimer mouse model. <i>Neurology</i> , 2020 , 95, e2666-e2674	6.5	9
14	Patients with anxiety disorders rely on bilateral dlPFC activation during verbal working memory. <i>Social Cognitive and Affective Neuroscience</i> , 2020 , 15, 1288-1298	4	5
13	Reply to comment on "An antidepressant decreases CSF A β production in healthy individuals and in transgenic AD mice". <i>Science Translational Medicine</i> , 2014 , 6, 268lr4	17.5	4
12	Structural brain measures linked to clinical phenotypes in major depression replicate across clinical centres. <i>Molecular Psychiatry</i> , 2021 , 26, 2764-2775	15.1	4
11	Dimensional connectomics of anxious misery, a human connectome study related to human disease: Overview of protocol and data quality. <i>NeuroImage: Clinical</i> , 2020 , 28, 102489	5.3	3
10	Linking antidepressant performance with pain network connectivity. <i>Lancet Psychiatry</i> , 2019 , 6, 635-636	6.3	3
9	IC-01-02: Longitudinal PIB PET imaging of the appearance and accumulation of beta-amyloid in cognitively normal middle and late life adults 2010 , 6, S2-S2		3
8	Cloud-Based Functional Magnetic Resonance Imaging Neurofeedback to Reduce the Negative Attentional Bias in Depression: A Proof-of-Concept Study. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2021 , 6, 490-497	3.4	3
7	Affect and neural activity in women with PTSD during a task of emotional interference. <i>Journal of Affective Disorders</i> , 2016 , 204, 9-15	6.6	2
6	Longitudinal ComBat: A Method for Harmonizing Longitudinal Multi-scanner Imaging Data		2
5	Combining transcranial magnetic stimulation with functional magnetic resonance imaging for probing and modulating neural circuits relevant to affective disorders. <i>Wiley Interdisciplinary Reviews: Cognitive Science</i> , 2021 , 12, e1553	4.5	2
4	Convergence, preliminary findings and future directions across the four human connectome projects investigating mood and anxiety disorders. <i>NeuroImage</i> , 2021 , 245, 118694	7.9	1
3	Differential impact of transdiagnostic, dimensional psychopathology on multiple scales of functional connectivity		1
2	Characterizing Heterogeneity in Neuroimaging, Cognition, Clinical Symptoms, and Genetics Among Patients With Late-Life Depression.. <i>JAMA Psychiatry</i> , 2022 ,	14.5	1

- 1 No increase in inflammation in late-life major depression screened to exclude physical illness.. *Translational Psychiatry*, **2022**, 12, 118 8.6 o