Carsten Hjorthj

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/5718751/carsten-hjorthoj-publications-by-year.pdf

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

135
papers3,820
citations34
h-index58
g-index148
ext. papers4,982
ext. citations6
avg, IF6.11
L-index

#	Paper	IF	Citations
135	Group cognitive behavioural therapy with virtual reality exposure versus group cognitive behavioural therapy with in vivo exposure for social anxiety disorder and agoraphobia: a protocol for a randomised clinical trial <i>BMJ Open</i> , 2022 , 12, e051147	3	O
134	The association between birth weight, ponderal index, psychotropic medication, and type 2 diabetes in individuals with severe mental illness <i>Journal of Diabetes and Its Complications</i> , 2022 , 10818	3 ^{3.2}	О
133	Consensus paper of the WFSBP task force on cannabis, cannabinoids and psychosis <i>World Journal of Biological Psychiatry</i> , 2022 , 1-24	3.8	1
132	The Danish High-Risk and Resilience Study-VIA 15 - A Study Protocol for the Third Clinical Assessment of a Cohort of 522 Children Born to Parents Diagnosed With Schizophrenia or Bipolar Disorder and Population-Based Controls <i>Frontiers in Psychiatry</i> , 2022 , 13, 809807	5	О
131	Vaccination against SARS-CoV-2 infection among vulnerable and marginalised population groups in Denmark: A nationwide population-based study <i>Lancet Regional Health - Europe, The</i> , 2022 , 16, 100355		O
130	People exposed to suicide attempts: Frequency, impact, and the support received. <i>Suicide and Life-Threatening Behavior</i> , 2021 , 51, 467-477	3.9	2
129	A Qualitative Study on the Implementation of Flexible Assertive Community Treatment - an Integrated Community-based Treatment Model for Patients with Severe Mental Illness. <i>International Journal of Integrated Care</i> , 2021 , 21, 13	2	O
128	The effect of sodium-glucose transport protein 2 inhibitors on mortality and heart failure in randomized trials versus observational studies. <i>Diabetic Medicine</i> , 2021 , 38, e14600	3.5	O
127	Mortality in substance-induced psychosis: a register-based national cohort study. <i>Addiction</i> , 2021 , 116, 3515-3524	4.6	O
126	Transdiagnostic versus Diagnosis-Specific Group Cognitive Behavioral Therapy for Anxiety Disorders and Depression: A Randomized Controlled Trial. <i>Psychotherapy and Psychosomatics</i> , 2021 , 1-14	49.4	7
125	Association between neonatal homecare for preterm infants and incidence of severe postpartum depression in mothers. <i>Journal of Affective Disorders</i> , 2021 , 278, 453-459	6.6	O
124	Associations between duration of untreated psychosis and domains of positive and negative symptoms persist after 10 years of follow-up: A secondary analysis from the OPUS trial. <i>Schizophrenia Research</i> , 2021 , 228, 575-580	3.6	1
123	Are attenuated positive symptoms and cortisol levels associated?. <i>Schizophrenia Research</i> , 2021 , 228, 621-623	3.6	
122	The effect of flexible assertive community treatment in Denmark: a quasi-experimental controlled study. <i>Lancet Psychiatry,the</i> , 2021 , 8, 27-35	23.3	11
121	Annual incidence of cannabis-induced psychosis, other substance-induced psychoses and dually diagnosed schizophrenia and cannabis use disorder in Denmark from 1994 to 2016. <i>Psychological Medicine</i> , 2021 , 51, 617-622	6.9	9
120	No evidence of associations between genetic liability for schizophrenia and development of cannabis use disorder. <i>Psychological Medicine</i> , 2021 , 51, 479-484	6.9	7
119	Predictors of remission from the ultra-high risk state for psychosis. <i>Microbial Biotechnology</i> , 2021 , 15, 104-112	3.3	5

118	Generalized neurocognitive impairment in individuals at ultra-high risk for psychosis: The possible key role of slowed processing speed. <i>Brain and Behavior</i> , 2021 , 11, e01962	3.4	3
117	Treatment of difficult-to-treat depression - clinical guideline for selected interventions. <i>Nordic Journal of Psychiatry</i> , 2021 , 1-12	2.3	
116	Digital Shared Decision-Making Interventions in Mental Healthcare: A Systematic Review and Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2021 , 12, 691251	5	1
115	Development Over Time of the Population-Attributable Risk Fraction for Cannabis Use Disorder in Schizophrenia in Denmark. <i>JAMA Psychiatry</i> , 2021 , 78, 1013-1019	14.5	11
114	The Danish OPUS Early Intervention Services for First-Episode Psychosis: A Phase 4 Prospective Cohort Study With Comparison of Randomized Trial and Real-World Data. <i>American Journal of Psychiatry</i> , 2021 , 178, 941-951	11.9	3
113	Medical cannabis and cannabis-based medicine show both potential efficacy and potential harms: Cross-sectional comparison with controls on self-rated and interviewer-rated outcomes within the Danish pilot program on medical cannabis. <i>Complementary Therapies in Clinical Practice</i> , 2021 , 45, 1014.	3.5 7 6	1
112	Repetitive transcranial magnetic stimulation and transcranial direct current stimulation for auditory hallucinations in schizophrenia: Systematic review and meta-analysis. <i>Journal of Psychiatric Research</i> , 2021 , 143, 163-175	5.2	3
111	Substance-induced psychosis as a risk factor for unipolar depression or anxiety disorders-A nationwide register-based prospective cohort study. <i>Journal of Affective Disorders</i> , 2021 , 295, 960-966	6.6	Ο
110	Etetrahydrocannabinol: harmful even in low doses?. Lancet Psychiatry,the, 2020, 7, 296-297	23.3	2
109	Children of Patients Undergoing Psychiatric Treatment: An Investigation of Statutory Support Services After Referrals to Child Protection Services. <i>Frontiers in Psychiatry</i> , 2020 , 11, 527	5	0
108	Infections as a Risk Factor for and Prognostic Factor After Substance-Induced Psychoses. <i>American Journal of Psychiatry</i> , 2020 , 177, 335-341	11.9	4
107	Experiential negative symptoms are more predictive of real-life functional outcome than expressive negative symptoms in clinical high-risk states. <i>Schizophrenia Research</i> , 2020 , 218, 151-156	3.6	7
106	Associations between facial affect recognition and neurocognition in subjects at ultra-high risk for psychosis: A case-control study. <i>Psychiatry Research</i> , 2020 , 290, 112969	9.9	2
105	Basic symptoms influence real-life functioning and symptoms in individuals at high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2020 , 141, 231-240	6.5	3
104	Supplementary data for a focused review and meta-analysis of H-MRS studies on cerebral glutamate and GABA levels in high-risk of psychosis states. <i>Data in Brief</i> , 2020 , 28, 104920	1.2	1
103	Cerebral glutamate and GABA levels in high-risk of psychosis states: Alfocused review and meta-analysis of H-MRS studies. <i>Schizophrenia Research</i> , 2020 , 215, 38-48	3.6	21
102	Cerebral Glutamate and Gamma-Aminobutyric Acid Levels in Individuals at Ultra-high Risk for Psychosis and the Association With Clinical Symptoms and Cognition. <i>Biological Psychiatry:</i> Cognitive Neuroscience and Neuroimaging, 2020 , 5, 569-579	3.4	7
101	T18. EFFECTS OF COGNITIVE REMEDIATION ON WHITE MATTER IN INDIVIDUALS AT ULTRA-HIGH RISK FOR PSYCHOSIS IA RANDOMIZED, CONTROLLED CLINICAL TRIAL. <i>Schizophrenia Bulletin</i> , 2020 , 46, S237-S238	1.3	78

100	T46. THE EFFECT OF COMPREHENSIVE COGNITIVE REMEDIATION IN INDIVIDUALS AT ULTRA-HIGH RISK FOR PSYCHOSIS: A SINGLE-BLIND, RANDOMISED, CLINICAL TRIAL (FOCUS). <i>Schizophrenia Bulletin</i> , 2020 , 46, S249-S249	1.3	78
99	T83. SUBSTANCE-INDUCED PSYCHOSIS LINKED TO BOTH INFECTIONS AND SCHIZOPHRENIA. <i>Schizophrenia Bulletin</i> , 2020 , 46, S263-S263	1.3	78
98	M94. NO EVIDENCE OF ASSOCIATIONS BETWEEN GENETIC LIABILITY FOR SCHIZOPHRENIA RISK OF CANNABIS USE DISORDER. <i>Schizophrenia Bulletin</i> , 2020 , 46, S170-S170	1.3	78
97	S121. OPUS 20, 20 YEAR FOLLOW-UP ON THE OPUS TRIAL. Schizophrenia Bulletin, 2020 , 46, S81-S82	1.3	78
96	Investigating Cognitive and Clinical Predictors of Real-Life Functioning, Functional Capacity, and Quality of Life in Individuals at Ultra-High Risk for Psychosis. <i>Schizophrenia Bulletin Open</i> , 2020 , 1,	2.2	3
95	Effectiveness of cognitive remediation in the ultra-high risk state for psychosis. <i>World Psychiatry</i> , 2020 , 19, 401-402	14.4	7
94	Efficacy and safety of anti-inflammatory agents in treatment of psychotic disorders - A comprehensive systematic review and meta-analysis. <i>Brain, Behavior, and Immunity,</i> 2020 , 90, 364-380	16.6	25
93	Cognitive remediation plus standard treatment versus standard treatment alone for individuals at ultra-high risk of developing psychosis: Results of the FOCUS randomised clinical trial. <i>Schizophrenia Research</i> , 2020 , 224, 151-158	3.6	10
92	No Effects of Cognitive Remediation on Cerebral White Matter in Individuals at Ultra-High Risk for Psychosis-A Randomized Clinical Trial. <i>Frontiers in Psychiatry</i> , 2020 , 11, 873	5	4
91	Baseline measures of cerebral glutamate and GABA levels in individuals at ultrahigh risk for psychosis: Implications for clinical outcome after 12 months. <i>European Psychiatry</i> , 2020 , 63, e83	6	3
90	Interview and questionnaire assessment of cognitive impairment in subjects at ultra-high risk for psychosis: Associations with cognitive test performance, psychosocial functioning, and positive symptoms. <i>Psychiatry Research</i> , 2020 , 294, 113498	9.9	3
89	High suicide rates during psychiatric inpatient stay and shortly after discharge. <i>Acta Psychiatrica Scandinavica</i> , 2020 , 142, 355-365	6.5	10
88	Sex differences in trauma exposure and symptomatology in trauma-affected refugees. <i>Psychiatry Research</i> , 2020 , 293, 113445	9.9	7
87	Cannabidiol for cannabis use disorder: too high hopes?. <i>Lancet Psychiatry,the</i> , 2020 , 7, 838-839	23.3	3
86	Morbidity and Mortality in the Children and Young Adult Offspring of Parents With Schizophrenia or Affective Disorders-A Nationwide Register-Based Cohort Study in 2 Million Individuals. <i>Schizophrenia Bulletin</i> , 2020 , 46, 130-139	1.3	14
85	Prevalence, predictors, and patterns of mechanical restraint use for inpatients with dual diagnosis. <i>Perspectives in Psychiatric Care</i> , 2020 , 56, 20-27	2.2	1
84	Cannabis use disorders may protect against certain disorders of the digestive organs in people with schizophrenia but not in healthy controls. <i>Psychological Medicine</i> , 2020 , 50, 499-506	6.9	3
83	Lifestyle Interventions for Weight Management in People with Serious Mental Illness: A Systematic Review with Meta-Analysis, Trial Sequential Analysis, and Meta-Regression Analysis Exploring the Mediators and Moderators of Treatment Effects. <i>Psychotherapy and Psychosomatics</i> , 2019 , 88, 350-362	9.4	26

(2018-2019)

82	Polygenic risk for psychiatric disorder and singleness in patients with severe mental illness and controls. <i>Journal of Psychiatric Research</i> , 2019 , 119, 60-66	5.2	5
81	Genome-wide association study implicates CHRNA2 in cannabis use disorder. <i>Nature Neuroscience</i> , 2019 , 22, 1066-1074	25.5	60
80	Dual diagnosis and mechanical restraint - a register based study of 31,793 patients and 6562 episodes of mechanical restraint in the Capital region of Denmark from 2010-2014. <i>Nordic Journal of Psychiatry</i> , 2019 , 73, 169-177	2.3	1
79	Efficacy of anti-inflammatory treatment on major depressive disorder or depressive symptoms: meta-analysis of clinical trials. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 139, 404-419	6.5	143
78	Cognitive functioning following discontinuation of antipsychotic medication. A naturalistic sub-group analysis from the OPUS II trial. <i>Psychological Medicine</i> , 2019 , 49, 1138-1147	6.9	6
77	Irritable bowel symptoms, use of healthcare, costs, sickness and disability pension benefits: A long-term population-based study. <i>Scandinavian Journal of Public Health</i> , 2019 , 47, 867-875	3	13
76	Individual-Level Predictors for Becoming Homeless and Exiting Homelessness: a Systematic Review and Meta-analysis. <i>Journal of Urban Health</i> , 2019 , 96, 741-750	5.8	38
75	Risk of homelessness after discharge from psychiatric wards in Denmark: a nationwide register-based cohort study. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 140, 477-489	6.5	6
74	Schizophrenia is associated with increased risk of subsequent substance abuse diagnosis: A nation-wide population-based register study. <i>Addiction</i> , 2019 , 114, 2217-2226	4.6	9
73	Illness management and recovery: one-year follow-up of a randomized controlled trial in Danish community mental health centers: long-term effects on clinical and personal recovery. <i>BMC Psychiatry</i> , 2019 , 19, 65	4.2	11
72	Emotion recognition latency, but not accuracy, relates to real life functioning in individuals at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2019 , 210, 197-202	3.6	8
71	Streptococcal Infections and Exacerbations in PANDAS: A Systematic Review and Meta-analysis. <i>Pediatric Infectious Disease Journal</i> , 2019 , 38, 189-194	3.4	5
70	Association between alcohol and substance use disorders and psychiatric service use in patients with severe mental illness: a nationwide Danish register-based cohort study. <i>Psychological Medicine</i> , 2018 , 48, 2592-2600	6.9	17
69	Associations between clinical and psychosocial factors and metabolic and cardiovascular risk factors in overweight patients with schizophrenia spectrum disorders - Baseline and two-years findings from the CHANGE trial. <i>Schizophrenia Research</i> , 2018 , 199, 96-102	3.6	24
68	Polygenic Risk Scores, School Achievement, and Risk for Schizophrenia: A Danish Population-Based Study. <i>Biological Psychiatry</i> , 2018 , 84, 684-691	7.9	18
67	Association of Substance Use Disorders With Conversion From Schizotypal Disorder to Schizophrenia. <i>JAMA Psychiatry</i> , 2018 , 75, 733-739	14.5	14
66	Dietary patterns and physical activity in people with schizophrenia and increased waist circumference. <i>Schizophrenia Research</i> , 2018 , 199, 109-115	3.6	33
65	Participants' and staffs' evaluation of the Illness Management and Recovery program: a randomized clinical trial. <i>Journal of Mental Health</i> , 2018 , 27, 30-37	2.7	11

64	T123. PERSISTENT NEGATIVE SYMPTOMS IN FIRST EPISODE PSYCHOSIS: PREVALENCE, PREDICTORS AND PROGNOSIS. <i>Schizophrenia Bulletin</i> , 2018 , 44, S163-S163	1.3	78
63	O12.7. RISK OF PSYCHOSIS IN OFFSPRING OF PARENTS WITH A HISTORY OF HOMELESSNESS DURING CHILDHOOD AND ADOLESCENCE: A NATIONWIDE, REGISTER-BASED, COHORT STUDY. <i>Schizophrenia Bulletin</i> , 2018 , 44, S111-S111	1.3	78
62	Illness management and recovery: Clinical outcomes of a randomized clinical trial in community mental health centers. <i>PLoS ONE</i> , 2018 , 13, e0194027	3.7	12
61	Rates and Predictors of Conversion to Schizophrenia or Bipolar Disorder Following Substance-Induced Psychosis. <i>American Journal of Psychiatry</i> , 2018 , 175, 343-350	11.9	93
60	School performance from primary education in the adolescent offspring of parents with schizophrenia and bipolar disorder- a national, register-based study. <i>Psychological Medicine</i> , 2018 , 48, 1993-2000	6.9	18
59	Examining speed of processing of facial emotion recognition in individuals at ultra-high risk for psychosis: Associations with symptoms and cognition. <i>Schizophrenia Research</i> , 2018 , 195, 562-563	3.6	4
58	Homelessness as a predictor of mortality: an 11-year register-based cohort study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018 , 53, 63-75	4.5	15
57	Implementation of the safewards model to reduce the use of coercive measures in adult psychiatric inpatient units: An interrupted time-series analysis. <i>Journal of Psychiatric Research</i> , 2018 , 105, 147-152	5.2	18
56	Mortality in unipolar depression preceding and following chronic somatic diseases. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 138, 500-508	6.5	11
55	The Self-help Online against Suicidal thoughts (SOS) trial: study protocol for a randomized controlled trial. <i>Trials</i> , 2017 , 18, 45	2.8	9
54	Years of potential life lost and life expectancy in schizophrenia: a systematic review and meta-analysis. <i>Lancet Psychiatry,the</i> , 2017 , 4, 295-301	23.3	458
53	Association between alcohol, cannabis, and other illicit substance abuse and risk of developing schizophrenia: a nationwide population based register study. <i>Psychological Medicine</i> , 2017 , 47, 1668-16	7 ^{6.9}	42
52	Associations between substance use disorders and suicide or suicide attempts in people with mental illness: a Danish nation-wide, prospective, register-based study of patients diagnosed with schizophrenia, bipolar disorder, unipolar depression or personality disorder. <i>Addiction</i> , 2017 , 112, 1250-	4.6 ·1259	83
51	The effect of cognitive remediation in individuals at ultra-high risk for psychosis: a systematic review. <i>NPJ Schizophrenia</i> , 2017 , 3, 20	5.5	41
50	Effect on return to work or education of Individual Placement and Support modified for people with mood and anxiety disorders: results of a randomised clinical trial. <i>Occupational and Environmental Medicine</i> , 2017 , 74, 717-725	2.1	24
49	Stability and development of psychotic symptoms and the use of antipsychotic medication - long-term follow-up. <i>Psychological Medicine</i> , 2017 , 47, 2118-2129	6.9	14
48	Negative symptoms mediate the relationship between neurocognition and function in individuals at ultrahigh risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 135, 250-258	6.5	17
47	The effect of duration of untreated psychosis and treatment delay on the outcomes of prolonged early intervention in psychotic disorders. <i>NPJ Schizophrenia</i> , 2017 , 3, 34	5.5	45

(2016-2017)

Effect of lifestyle coaching versus care coordination versus treatment as usual in people with severe mental illness and overweight: Two-years follow-up of the randomized CHANGE trial. <i>PLoS ONE</i> , 2017 , 12, e0185881	3.7	19
TAILOR - tapered discontinuation versus maintenance therapy of antipsychotic medication in patients with newly diagnosed schizophrenia or persistent delusional disorder in remission of psychotic symptoms: study protocol for a randomized clinical trial. <i>Trials</i> , 2017 , 18, 445	2.8	14
Irritable bowel symptoms and the development of common mental disorders and functional somatic syndromes identified in secondary care - a long-term, population-based study. <i>Clinical Epidemiology</i> , 2017 , 9, 393-402	5.9	6
Exercise for patients with major depression: a systematic review with meta-analysis and trial sequential analysis. <i>BMJ Open</i> , 2017 , 7, e014820	3	62
Letter to the Editor. European Journal of Gastroenterology and Hepatology, 2017, 29, 615	2.2	
Antipsychotic medication and remission of psychotic symptoms 10years after a first-episode psychosis. <i>Schizophrenia Research</i> , 2017 , 182, 42-48	3.6	55
Course of illness in a sample of patients diagnosed with a schizotypal disorder and treated in a specialized early intervention setting. Findings from the 3.5year follow-up of the OPUS II study. <i>Schizophrenia Research</i> , 2017 , 182, 24-30	3.6	13
Risk of psychiatric disorders in offspring of parents with a history of homelessness during childhood and adolescence in Denmark: a nationwide, register-based, cohort study. <i>Lancet Public Health, The</i> , 2017 , 2, e541-e550	22.4	15
Five years of specialised early intervention versus two years of specialised early intervention followed by three years of standard treatment for patients with a first episode psychosis: randomised, superiority, parallel group trial in Denmark (OPUS II). <i>BMJ, The</i> , 2017 , 356, i6681	5.9	53
High loading of polygenic risk in cases with chronic schizophrenia. <i>Molecular Psychiatry</i> , 2016 , 21, 969-7	415.1	44
High loading of polygenic risk in cases with chronic schizophrenia. <i>Molecular Psychiatry</i> , 2016 , 21, 969-70. The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. <i>World Psychiatry</i> , 2016 , 15, 155-65	'415.1 14.4	93
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with		
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. <i>World Psychiatry</i> , 2016 , 15, 155-65 Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2016 ,	14.4	93
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. World Psychiatry, 2016, 15, 155-65 Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2016, 55, 953-961 Gastrointestinal symptoms related to the irritable bowel syndrome - a longitudinal	14.4 7.2	93
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. World Psychiatry, 2016, 15, 155-65 Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2016, 55, 953-961 Gastrointestinal symptoms related to the irritable bowel syndrome - a longitudinal population-based register study. Scandinavian Journal of Gastroenterology, 2016, 51, 420-6 Effectiveness of Mental Health First Aid training in Denmark: a randomized trial in waitlist design.	7.2	93 28 5
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. World Psychiatry, 2016, 15, 155-65 Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2016, 55, 953-961 Gastrointestinal symptoms related to the irritable bowel syndrome - a longitudinal population-based register study. Scandinavian Journal of Gastroenterology, 2016, 51, 420-6 Effectiveness of Mental Health First Aid training in Denmark: a randomized trial in waitlist design. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 597-606 Prevalence of substance use disorders in psychiatric patients: a nationwide Danish	7.2 2.4 4.5	9328521
The CHANGE trial: no superiority of lifestyle coaching plus care coordination plus treatment as usual compared to treatment as usual alone in reducing risk of cardiovascular disease in adults with schizophrenia spectrum disorders and abdominal obesity. World Psychiatry, 2016, 15, 155-65 Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2016, 55, 953-961 Gastrointestinal symptoms related to the irritable bowel syndrome - a longitudinal population-based register study. Scandinavian Journal of Gastroenterology, 2016, 51, 420-6 Effectiveness of Mental Health First Aid training in Denmark: a randomized trial in waitlist design. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 597-606 Prevalence of substance use disorders in psychiatric patients: a nationwide Danish population-based study. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 129-40 Research Review: Do motor deficits during development represent an endophenotype for schizophrenia? A meta-analysis. Journal of Child Psychology and Psychiatry and Allied Disciplines,	14.4 7.2 2.4 4.5	9328521103
	TAILOR - tapered discontinuation versus maintenance therapy of antipsychotic medication in patients with newly diagnosed schizophrenia or persistent delusional disorder in remission of psychotic symptoms: study protocol for a randomized clinical trial. <i>Trials</i> , 2017, 18, 445 Irritable bowel symptoms and the development of common mental disorders and functional somatic syndromes identified in secondary care - a long-term, population-based study. <i>Clinical Epidemiology</i> , 2017, 9, 393-402 Exercise for patients with major depression: a systematic review with meta-analysis and trial sequential analysis. <i>BMJ Open</i> , 2017, 7, e014820 Letter to the Editor. <i>European Journal of Gastroenterology and Hepatology</i> , 2017, 29, 615 Antipsychotic medication and remission of psychotic symptoms 10years after a first-episode psychosis. <i>Schizophrenia Research</i> , 2017, 182, 42-48 Course of illness in a sample of patients diagnosed with a schizotypal disorder and treated in a specialized early intervention setting. Findings from the 3.5year follow-up of the OPUS II study. <i>Schizophrenia Research</i> , 2017, 182, 24-30 Risk of psychiatric disorders in offspring of parents with a history of homelessness during childhood and adolescence in Denmark: a nationwide, register-based, cohort study. <i>Lancet Public Health</i> , <i>The</i> , 2017, 2, e541-e550 Five years of specialised early intervention versus two years of specialised early intervention followed by three years of standard treatment for patients with a first episode psychosis:	TAILOR - tapered discontinuation versus maintenance therapy of antipsychotic medication in patients with newly diagnosed schizophrenia or persistent delusional disorder in remission of psychotic symptoms: study protocol for a randomized clinical trial. <i>Trials, 2017, 18, 445</i> Irritable bowel symptoms and the development of common mental disorders and functional somatic syndromes identified in secondary care - a long-term, population-based study. <i>Clinical Epidemiology, 2017, 9, 393-402</i> Exercise for patients with major depression: a systematic review with meta-analysis and trial sequential analysis. <i>BMJ Open, 2017, 7,</i> e014820 3 Letter to the Editor. <i>European Journal of Gastroenterology and Hepatology, 2017, 29,</i> 615 2.2 Antipsychotic medication and remission of psychotic symptoms 10years after a first-episode psychosis. <i>Schizophrenia Research, 2017, 182, 42-48</i> Course of illness in a sample of patients diagnosed with a schizotypal disorder and treated in a specialized early intervention setting. Findings from the 3.5year follow-up of the OPUS II study. <i>Schizophrenia Research, 2017, 182, 24-30</i> Risk of psychiatric disorders in offspring of parents with a history of homelessness during childhood and adolescence in Denmark: a nationwide, register-based, cohort study. <i>Lancet Public Health, The, 2017, 2, e541-e550</i> Five years of specialised early intervention versus two years of specialised early intervention followed by three years of standard treatment for patients with a first episode psychosis: 5.9

28	Systemic oxidative DNA and RNA damage are not increased during early phases of psychosis: A case control study. <i>Psychiatry Research</i> , 2016 , 241, 201-6	9.9	17
27	Social cognition in patients at ultra-high risk for psychosis: What is the relation to social skills and functioning?. <i>Schizophrenia Research: Cognition</i> , 2016 , 5, 21-27	2.8	37
26	Ten-year follow-up of the OPUS specialized early intervention trial for patients with a first episode of psychosis. <i>Schizophrenia Bulletin</i> , 2015 , 41, 617-26	1.3	94
25	DEPERROR: Risks of systematic errors in drug and non-drug randomized clinical trials assessing intervention effects in patients with unipolar depression. <i>Journal of Affective Disorders</i> , 2015 , 179, 121-	7 6.6	5
24	From research to practice: how OPUS treatment was accepted and implemented throughout Denmark. <i>Microbial Biotechnology</i> , 2015 , 9, 156-62	3.3	60
23	Protocol for CHANGE: a randomized clinical trial assessing lifestyle coaching plus care coordination versus care coordination alone versus treatment as usual to reduce risks of cardiovascular disease in adults with schizophrenia and abdominal obesity. <i>BMC Psychiatry</i> , 2015 , 15, 119	4.2	10
22	Association between alcohol and substance use disorders and all-cause and cause-specific mortality in schizophrenia, bipolar disorder, and unipolar depression: a nationwide, prospective, register-based study. <i>Lancet Psychiatry,the</i> , 2015 , 2, 801-8	23.3	91
21	Prevention of depression in patients with acute coronary syndrome (DECARD) randomized trial: effects on and by self-reported health. <i>Microbial Biotechnology</i> , 2015 , 9, 370-7	3.3	1
20	Serious mental illness and disrupted caregiving for children: a nationwide, register-based cohort study. <i>Journal of Clinical Psychiatry</i> , 2015 , 76, e1006-14	4.6	28
19	Risk of suicide according to level of psychiatric treatment: a nationwide nested case-control study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014 , 49, 1357-65	4.5	64
18	Suicide and unintentional injury mortality among homeless people: a Danish nationwide register-based cohort study. <i>European Journal of Public Health</i> , 2014 , 24, 50-6	2.1	26
17	Predictors of recovery in first episode psychosis: the OPUS cohort at 10 year follow-up. <i>Schizophrenia Research</i> , 2013 , 150, 163-8	3.6	107
16	Test-retest reliability of the Danish Adult Reading Test in patients with comorbid psychosis and cannabis-use disorder. <i>Nordic Journal of Psychiatry</i> , 2013 , 67, 159-63	2.3	12
15	Psychiatric treatment following participation in the CapOpus randomized trial for patients with comorbid cannabis use disorder and psychosis. <i>Schizophrenia Research</i> , 2013 , 151, 191-6	3.6	8
14	Health care use by patients with somatoform disorders: a register-based follow-up study. <i>Psychosomatics</i> , 2013 , 54, 132-41	2.6	22
13	Specialized psychosocial treatment plus treatment as usual (TAU) versus TAU for patients with cannabis use disorder and psychosis: the CapOpus randomized trial. <i>Psychological Medicine</i> , 2013 , 43, 1499-510	6.9	39
12	Validity of Timeline Follow-Back for self-reported use of cannabis and other illicit substancessystematic review and meta-analysis. <i>Addictive Behaviors</i> , 2012 , 37, 225-33	4.2	229
11	Correlations and agreement between delta-9-tetrahydrocannabinol (THC) in blood plasma and timeline follow-back (TLFB)-assisted self-reported use of cannabis of patients with cannabis use disorder and psychotic illness attending the CapOpus randomized clinical trial. <i>Addiction</i> , 2012 , 107, 113	4.6 23-31	43

LIST OF PUBLICATIONS

10	Psychiatric disorders and mortality among people in homeless shelters in Denmark: a nationwide register-based cohort study. <i>Lancet, The</i> , 2011 , 377, 2205-14	40	125
9	Does the EQ-5D measure quality of life in schizophrenia?. <i>Journal of Mental Health Policy and Economics</i> , 2011 , 14, 187-96	1.3	7
8	Treatment of cannabis use disorders in people with schizophrenia spectrum disorders - a systematic review. <i>Addictive Behaviors</i> , 2009 , 34, 520-5	4.2	31
7	Reprint of "Treatment of cannabis use disorders in people with schizophrenia spectrum disordersa systematic review". <i>Addictive Behaviors</i> , 2009 , 34, 846-51	4.2	8
6	CAPOPUS Trial: an Observer-blinded RCT of Specialized Addiction Treatment Versus Standard Treatment for Young Patients with Cannabis Abuse and Psychosis. <i>European Psychiatry</i> , 2009 , 24, 1-1	6	1
5	Design paper: The CapOpus trial: a randomized, parallel-group, observer-blinded clinical trial of specialized addiction treatment versus treatment as usual for young patients with cannabis abuse and psychosis. <i>Trials</i> , 2008 , 9, 42	2.8	11
4	Cannabis use and risk of psychosis in later life. Lancet, The, 2007, 370, 293-4	40	15
3	Home environment of 11-year-old children born to parents with schizophrenia or bipolar disorder a controlled, 4-year follow-up study: The Danish High Risk and Resilience Study IVIA 11. <i>Psychological Medicine</i> ,1-11	6.9	
2	Rating the importance of outcomes from diabetes trials. A survey of patients and doctors opinions. <i>Journal of Diabetes and Metabolic Disorders</i> , 1	2.5	
1	Genome-wide association study implicates CHRNA2 in cannabis use disorder		5