

Andrew C Cogswell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5722640/publications.pdf>

Version: 2024-02-01

10
papers

338
citations

1307366

7
h-index

1474057

9
g-index

10
all docs

10
docs citations

10
times ranked

604
citing authors

#	ARTICLE	IF	CITATIONS
1	Presence of Natural Killer B Cells in Simian Immunodeficiency Virus-Infected Colon That Have Properties and Functions Similar to Those of Natural Killer Cells and B Cells but Are a Distinct Cell Population. <i>Journal of Virology</i> , 2022, 96, e0023522.	1.5	2
2	Contribution of Innate Lymphoid Cells in Supplementing Cytokines Produced by CD4+ T-cells During Acute and Chronic SIV Infection of the Colon. <i>AIDS Research and Human Retroviruses</i> , 2022, , .	0.5	0
3	Enteric bacteria induce IFN γ and Granzyme B from human colonic Group 1 Innate Lymphoid Cells. <i>Gut Microbes</i> , 2020, 12, 1667723.	4.3	15
4	Presence of Inflammatory Group I and III Innate Lymphoid Cells in the Colon of Simian Immunodeficiency Virus-Infected Rhesus Macaques. <i>Journal of Virology</i> , 2020, 94, .	1.5	7
5	Children with Narcolepsy type 1 have increased T α cell responses to orexins. <i>Annals of Clinical and Translational Neurology</i> , 2019, 6, 2566-2572.	1.7	17
6	Commensal and Pathogenic Bacteria Indirectly Induce IL-22 but Not IFN γ Production From Human Colonic ILC3s via Multiple Mechanisms. <i>Frontiers in Immunology</i> , 2019, 10, 649.	2.2	42
7	The neurobiological basis of narcolepsy. <i>Nature Reviews Neuroscience</i> , 2019, 20, 83-93.	4.9	151
8	CD155 on HIV-Infected Cells Is Not Modulated by HIV-1 Vpu and Nef but Synergizes with NKG2D Ligands to Trigger NK Cell Lysis of Autologous Primary HIV-Infected Cells. <i>AIDS Research and Human Retroviruses</i> , 2017, 33, 93-100.	0.5	21
9	Brief Report: Inflammatory Colonic Innate Lymphoid Cells Are Increased During Untreated HIV-1 Infection and Associated With Markers of Gut Dysbiosis and Mucosal Immune Activation. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2017, 76, 431-437.	0.9	16
10	A Conserved HIV-1-Derived Peptide Presented by HLA-E Renders Infected T-cells Highly Susceptible to Attack by NKG2A/CD94-Bearing Natural Killer Cells. <i>PLoS Pathogens</i> , 2016, 12, e1005421.	2.1	67