

Jens Matthiessen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5722927/publications.pdf>

Version: 2024-02-01

76
papers

6,392
citations

87723

38
h-index

88477

70
g-index

83
all docs

83
docs citations

83
times ranked

4789
citing authors

#	ARTICLE	IF	CITATIONS
1	Reply to: No freshwater-filled glacial Arctic Ocean. <i>Nature</i> , 2022, 602, E4-E6.	13.7	4
2	Reply to "Challenging the hypothesis of an arctic ocean lake during recent glacial episodes" by Hillaire-Marcel, <i>Journal of Quaternary Science</i> , 2022, 37, 568-571.	1.1	3
3	A revised core-seismic integration in the Molloy Basin (ODP Site 909): Implications for the history of ice rafting and ocean circulation in the Atlantic-Arctic gateway. <i>Global and Planetary Change</i> , 2022, 215, 103876.	1.6	3
4	Glacial episodes of a freshwater Arctic Ocean covered by a thick ice shelf. <i>Nature</i> , 2021, 590, 97-102.	13.7	32
5	Distribution of common modern dinoflagellate cyst taxa in surface sediments of the Northern Hemisphere in relation to environmental parameters: The new n=1968 database. <i>Marine Micropaleontology</i> , 2020, 159, 101796.	0.5	65
6	An overview and brief description of common marine organic-walled dinoflagellate cyst taxa occurring in surface sediments of the Northern Hemisphere. <i>Marine Micropaleontology</i> , 2020, 159, 101814.	0.5	45
7	Arctic Continental Margin Sediments as Possible Fe and Mn Sources to Seawater as Sea Ice Retreats: Insights From the Eurasian Margin. <i>Global Biogeochemical Cycles</i> , 2020, 34, e2020GB006581.	1.9	5
8	Natural variability of the Arctic Ocean sea ice during the present interglacial. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 26069-26075.	3.3	28
9	Benthic phosphorus cycling within the Eurasian marginal sea ice zone. <i>Philosophical Transactions Series A, Mathematical, Physical, and Engineering Sciences</i> , 2020, 378, 20190358.	1.6	6
10	Palynology, biostratigraphy, and paleoceanography of the Plio-Pleistocene at Ocean Drilling Program Site 887, Gulf of Alaska. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 546, 109605.	1.0	1
11	<i>Impagidinium detroitense</i> and <i>I. diaphanum</i> : Two new dinoflagellate cyst species from the Pliocene of the North Pacific Ocean, and their biostratigraphic significance. <i>Review of Palaeobotany and Palynology</i> , 2019, 264, 24-37.	0.8	5
12	Amino acid racemization in Quaternary foraminifera from the Yermak Plateau, Arctic Ocean. <i>Geochronology</i> , 2019, 1, 53-67.	1.0	11
13	Changes in sea ice cover and ice sheet extent at the Yermak Plateau during the last 160 ka "Reconstructions from biomarker records. <i>Quaternary Science Reviews</i> , 2018, 182, 93-108.	1.4	43
14	Distribution and (palaeo)ecological affinities of the main <i>Spiniferites</i> taxa in the mid-high latitudes of the Northern Hemisphere. <i>Palynology</i> , 2018, 42, 182-202.	0.7	16
15	Quaternary dinoflagellate cysts in the Arctic Ocean: Potential and limitations for stratigraphy and paleoenvironmental reconstructions. <i>Quaternary Science Reviews</i> , 2018, 192, 1-26.	1.4	15
16	Ballasting by cryogenic gypsum enhances carbon export in a <i>Phaeocystis</i> under-ice bloom. <i>Scientific Reports</i> , 2018, 8, 7703.	1.6	48
17	Improved Pleistocene sediment stratigraphy and paleoenvironmental implications for the western Arctic Ocean off the East Siberian and Chukchi margins. <i>Arktos</i> , 2018, 4, 1-20.	1.0	30
18	Neogene dinoflagellate cysts and acritarchs from the high northern latitudes and their relation to sea surface temperature. <i>Marine Micropaleontology</i> , 2017, 136, 51-65.	0.5	12

#	ARTICLE	IF	CITATIONS
19	Regional seesaw between the North Atlantic and Nordic Seas during the last glacial abrupt climate events. <i>Climate of the Past</i> , 2017, 13, 729-739.	1.3	10
20	Sea surface conditions in the southern Nordic Seas during the Holocene based on dinoflagellate cyst assemblages. <i>Holocene</i> , 2016, 26, 722-735.	0.9	21
21	Evidence for ice-free summers in the late Miocene central Arctic Ocean. <i>Nature Communications</i> , 2016, 7, 11148.	5.8	96
22	Dinoflagellates. <i>Encyclopedia of Earth Sciences Series</i> , 2016, , 189-193.	0.1	0
23	Early Pliocene onset of modern Nordic Seas circulation related to ocean gateway changes. <i>Nature Communications</i> , 2015, 6, 8659.	5.8	59
24	Dinoflagellates. , 2015, , 1-7.		1
25	<i>Batiacasphaera bergenensis</i> and <i>Lavradosphaera elongata</i> – New dinoflagellate cyst and acritarch species from the Miocene of the Iceland Sea (ODP Hole 907A). <i>Review of Palaeobotany and Palynology</i> , 2014, 211, 97-106.	0.8	3
26	Effect of early Pliocene uplift on late Pliocene cooling in the Arctic – Atlantic gateway. <i>Earth and Planetary Science Letters</i> , 2014, 387, 132-144.	1.8	71
27	Statistically assessing the correlation between salinity and morphology in cysts produced by the dinoflagellate <i>Protoceratium reticulatum</i> from surface sediments of the North Atlantic Ocean, Mediterranean – Marmara – Black Sea region, and Baltic – Kattegat – Skagerrak estuarine system. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 399, 202-213.	1.0	25
28	Changes in current patterns in the Fram Strait at the Pliocene/Pleistocene boundary. <i>Quaternary Science Reviews</i> , 2014, 92, 179-189.	1.4	14
29	Repeated Pleistocene glaciation of the East Siberian continental margin. <i>Nature Geoscience</i> , 2013, 6, 842-846.	5.4	140
30	Response of marine palynomorphs to Neogene climate cooling in the Iceland Sea (ODP Hole 907A). <i>Marine Micropaleontology</i> , 2013, 101, 49-67.	0.5	35
31	Glacial freshwater discharge events recorded by authigenic neodymium isotopes in sediments from the Mendeleev Ridge, western Arctic Ocean. <i>Earth and Planetary Science Letters</i> , 2013, 369-370, 148-157.	1.8	28
32	Atlas of modern dinoflagellate cyst distribution based on 2405 data points. <i>Review of Palaeobotany and Palynology</i> , 2013, 191, 1-197.	0.8	369
33	A magnetostratigraphic calibration of Middle Miocene through Pliocene dinoflagellate cyst and acritarch events in the Iceland Sea (Ocean Drilling Program Hole 907A). <i>Review of Palaeobotany and Palynology</i> , 2012, 187, 66-94.	0.8	55
34	Manganese-rich brown layers in Arctic Ocean sediments: Composition, formation mechanisms, and diagenetic overprint. <i>Geochimica Et Cosmochimica Acta</i> , 2011, 75, 7668-7687.	1.6	94
35	Deciphering the palaeoecology of Late Pliocene and Early Pleistocene dinoflagellate cysts. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2011, 309, 17-32.	1.0	52
36	Ice sheet grounding and iceberg plow marks on the northern and central Yermak Plateau revealed by geophysical data. <i>Quaternary Science Reviews</i> , 2011, 30, 1726-1738.	1.4	20

#	ARTICLE	IF	CITATIONS
37	Determining the absolute abundance of dinoflagellate cysts in recent marine sediments: The Lycopodium marker-grain method put to the test. <i>Review of Palaeobotany and Palynology</i> , 2009, 157, 238-252.	0.8	141
38	Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum. <i>Nature Geoscience</i> , 2009, 2, 127-132.	5.4	517
39	The Plio-Pleistocene glaciation of the Barents Sea – Svalbard region: a new model based on revised chronostratigraphy. <i>Quaternary Science Reviews</i> , 2009, 28, 812-829.	1.4	183
40	Pliocene palaeoceanography of the Arctic Ocean and subarctic seas. <i>Philosophical Transactions Series A, Mathematical, Physical, and Engineering Sciences</i> , 2009, 367, 21-48.	1.6	72
41	Biogenic barium in surface sediments of the European Nordic Seas. <i>Marine Geology</i> , 2008, 250, 89-103.	0.9	19
42	Last interglacial surface water conditions in the eastern Nordic Seas inferred from dinocyst and foraminiferal assemblages. <i>Marine Micropaleontology</i> , 2008, 66, 247-263.	0.5	24
43	Re-advance of the Fennoscandian Ice Sheet during Heinrich Event 1. <i>Marine Geology</i> , 2007, 240, 1-18.	0.9	37
44	Effects of Arctic freshwater forcing on thermohaline circulation during the Pleistocene. <i>Geology</i> , 2007, 35, 1075.	2.0	48
45	Last Interglacial Arctic warmth confirms polar amplification of climate change. <i>Quaternary Science Reviews</i> , 2006, 25, 1383-1400.	1.4	215
46	Subtropical Arctic Ocean temperatures during the Palaeocene/Eocene thermal maximum. <i>Nature</i> , 2006, 441, 610-613.	13.7	578
47	Episodic fresh surface waters in the Eocene Arctic Ocean. <i>Nature</i> , 2006, 441, 606-609.	13.7	284
48	The Cenozoic palaeoenvironment of the Arctic Ocean. <i>Nature</i> , 2006, 441, 601-605.	13.7	471
49	Modern organic-walled dinoflagellate cysts in arctic marine environments and their (paleo-) environmental significance. <i>Palaontologische Zeitschrift</i> , 2005, 79, 3-51.	0.8	98
50	Reconstruction of sea-surface conditions at middle to high latitudes of the Northern Hemisphere during the Last Glacial Maximum (LGM) based on dinoflagellate cyst assemblages. <i>Quaternary Science Reviews</i> , 2005, 24, 897-924.	1.4	283
51	Arctic (palaeo) river discharge and environmental change: evidence from the Holocene Kara Sea sedimentary record. <i>Quaternary Science Reviews</i> , 2004, 23, 1485-1511.	1.4	81
52	A multi-proxy study of Pliocene sediments from Ålde de France, North-East Greenland. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2002, 186, 1-23.	1.0	49
53	Norwegian sea-surface palaeoenvironments of marine oxygen-isotope stage 3: the paradoxical response of dinoflagellate cysts. <i>Journal of Quaternary Science</i> , 2002, 17, 349-359.	1.1	25
54	Magnetic susceptibility and ice-rafted debris in surface sediments of the Nordic Seas: implications for Isotope Stage 3 oscillations. <i>Geo-Marine Letters</i> , 2002, 22, 1-11.	0.5	29

#	ARTICLE	IF	CITATIONS
55	Evidence of 'Mid-Pliocene (~3 Ma) global warmth' in the eastern Arctic Ocean and implications for the Svalbard/Barents Sea ice sheet during the late Pliocene and early Pleistocene (~3-1.7 Ma). <i>Boreas</i> , 2002, 31, 82-93.	1.2	4
56	Evidence of 'Mid-Pliocene (~3 Ma) global warmth' in the eastern Arctic Ocean and implications for the Svalbard/Barents Sea ice sheet during the late Pliocene and early Pleistocene (~3-1.7 Ma). <i>Boreas</i> , 2002, 31, 82-93.	1.2	28
57	A multiproxy reconstruction of the evolution of deep and surface waters in the subarctic Nordic seas over the last 30,000yr. <i>Quaternary Science Reviews</i> , 2001, 20, 659-678.	1.4	183
58	Marine ice-rafted debris records constrain maximum extent of Saalian and Weichselian ice-sheets along the northern Eurasian margin. <i>Global and Planetary Change</i> , 2001, 31, 45-64.	1.6	96
59	Late Quaternary dinoflagellate cyst stratigraphy at the Eurasian continental margin, Arctic Ocean: indications for Atlantic water inflow in the past 150,000 years. <i>Global and Planetary Change</i> , 2001, 31, 65-86.	1.6	62
60	Late Holocene dinoflagellate cysts as indicators for short-term climate variability in the eastern Laptev Sea (Arctic Ocean). <i>Journal of Quaternary Science</i> , 2001, 16, 711-716.	1.1	13
61	Dinoflagellate cyst evidence for warm interglacial conditions at the northern Barents Sea margin during marine oxygen isotope stage 5. <i>Journal of Quaternary Science</i> , 2001, 16, 727-737.	1.1	42
62	Cold marine indicators of the late Quaternary: the new dinoflagellate cyst genus <i>Islandinium</i> and related morphotypes. <i>Journal of Quaternary Science</i> , 2001, 16, 621-636.	1.1	189
63	Dinoflagellate cyst assemblages as tracers of sea-surface conditions in the northern North Atlantic, Arctic and sub-Arctic seas: the new $n=677$ data base and its application for quantitative palaeoceanographic reconstruction. <i>Journal of Quaternary Science</i> , 2001, 16, 681-698.	1.1	303
64	Marine dinoflagellate cysts and high latitude Quaternary paleoenvironmental reconstructions: an introduction. <i>Journal of Quaternary Science</i> , 2001, 16, 595-602.	1.1	52
65	Distribution, Export and Alteration of Fossilizable Plankton in the Nordic Seas. , 2001, , 81-104.		20
66	The Potential of Synoptic Plankton Analyses for Paleoclimatic Investigations: Five Plankton Groups from the Holocene Nordic Seas. , 2001, , 291-318.		9
67	Distribution of Calcareous, Siliceous and Organic-Walled Planktic Microfossils in Surface Sediments of the Nordic Seas and their Relation to Surface-Water Masses. , 2001, , 105-127.		15
68	Freshwater chlorophycean algae in recent marine sediments of the Beaufort, Laptev and Kara Seas (Arctic Ocean) as indicators of river runoff. <i>International Journal of Earth Sciences</i> , 2000, 89, 470-485.	0.9	88
69	Reconstruction of sea-surface temperature, salinity, and sea-ice cover in the northern North Atlantic during the last glacial maximum based on dinocyst assemblages. <i>Canadian Journal of Earth Sciences</i> , 2000, 37, 725-750.	0.6	130
70	Organic-walled dinoflagellate cysts: Palynological tracers of sea-surface conditions in middle to high latitude marine environments. <i>Geobios</i> , 1997, 30, 905-920.	0.7	157
71	Plankton in the Norwegian-Greenland Sea: from living communities to sediment assemblages ?an actualistic approach. <i>Geologische Rundschau: Zeitschrift Fur Allgemeine Geologie</i> , 1995, 84, 108.	1.3	119
72	Distribution patterns of dinoflagellate cysts and other organic-walled microfossils in recent Norwegian-Greenland Sea sediments. <i>Marine Micropaleontology</i> , 1995, 24, 307-334.	0.5	129

#	ARTICLE	IF	CITATIONS
73	A forum on Neogene and quaternary dinoflagellate cysts: The edited transcript of a round table discussion held at the third workshop on Neogene and Quaternary dinoflagellates; with taxonomic appendix. <i>Palynology</i> , 1993, 17, 201-239.	0.7	45
74	Variations in surface water mass conditions in the Norwegian Sea: Evidence from Holocene coccolith and dinoflagellate cyst assemblages. <i>Marine Micropaleontology</i> , 1992, 20, 129-146.	0.5	61
75	Dinoflagellate Cyst Ecostratigraphy of Pliocene–Pleistocene Sediments from the Yermak Plateau (Arctic Ocean, Hole 911A). , 0, , .		13
76	<i>Batiacasphaera micropapillata</i> . , 0, , 301-314.		4