List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5728898/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Biological monitoring of exposure to benzene: a comparison between S-phenylmercapturic acid, trans,trans-muconic acid, and phenol Occupational and Environmental Medicine, 1995, 52, 611-620.	1.3	159
2	Guidelines for the derivation of Biomonitoring Equivalents: Report from the Biomonitoring Equivalents Expert Workshop. Regulatory Toxicology and Pharmacology, 2008, 51, S4-S15.	1.3	147
3	The use of biomonitoring data in exposure and human health risk assessment: benzene case study. Critical Reviews in Toxicology, 2013, 43, 119-153.	1.9	107
4	Guidelines for the communication of Biomonitoring Equivalents: Report from the Biomonitoring Equivalents Expert Workshop. Regulatory Toxicology and Pharmacology, 2008, 51, S16-S26.	1.3	99
5	Exposure to polycyclic aromatic hydrocarbons in petrochemical industries by measurement of urinary 1-hydroxypyrene Occupational and Environmental Medicine, 1994, 51, 250-258.	1.3	91
6	Biomarkers of Exposure to 1,3-Butadiene as a Basis for Cancer Risk Assessment. Toxicological Sciences, 2000, 56, 189-202.	1.4	80
7	The Role of Hydrolysis in the Detoxification of 1,2:3,4-Diepoxybutane by Human, Rat, and Mouse Liver and Lungin Vitro. Toxicology and Applied Pharmacology, 1996, 141, 617-627.	1.3	78
8	Creating context for the use of DNA adduct data in cancer risk assessment: II. Overview of methods of identification and quantitation of DNA damage. Critical Reviews in Toxicology, 2009, 39, 679-694.	1.9	75
9	Glutathione Conjugation of 1,2:3,4-Diepoxybutane in Human Liver and Rat and Mouse Liver and Lungin Vitro. Toxicology and Applied Pharmacology, 1996, 136, 307-316.	1.3	73
10	Biomonitoring of exposure to ethylene oxide and propylene oxide by determination of hemoglobin adducts: correlations between airborne exposure and adduct levels. International Archives of Occupational and Environmental Health, 1999, 72, 142-150.	1.1	69
11	A chemical–biological similarity-based grouping of complex substances as a prototype approach for evaluating chemical alternatives. Green Chemistry, 2016, 18, 4407-4419.	4.6	69
12	Biomarkers for assessing occupational exposures to 1,3-butadiene. Chemico-Biological Interactions, 2001, 135-136, 429-453.	1.7	64
13	Use of haemoglobin adducts in exposure monitoring and risk assessment. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2002, 778, 309-322.	1.2	59
14	Building scientific confidence in the development and evaluation of read-across. Regulatory Toxicology and Pharmacology, 2015, 72, 117-133.	1.3	56
15	Biological effect monitoring in industrial workers following incidental exposure to high concentrations of ethylene oxide. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 1995, 329, 63-77.	0.4	55
16	Toxicity of the cysteine-S-conjugates and mercapturic acids of four structurally related difluoroethylenes in isolated proximal tubular cells from rat kidney. Biochemical Pharmacology, 1989, 38, 3731-3741.	2.0	54
17	Urinary excretion of N -acetyl- S -allyl- L -cysteine upon garlic consumption by human volunteers. Archives of Toxicology, 1996, 70, 635-639.	1.9	54
18	The role of metallothionein in the reduction of cisplatin-induced nephrotoxicity by Bi3+-pretreatment in the rat in vivo and in vitro. Biochemical Pharmacology, 1991, 41, 369-375.	2.0	50

#	Article	IF	CITATIONS
19	Mutagenicity and cytotoxicity of two regioisomeric mercapturic acids and cysteine S-conjugates of trichloroethylene. Archives of Toxicology, 1991, 65, 373-380.	1.9	49
20	Human biomonitoring as a pragmatic tool to support health risk management of chemicals – Examples under the EU REACH programme. Regulatory Toxicology and Pharmacology, 2011, 59, 125-132.	1.3	49
21	Renal proximal tubular cells in suspension or in primary culture as in vitro models to study nephrotoxicity. Chemico-Biological Interactions, 1990, 76, 251-291.	1.7	48
22	Urinary 1-hydroxypyrene as biomarker of exposure to polycyclic aromatic hydrocarbons in workers in petrochemical industries: baseline values and dermal uptake. Science of the Total Environment, 1995, 163, 203-209.	3.9	48
23	Comparative assessment of gastrointestinal irritant potency in man of tin(II) chloride and tin migrated from packaging. Food and Chemical Toxicology, 2003, 41, 1663-1670.	1.8	47
24	Formation of DNA adducts and induction of mutagenic effects in rats following 4 weeks inhalation exposure to ethylene oxide as a basis for cancer risk assessment. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 2000, 447, 27-48.	0.4	45
25	Heavy hydrocarbon fate and transport in the environment. Quarterly Journal of Engineering Geology and Hydrogeology, 2017, 50, 333-346.	0.8	44
26	Proposal for the assessment of quantitative dermal exposure limits in occupational environments: Part 1. Development of a concept to derive a quantitative dermal occupational exposure limit. Occupational and Environmental Medicine, 1998, 55, 795-804.	1.3	43
27	Isolated proximal tubular cells from rat kidney as an in vitro model for studies on nephrotoxicity. Toxicology and Applied Pharmacology, 1989, 101, 135-143.	1.3	41
28	Quantitative and qualitative differences in the metabolism of 14C-1,3- butadiene in rats and mice: relevance to cancer susceptibility. Toxicological Sciences, 1999, 49, 186-201.	1.4	38
29	A systematic approach for evaluating and scoring human data. Regulatory Toxicology and Pharmacology, 2013, 66, 241-247.	1.3	36
30	Primary culture of proximal tubular cells from normal rat kidney as an in vitro model to study mechanisms of nephrotoxicity. Biochemical Pharmacology, 1990, 39, 1335-1345.	2.0	35
31	Dermal penetration and metabolism of five glycidyl ethers in human, rat and mouse skin. Xenobiotica, 2000, 30, 469-483.	0.5	35
32	Quantification of DNA Adducts Formed in Liver, Lungs, and Isolated Lung Cells of Rats and Mice Exposed to 14C-Styrene by Nose-Only Inhalation. Toxicological Sciences, 2000, 57, 203-216.	1.4	35
33	Application of the urinary S-phenylmercapturic acid test as a biomarker for low levels of exposure to benzene in industry Occupational and Environmental Medicine, 1993, 50, 460-469.	1.3	32
34	Urinary metabolites and haemoglobin adducts as biomarkers of exposure to 1,3-butadiene: a basis for 1,3-butadiene cancer risk assessment Chemico-Biological Interactions, 2001, 135-136, 695-701.	1.7	31
35	Identification and quantitative determination of mercapturic acids formed from Z- and E-1,3-dichloropropene by the rat, using gas chromatography with three different detection techniques. Archives of Toxicology, 1986, 59, 235-241.	1.9	30
36	Prenatal developmental toxicity testing of petroleum substances: Application of the mouse embryonic stem cell test (EST) to compare in vitro potencies with potencies observed in vivo. Toxicology in Vitro, 2017, 44, 303-312.	1.1	30

#	Article	IF	CITATIONS
37	4-methylthiobenzoic acid reduces cisplatin nephrotoxicity in rats without compromising anti-tumour activity. Biochemical Pharmacology, 1991, 41, 1997-2003.	2.0	29
38	Allylmercapturic acid as urinary biomarker of human exposure to allyl chloride Occupational and Environmental Medicine, 1997, 54, 653-661.	1.3	26
39	Comparison of the metabolism of ethylene glycol and glycolic acidin vitroby precision-cut tissue slices from female rat, rabbit and human liver. Xenobiotica, 2004, 34, 31-48.	0.5	26
40	An organizational approach for the assessment of DNA adduct data in risk assessment: case studies for aflatoxin B ₁ , tamoxifen and vinyl chloride. Critical Reviews in Toxicology, 2014, 44, 348-391.	1.9	26
41	Toxicogenomics in vitro as an alternative tool for safety evaluation of petroleum substances and PAHs with regard to prenatal developmental toxicity. Toxicology in Vitro, 2015, 29, 299-307.	1.1	26
42	The Role of Endocrine and Dioxin-Like Activity of Extracts of Petroleum Substances in Developmental Toxicity as Detected in a Panel of CALUX Reporter Gene Assays. Toxicological Sciences, 2018, 164, 576-591.	1.4	26
43	Identification of Novel Metabolites of Butadiene Monoepoxide in Rats and Mice. Chemical Research in Toxicology, 1998, 11, 1543-1555.	1.7	25
44	The selective determination of potentially carcinogenic polycyclic aromatic compounds in lubricant base oils by the DMSO extraction method IP346 and its correlation to mouse skin painting carcinogenicity assays. Regulatory Toxicology and Pharmacology, 2019, 106, 316-333.	1.3	25
45	In vitro metabolism of naphthalene and its alkylated congeners by human and rat liver microsomes via alkyl side chain or aromatic oxidation. Chemico-Biological Interactions, 2020, 315, 108905.	1.7	25
46	Isolated proximal tubular cells from rat kidney as an in vitro model for studies on nephrotoxicity. Toxicology and Applied Pharmacology, 1989, 101, 144-157.	1.3	24
47	Proposal for the assessment to quantitative dermal exposure limits in occupational environments: Part 2. Feasibility study for application in an exposure scenario for MDA by two different dermal exposure sampling methods. Occupational and Environmental Medicine, 1998, 55, 805-811.	1.3	24
48	Grouping of Petroleum Substances as Example UVCBs by Ion Mobility-Mass Spectrometry to Enable Chemical Composition-Based Read-Across. Environmental Science & Technology, 2017, 51, 7197-7207.	4.6	23
49	In vitro prenatal developmental toxicity induced by some petroleum substances is mediated by their 3- to 7-ring PAH constituent with a potential role for the aryl hydrocarbon receptor (AhR). Toxicology Letters, 2019, 315, 64-76.	0.4	23
50	Toxicological and ecotoxicological properties of gas-to-liquid (GTL) products. 1. Mammalian toxicology. Critical Reviews in Toxicology, 2017, 47, 121-144.	1.9	22
51	Regeneration experiments of the platinated enzyme fumarase, using sodium diethyldithiocarbamate, thiourea, and sodium thiosulfate. Journal of Inorganic Biochemistry, 1991, 41, 17-24.	1.5	20
52	Effects of Exposure to Elemental Mercury on the Nervous System and the Kidneys of Workers Producing Natural Gas. Archives of Environmental Health, 1996, 51, 108-115.	0.4	20
53	Urinary biomarkers in the risk assessment of PAHs. Occupational and Environmental Medicine, 2008, 65, 221-222.	1.3	20
54	Dermal uptake of petroleum substances. Toxicology Letters, 2015, 235, 123-139.	0.4	20

4

#	Article	IF	CITATIONS
55	Increased albumin excretion in industrial workers due to shift work rather than to prolonged exposure to low concentrations of chlorinated hydrocarbons Occupational and Environmental Medicine, 1994, 51, 638-641.	1.3	18
56	Metabolism of butadiene by mice, rats, and humans: a comparison of physiologically based toxicokinetic model predictions and experimental data. Toxicology, 1996, 113, 48-54.	2.0	17
57	DNA adducts in rats and mice following exposure to [4-14C]-1,2-epoxy-3-butene and to [2,3-14C]-1,3-butadiene. Chemico-Biological Interactions, 2004, 148, 69-92.	1.7	17
58	Metabolic inactivation of five glycidyl ethers in lung and liver of humans, rats and micein vitro. Xenobiotica, 2000, 30, 485-502.	0.5	16
59	Disposition of [Ring-U-14C]styrene in Rats and Mice Exposed by Recirculating Nose-Only Inhalation. Toxicological Sciences, 2000, 58, 161-172.	1.4	16
60	3-Chloro-2-hydroxypropylmercapturic acid and α-chlorohydrin as biomarkers of occupational exposure to epichlorohydrin. Environmental Toxicology and Pharmacology, 1997, 3, 175-185.	2.0	15
61	Comparative toxicokinetics of low-viscosity mineral oil in Fischer 344 rats, Sprague–Dawley rats, and humans – Implications for an Acceptable Daily Intake (ADI). Regulatory Toxicology and Pharmacology, 2012, 63, 69-77.	1.3	15
62	A proposed framework for the interpretation of biomonitoring data. Environmental Health, 2008, 7, S12.	1.7	14
63	Assessment of petroleum streams for thyroid toxicity. Toxicology Letters, 2016, 254, 52-62.	0.4	14
64	Grouping of UVCB substances with new approach methodologies (NAMs) data. ALTEX: Alternatives To Animal Experimentation, 2021, 38, 123-137.	0.9	13
65	Disposition of butadiene epoxides in Sprague-Dawley rats. Chemico-Biological Interactions, 1997, 104, 103-115.	1.7	12
66	Occupational exposure to cis-1,3-dichloropropene: biological effect monitoring of kidney and liver function. Occupational and Environmental Medicine, 2000, 57, 745-751.	1.3	12
67	A novel DNA adduct, originating from 1,2-epoxy-3,4-butanediol, is the major DNA adduct after exposure to [2,3-14C]-1,3-butadiene, but not after exposure to [4-14C]-1,2-epoxy-3-butene. Chemico-Biological Interactions, 2001, 135-136, 687-693.	1.7	12
68	Human biomonitoring activities – Programmes by industry. International Journal of Hygiene and Environmental Health, 2007, 210, 259-261.	2.1	11
69	The role of metabolism in the developmental toxicity of polycyclic aromatic hydrocarbonâ€containing extracts of petroleum substances. Journal of Applied Toxicology, 2020, 40, 330-341.	1.4	11
70	Nephrotoxicity of halogenated alkenyl cysteine-S-conjugates. Life Sciences, 1991, 49, 1769-1776.	2.0	10
71	Effects of exposure to low concentrations of chlorinated hydrocarbons on the kidney and liver of industrial workers Occupational and Environmental Medicine, 1993, 50, 331-339.	1.3	10
72	Personal air sampling and biological monitoring of occupational exposure to the soil fumigant cis-1,3-dichloropropene. Occupational and Environmental Medicine, 2000, 57, 738-744.	1.3	10

#	Article	IF	CITATIONS
73	Gas chromatography-electron capture determination of styrene-7,8-oxide enantiomers. Biomedical Applications, 2000, 749, 265-274.	1.7	9
74	Development of a competitive immunoassay for the determination of N-(2-hydroxypropyl)valine adducts in human haemoglobin and its application in biological monitoring. Biomarkers, 2005, 10, 127-137.	0.9	9
75	Application of human biomonitoring (HBM) of chemical exposure in the characterisation of health risks under REACH. International Journal of Hygiene and Environmental Health, 2012, 215, 238-241.	2.1	9
76	Developmental toxicity testing of unsubstituted and methylated 4- and 5-ring polycyclic aromatic hydrocarbons using the zebrafish embryotoxicity test. Toxicology in Vitro, 2022, 80, 105312.	1.1	9
77	Hepatic and pulmonary glutathione conjugation of 1,2:3,4-diepoxybutane in human, rat, and mouse in vitro. Toxicology, 1996, 113, 297-299.	2.0	8
78	Metabolic inactivation of 2-oxiranylmethyl 2-ethyl2,5-dimethylhexanoate (C10GE) in skin, lung and liver of human, rat and mouse. Xenobiotica, 1999, 29, 987-1006.	0.5	8
79	Guidance on the selection of cohorts for the extended one-generation reproduction toxicity study (OECD test guideline 443). Regulatory Toxicology and Pharmacology, 2016, 80, 32-40.	1.3	8
80	Toxicological and ecotoxicological properties of gas-to-liquid (GTL) products. 2. Ecotoxicology. Critical Reviews in Toxicology, 2018, 48, 273-296.	1.9	8
81	Derivation of an occupational exposure limit for benzene using epidemiological study quality assessment tools. Toxicology Letters, 2020, 334, 117-144.	0.4	8
82	Prenatal developmental toxicity testing of petroleum substances using the zebrafish embryotoxicity test. ALTEX: Alternatives To Animal Experimentation, 2019, 36, 245-260.	0.9	8
83	The effect of alkyl substitution on the oxidative metabolism and mutagenicity of phenanthrene. Archives of Toxicology, 2022, 96, 1109-1131.	1.9	8
84	Development of a competitive immunoassay for the determination of N-(2-hydroxyethyl)valine adducts in human haemoglobin and its application in biological monitoring. Biomarkers, 2004, 9, 407-417.	0.9	7
85	Determination of Exposure to Bitumen and Fume from Bitumen in the Oil Industry Through Determination of Urinary 1-Hydroxypyrene. Journal of Occupational and Environmental Hygiene, 2007, 4, 111-117.	0.4	7
86	Guidance on classification for reproductive toxicity under the globally harmonized system of classification and labelling of chemicals (GHS). Critical Reviews in Toxicology, 2013, 43, 850-891.	1.9	7
87	Biological monitoring to assess dermal exposure to ethylene oxide vapours during an incidental release. Toxicology Letters, 2014, 231, 387-390.	0.4	7
88	Risk assessment of workers exposed to crystalline silica aerosols. Human and Ecological Risk Assessment (HERA), 2016, 22, 1678-1686.	1.7	7
89	Non-parametric estimation of low-concentration benzene metabolism. Chemico-Biological Interactions, 2017, 278, 242-255.	1.7	7
90	Predicting the in vivo developmental toxicity of benzo[a]pyrene (BaP) in rats by an in vitro–in silico approach. Archives of Toxicology, 2021, 95, 3323-3340.	1.9	7

#	Article	IF	CITATIONS
91	Synthesis of [14C]-labelled glycidyl and glycerol ethers of aliphatic and aromatic alcohols. Journal of Labelled Compounds and Radiopharmaceuticals, 2000, 43, 147-167.	0.5	5
92	Biomonitoring as a tool in the human health risk characterization of dermal exposure. Human and Experimental Toxicology, 2008, 27, 297-305.	1.1	5
93	The influence of alkyl substitution on the in vitro metabolism and mutagenicity of benzo[a]pyrene. Chemico-Biological Interactions, 2022, 363, 110007.	1.7	5
94	Ceramic Fibers and Other Respiratory Hazards During the Renewal of the Refractory Lining in a Large Industrial Furnace. Journal of Occupational and Environmental Hygiene, 1994, 9, 32-35.	0.5	4
95	Reproductive and developmental toxicity assessment of gas-to-liquid diesel and base oil. Reproductive Toxicology, 2015, 56, 30.	1.3	4
96	The low-dose benzene debate needs a sharp blade. Chemico-Biological Interactions, 2017, 278, 239-241.	1.7	4
97	Cisplatin Nephrotoxicity and Platinum-Metallothioneins: Uptake and Toxicity in Proximal Tubular Cells from Rat Kidney. Contributions To Nephrology, 1990, 83, 208-212.	1.1	4
98	A consistent and transparent approach for calculation of Derived No-Effect Levels (DNELs) for petroleum substances. Regulatory Toxicology and Pharmacology, 2012, 62, 85-98.	1.3	3
99	Prenatal developmental toxicity studies on fumes from oxidised asphalt (OA) in the rat. Reproductive Toxicology, 2021, 102, 67-79.	1.3	3
100	Developmental toxicity testing of the fume condensate extracts of bitumen and oxidized asphalt in a series of in vitro alternative assays. Toxicology in Vitro, 2021, 75, 105195.	1.1	3
101	Skin irritation by kerosine. Toxicology Letters, 2006, 164, S94.	0.4	2
102	Getting under the skin. Human and Experimental Toxicology, 2008, 27, 267-268.	1.1	2
103	Interpretation of Human Biological Monitoring Data Using a Newly Developed Generic Physiological-Based Toxicokinetic Model. , 2013, , 137-150.		2
104	Prenatal developmental toxicity studies on fumes from bitumen in the rat. Reproductive Toxicology, 2021, 99, 15-26.	1.3	2
105	Molecular Dosimetry of 2,4-Difluoroaniline in Humans and Rats by Determination of Hemoglobin Adducts. Environmental Health Perspectives, 1994, 102, 27.	2.8	1
106	Assessment of Asphalt Workers' Dermal Exposure. Annals of Occupational Hygiene, 2005, 49, 93; author reply 93-4.	1.9	0
107	Biomonitoring as a tool in the human health risk assessment of dermal exposure. Toxicology Letters, 2006, 164, S323.	0.4	0