Tadeusz S Porowski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5740818/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Urinary citrate excretion in healthy children depends on age and gender. Pediatric Nephrology, 2014, 29, 1575-1582.	0.9	38
2	An association between kidney stone composition and urinary metabolic disturbances in children. Journal of Pediatric Urology, 2014, 10, 130-135.	0.6	38
3	A potential pathogenic role of oxalate in autism. European Journal of Paediatric Neurology, 2012, 16, 485-491.	0.7	34
4	Assessment of serum cystatinÂC in children with congenital solitary kidney. Pediatric Nephrology, 2006, 21, 688-693.	0.9	25
5	A new approach to the diagnosis of children's urolithiasis based on the Bonn Risk Index. Pediatric Nephrology, 2008, 23, 1123-1128.	0.9	21
6	Does obesity or hyperuricemia influence lithogenic risk profile in children with urolithiasis?. Pediatric Nephrology, 2015, 30, 797-803.	0.9	20
7	Normative data on the Bonn Risk Index for calcium oxalate crystallization in healthy children. Pediatric Nephrology, 2007, 22, 514-520.	0.9	14
8	Reference values of plasma oxalate in children and adolescents. Pediatric Nephrology, 2008, 23, 1787-1794.	0.9	14
9	Correspondence between Ca2+ and calciuria, citrate level and pH of urine in pediatric urolithiasis. Pediatric Nephrology, 2013, 28, 1079-1084.	0.9	14
10	Upper metastable limit osmolality of urine as a predictor of kidney stone formation in children. Urolithiasis, 2019, 47, 155-163.	1.2	14
11	Laminin and transforming growth factor beta-1 in children with vesicoureteric reflux. Pediatric Nephrology, 2008, 23, 769-774.	0.9	12
12	Urinary nerve growth factor level in children with neurogenic bladder due to myelomeningocele. Scandinavian Journal of Urology, 2013, 47, 411-417.	0.6	10
13	Urodynamic Findings and Renal Function in Children with Neurogenic Bladder after Myelomeningocele. Urologia Internationalis, 2015, 95, 146-152.	0.6	10
14	Isolation, purification, and characterization of glucosamine-6-phosphate-N-acetylase from pig liver. Biochemical Medicine and Metabolic Biology, 1990, 44, 1-12.	0.7	9
15	Determining normal values of urinary phosphorus excretion in 3913 healthy children aged 2–18 to aid early diagnosis and treatment for urolithiasis. Acta Paediatrica, International Journal of Paediatrics, 2017, 106, 1170-1175.	0.7	9
16	Induced urinary crystal formation as an analytical strategy for the prediction and monitoring of urolithiasis and other metabolism-related disorders. EPMA Journal, 2014, 5, 13.	3.3	8
17	Assessment of Lithogenic Risk in Children Based on a Morning Spot Urine Sample. Journal of Urology, 2010, 184, 2103-2108.	0.2	7
18	A semi-micromethod for determination of oxalate in human plasma. Acta Poloniae Pharmaceutica, 2003, 60, 239-45.	0.3	4

TADEUSZ S POROWSKI

#	Article	IF	CITATIONS
19	Spontaneous urinary calcium oxalate crystallization in hypercalciuric children. Pediatric Nephrology, 2009, 24, 1705-1710.	0.9	3
20	Clinical profile of a Polish cohort of children and young adults with cystinuria. Renal Failure, 2021, 43, 62-70.	0.8	2
21	Serum and urine fibronectin levels in children with vesicoureteral reflux. Pediatric Nephrology, 2007, 22, 1173-1179.	0.9	1
22	A Need to Establish Normative Data for Plasma Oxalates. American Journal of Kidney Diseases, 2008, 51, 1071-1072.	2.1	1
23	Citrate usage in the leading causes of blindness: new possibilities for the old metabolite. Metabolomics, 2018, 14, 82.	1.4	1
24	The role of selected parental and perinatal factors in the future mobility of children with myelomeningocele. Family Medicine and Primary Care Review, 2016, 1, 24-28.	0.1	0
25	Urinary risk factors for calcium oxalate urolithiasis in children with monosymptomatic enuresis. Journal of Nephropathology, 2021, 10, e42-e42.	0.1	Ο