

Ji-Liang Li

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/574378/ji-liang-li-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62

papers

5,368

citations

35

h-index

64

g-index

64

ext. papers

6,004

ext. citations

6.9

avg, IF

4.99

L-index

#	Paper	IF	Citations
62	Downregulation of Manic fringe impedes angiogenesis and cell migration of renal carcinoma.. <i>Microvascular Research</i> , 2022 , 104341	3.7	1
61	Role of lncSLCO1C1 in gastric cancer progression and resistance to oxaliplatin therapy.. <i>Clinical and Translational Medicine</i> , 2022 , 12, e691	5.7	0
60	Tspan5 promotes epithelial-mesenchymal transition and tumour metastasis of hepatocellular carcinoma by activating Notch signalling. <i>Molecular Oncology</i> , 2021 , 15, 3184-3202	7.9	4
59	Nuclear and stromal expression of Manic fringe in renal cell carcinoma. <i>Experimental and Molecular Pathology</i> , 2021 , 122, 104667	4.4	2
58	RHOQ is induced by DLL4 and regulates angiogenesis by determining the intracellular route of the Notch intracellular domain. <i>Angiogenesis</i> , 2020 , 23, 493-513	10.6	16
57	Development of Therapeutic Anti-JAGGED1 Antibodies for Cancer Therapy. <i>Molecular Cancer Therapeutics</i> , 2019 , 18, 2030-2042	6.1	16
56	RN181 is a tumour suppressor in gastric cancer by regulation of the ERK/MAPK-cyclin D1/CDK4 pathway. <i>Journal of Pathology</i> , 2019 , 248, 204-216	9.4	11
55	contributes to epithelial-mesenchymal transition (EMT) by activating AKT signaling pathway and regulating MMP-2 expression. <i>Cancer Management and Research</i> , 2019 , 11, 2415-2424	3.6	13
54	M2 macrophages mediate sorafenib resistance by secreting HGF in a feed-forward manner in hepatocellular carcinoma. <i>British Journal of Cancer</i> , 2019 , 121, 22-33	8.7	46
53	Disordered intestinal microbes are associated with the activity of Systemic Lupus Erythematosus. <i>Clinical Science</i> , 2019 , 133, 821-838	6.5	57
52	WAP four-disulfide core domain protein 2 promotes metastasis of human ovarian cancer by regulation of metastasis-associated genes. <i>Journal of Ovarian Research</i> , 2017 , 10, 40	5.5	9
51	Role of Delta-like 4 in Jagged1-induced tumour angiogenesis and tumour growth. <i>Oncotarget</i> , 2017 , 8, 40115-40131	3.3	24
50	hTERT mediates gastric cancer metastasis partially through the indirect targeting of ITGB1 by microRNA-29a. <i>Scientific Reports</i> , 2016 , 6, 21955	4.9	38
49	Tspan5 is an independent favourable prognostic factor and suppresses tumour growth in gastric cancer. <i>Oncotarget</i> , 2016 , 7, 40160-40173	3.3	15
48	Disrupting Hypoxia-Induced Bicarbonate Transport Acidifies Tumor Cells and Suppresses Tumor Growth. <i>Cancer Research</i> , 2016 , 76, 3744-55	10.1	63
47	WAP four-disulfide core domain protein 2 gene(WFDC2) is a target of estrogen in ovarian cancer cells. <i>Journal of Ovarian Research</i> , 2016 , 9, 10	5.5	7
46	Epstein-Barr virus-encoded microRNA BART1 induces tumour metastasis by regulating PTEN-dependent pathways in nasopharyngeal carcinoma. <i>Nature Communications</i> , 2015 , 6, 7353	17.4	150

45	Meta-analysis of Androgen Insensitivity in Preoperative Hormone Therapy in Hypospadias. <i>Urology</i> , 2015 , 85, 1166-1172	1.6	5
44	Carbonic anhydrase IX induction defines a heterogeneous cancer cell response to hypoxia and mediates stem cell-like properties and sensitivity to HDAC inhibition. <i>Oncotarget</i> , 2015 , 6, 19413-27	3.3	35
43	Estrogen receptor- α directly regulates the hypoxia-inducible factor 1 pathway associated with antiestrogen response in breast cancer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 15172-7	11.5	74
42	Combining lapatinib and pertuzumab to overcome lapatinib resistance due to NRG1-mediated signalling in HER2-amplified breast cancer. <i>Oncotarget</i> , 2015 , 6, 5678-94	3.3	25
41	IGF-1R inhibition induces schedule-dependent sensitization of human melanoma to temozolomide. <i>Oncotarget</i> , 2015 , 6, 39877-90	3.3	17
40	Nuclear HER4 mediates acquired resistance to trastuzumab and is associated with poor outcome in HER2 positive breast cancer. <i>Oncotarget</i> , 2014 , 5, 5934-49	3.3	48
39	ADAM10 mediates trastuzumab resistance and is correlated with survival in HER2 positive breast cancer. <i>Oncotarget</i> , 2014 , 5, 6633-46	3.3	54
38	Fatty acid uptake and lipid storage induced by HIF-1 α contribute to cell growth and survival after hypoxia-reoxygenation. <i>Cell Reports</i> , 2014 , 9, 349-365	10.6	324
37	Functional comparison of Notch ligands in tumour angiogenesis. <i>Asian Pacific Journal of Tropical Disease</i> , 2014 , 4, 229		
36	Dichloroacetate reverses the hypoxic adaptation to bevacizumab and enhances its antitumor effects in mouse xenografts. <i>Journal of Molecular Medicine</i> , 2013 , 91, 749-58	5.5	55
35	A core human primary tumor angiogenesis signature identifies the endothelial orphan receptor ELTD1 as a key regulator of angiogenesis. <i>Cancer Cell</i> , 2013 , 24, 229-41	24.3	164
34	Neratinib overcomes trastuzumab resistance in HER2 amplified breast cancer. <i>Oncotarget</i> , 2013 , 4, 15923-305	3.3	108
33	Glucose utilization via glycogen phosphorylase sustains proliferation and prevents premature senescence in cancer cells. <i>Cell Metabolism</i> , 2012 , 16, 751-64	24.6	241
32	Carbonic anhydrase IX promotes tumor growth and necrosis in vivo and inhibition enhances anti-VEGF therapy. <i>Clinical Cancer Research</i> , 2012 , 18, 3100-11	12.9	195
31	RN181 suppresses hepatocellular carcinoma growth by inhibition of the ERK/MAPK pathway. <i>Hepatology</i> , 2011 , 53, 1932-42	11.2	33
30	DLL4-Notch signaling mediates tumor resistance to anti-VEGF therapy in vivo. <i>Cancer Research</i> , 2011 , 71, 6073-83	10.1	181
29	MicroRNA-210 regulates mitochondrial free radical response to hypoxia and krebs cycle in cancer cells by targeting iron sulfur cluster protein ISCU. <i>PLoS ONE</i> , 2010 , 5, e10345	3.7	243
28	Effects of acute versus chronic hypoxia on DNA damage responses and genomic instability. <i>Cancer Research</i> , 2010 , 70, 925-35	10.1	142

27	Targeting DLL4 in tumors shows preclinical activity but potentially significant toxicity. <i>Future Oncology</i> , 2010 , 6, 1099-103	3.6	16
26	Expression of vascular notch ligand delta-like 4 and inflammatory markers in breast cancer. <i>American Journal of Pathology</i> , 2010 , 176, 2019-28	5.8	89
25	Crosstalk of VEGF and Notch pathways in tumour angiogenesis: therapeutic implications. <i>Frontiers in Bioscience - Landmark</i> , 2009 , 14, 3094-110	2.8	103
24	Conformation-dependent single-chain variable fragment antibodies specifically recognize beta-amyloid oligomers. <i>FEBS Letters</i> , 2009 , 583, 579-84	3.8	50
23	Nuclear and membrane expression of the angiogenesis regulator delta-like ligand 4 (DLL4) in normal and malignant human tissues. <i>Histopathology</i> , 2009 , 54, 598-606	7.3	14
22	Role of hypoxia-inducible factors in epigenetic regulation via histone demethylases. <i>Annals of the New York Academy of Sciences</i> , 2009 , 1177, 185-97	6.5	86
21	Regulation of multiple angiogenic pathways by Dll4 and Notch in human umbilical vein endothelial cells. <i>Microvascular Research</i> , 2008 , 75, 144-54	3.7	169
20	The potential of new tumor endothelium-specific markers for the development of antivascular therapy. <i>Cancer Cell</i> , 2007 , 11, 478-81	24.3	19
19	Delta-like 4 Notch ligand regulates tumor angiogenesis, improves tumor vascular function, and promotes tumor growth in vivo. <i>Cancer Research</i> , 2007 , 67, 11244-53	10.1	265
18	Up-regulation of endothelial delta-like 4 expression correlates with vessel maturation in bladder cancer. <i>Clinical Cancer Research</i> , 2006 , 12, 4836-44	12.9	112
17	Up-regulation of the Notch ligand Delta-like 4 inhibits VEGF-induced endothelial cell function. <i>Blood</i> , 2006 , 107, 931-9	2.2	295
16	Contrasting properties of hypoxia-inducible factor 1 (HIF-1) and HIF-2 in von Hippel-Lindau-associated renal cell carcinoma. <i>Molecular and Cellular Biology</i> , 2005 , 25, 5675-86	4.8	754
15	Notch signaling from tumor cells: a new mechanism of angiogenesis. <i>Cancer Cell</i> , 2005 , 8, 1-3	24.3	95
14	Up-regulation of delta-like 4 ligand in human tumor vasculature and the role of basal expression in endothelial cell function. <i>Cancer Research</i> , 2005 , 65, 8690-7	10.1	288
13	Physical and functional interaction between the Bloom syndrome gene product and the largest subunit of chromatin assembly factor 1. <i>Molecular and Cellular Biology</i> , 2004 , 24, 4710-9	4.8	42
12	Characterisation of a sexual stage-specific gene encoding ORC1 homologue in the human malaria parasite <i>Plasmodium falciparum</i> . <i>Parasitology International</i> , 2003 , 52, 41-52	2.1	11
11	Functional interaction between the Bloom syndrome helicase and the RAD51 paralog, RAD51L3 (RAD51D). <i>Journal of Biological Chemistry</i> , 2003 , 278, 48357-66	5.4	66
10	Identification of a second proliferating cell nuclear antigen in the human malarial pathogen <i>Plasmodium falciparum</i> . <i>International Journal for Parasitology</i> , 2002 , 32, 1683-92	4.3	8

9	Primary structure and sexual stage-specific expression of a LAMMER protein kinase of <i>Plasmodium falciparum</i> . <i>International Journal for Parasitology</i> , 2001 , 31, 387-92	4-3	18
8	Identification of an MCM4 homologue expressed specifically in the sexual stage of <i>Plasmodium falciparum</i> . <i>International Journal for Parasitology</i> , 2001 , 31, 1246-52	4-3	11
7	Molecular cloning of a gene encoding a 20S proteasome beta subunit from <i>Plasmodium falciparum</i> . <i>International Journal for Parasitology</i> , 2000 , 30, 729-33	4-3	7
6	Replication protein A physically interacts with the Bloom's syndrome protein and stimulates its helicase activity. <i>Journal of Biological Chemistry</i> , 2000 , 275, 23500-8	5-4	239
5	Guanylyl cyclase activity associated with putative bifunctional integral membrane proteins in <i>Plasmodium falciparum</i> . <i>Journal of Biological Chemistry</i> , 2000 , 275, 22147-56	5-4	65
4	A putative protein serine/threonine phosphatase from <i>Plasmodium falciparum</i> contains a large N-terminal extension and five unique inserts in the catalytic domain. <i>Molecular and Biochemical Parasitology</i> , 1998 , 95, 287-95	1-9	38
3	Protein phosphatase beta, a putative type-2A protein phosphatase from the human malaria parasite <i>Plasmodium falciparum</i> . <i>FEBS Journal</i> , 1997 , 249, 98-106		35
2	Pfmrk, a MO15-related protein kinase from <i>Plasmodium falciparum</i> . Gene cloning, sequence, stage-specific expression and chromosome localization. <i>FEBS Journal</i> , 1996 , 241, 805-13		53
1	Immunization strategies for the production of rat monoclonal anti-idiotope antibodies. <i>Journal of Immunological Methods</i> , 1991 , 142, 15-20	2-5	3