

Toshihiro Tanaka

List of Publications by Citations

Source: <https://exaly.com/author-pdf/575272/toshihiro-tanaka-publications-by-citations.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

165
papers

32,385
citations

66
h-index

174
g-index

174
ext. papers

36,333
ext. citations

10.9
avg, IF

7.43
L-index

#	Paper	IF	Citations
165	The International HapMap Project. <i>Nature</i> , 2003 , 426, 789-96	50.4	5039
164	A haplotype map of the human genome. <i>Nature</i> , 2005 , 437, 1299-320	50.4	4818
163	A second generation human haplotype map of over 3.1 million SNPs. <i>Nature</i> , 2007 , 449, 851-61	50.4	3647
162	Genetic studies of body mass index yield new insights for obesity biology. <i>Nature</i> , 2015 , 518, 197-206	50.4	2687
161	Genome-wide detection and characterization of positive selection in human populations. <i>Nature</i> , 2007 , 449, 913-8	50.4	1367
160	p53AIP1, a potential mediator of p53-dependent apoptosis, and its regulation by Ser-46-phosphorylated p53. <i>Cell</i> , 2000 , 102, 849-62	56.2	1005
159	Functional SNPs in the lymphotoxin-alpha gene that are associated with susceptibility to myocardial infarction. <i>Nature Genetics</i> , 2002 , 32, 650-4	36.3	755
158	Complete sequencing and characterization of 21,243 full-length human cDNAs. <i>Nature Genetics</i> , 2004 , 36, 40-5	36.3	695
157	GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. <i>Science</i> , 2013 , 340, 1467-71	33.3	563
156	Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. <i>Nature Genetics</i> , 2013 , 45, 145-54	36.3	505
155	Identification of a novel non-coding RNA, MIAT, that confers risk of myocardial infarction. <i>Journal of Human Genetics</i> , 2006 , 51, 1087-1099	4.3	491
154	Meta-analysis identifies six new susceptibility loci for atrial fibrillation. <i>Nature Genetics</i> , 2012 , 44, 670-5	36.3	429
153	A high-throughput SNP typing system for genome-wide association studies. <i>Journal of Human Genetics</i> , 2001 , 46, 471-7	4.3	405
152	Hyperglycemia causes oxidative stress in pancreatic beta-cells of GK rats, a model of type 2 diabetes. <i>Diabetes</i> , 1999 , 48, 927-32	0.9	376
151	A functional polymorphism in the 5UTR of GDF5 is associated with susceptibility to osteoarthritis. <i>Nature Genetics</i> , 2007 , 39, 529-33	36.3	370
150	Meta-analysis identifies common variants associated with body mass index in east Asians. <i>Nature Genetics</i> , 2012 , 44, 307-11	36.3	301
149	p53DINP1, a p53-inducible gene, regulates p53-dependent apoptosis. <i>Molecular Cell</i> , 2001 , 8, 85-94	17.6	283

148	Multi-ethnic genome-wide association study for atrial fibrillation. <i>Nature Genetics</i> , 2018 , 50, 1225-1233	36.3	277
147	Gene-based SNP discovery as part of the Japanese Millennium Genome Project: identification of 190,562 genetic variations in the human genome. Single-nucleotide polymorphism. <i>Journal of Human Genetics</i> , 2002 , 47, 605-10	4.3	253
146	Linkage of familial moyamoya disease (spontaneous occlusion of the circle of Willis) to chromosome 17q25. <i>Stroke</i> , 2000 , 31, 930-5	6.7	230
145	Large-scale genome-wide association studies in East Asians identify new genetic loci influencing metabolic traits. <i>Nature Genetics</i> , 2011 , 43, 990-5	36.3	229
144	A genome-wide association study identifies three new risk loci for Kawasaki disease. <i>Nature Genetics</i> , 2012 , 44, 517-21	36.3	217
143	JSNP: a database of common gene variations in the Japanese population. <i>Nucleic Acids Research</i> , 2002 , 30, 158-62	20.1	211
142	Functional variation in LGALS2 confers risk of myocardial infarction and regulates lymphotoxin-alpha secretion in vitro. <i>Nature</i> , 2004 , 429, 72-5	50.4	206
141	Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. <i>Nature Genetics</i> , 2012 , 44, 904-9	36.3	201
140	Common variants at CDKAL1 and KLF9 are associated with body mass index in east Asian populations. <i>Nature Genetics</i> , 2012 , 44, 302-6	36.3	192
139	Identification by cDNA microarray of genes involved in ovarian carcinogenesis. <i>Cancer Research</i> , 2000 , 60, 5007-11	10.1	187
138	Identification and characterization of the potential promoter regions of 1031 kinds of human genes. <i>Genome Research</i> , 2001 , 11, 677-84	9.7	184
137	Large-scale analyses of common and rare variants identify 12 new loci associated with atrial fibrillation. <i>Nature Genetics</i> , 2017 , 49, 946-952	36.3	176
136	Genome-wide meta-analysis identifies six novel loci associated with habitual coffee consumption. <i>Molecular Psychiatry</i> , 2015 , 20, 647-656	15.1	167
135	Alterations of gene expression during colorectal carcinogenesis revealed by cDNA microarrays after laser-capture microdissection of tumor tissues and normal epithelia. <i>Cancer Research</i> , 2001 , 61, 3544-9	10.1	160
134	Integrating genetic, transcriptional, and functional analyses to identify 5 novel genes for atrial fibrillation. <i>Circulation</i> , 2014 , 130, 1225-35	16.7	143
133	Meta-analysis of genome-wide association studies in East Asian-ancestry populations identifies four new loci for body mass index. <i>Human Molecular Genetics</i> , 2014 , 23, 5492-504	5.6	141
132	Diverse transcriptional initiation revealed by fine, large-scale mapping of mRNA start sites. <i>EMBO Reports</i> , 2001 , 2, 388-93	6.5	141
131	The p53 family member genes are involved in the Notch signal pathway. <i>Journal of Biological Chemistry</i> , 2002 , 277, 719-24	5.4	137

130	Identification of the gene responsible for gelatinous drop-like corneal dystrophy. <i>Nature Genetics</i> , 1999 , 21, 420-3	36.3	137
129	Homocysteine and coronary heart disease: meta-analysis of MTHFR case-control studies, avoiding publication bias. <i>PLoS Medicine</i> , 2012 , 9, e1001177	11.6	135
128	Novel calmodulin mutations associated with congenital arrhythmia susceptibility. <i>Circulation: Cardiovascular Genetics</i> , 2014 , 7, 466-74		133
127	Osteopenia and male-specific sudden cardiac death in mice lacking a zinc transporter gene, Znt5. <i>Human Molecular Genetics</i> , 2002 , 11, 1775-84	5.6	120
126	Mutations in the NMMHC-A gene cause autosomal dominant macrothrombocytopenia with leukocyte inclusions (May-Hegglin anomaly/Sebastian syndrome). <i>Blood</i> , 2001 , 97, 1147-9	2.2	118
125	Common variants in DVWA on chromosome 3p24.3 are associated with susceptibility to knee osteoarthritis. <i>Nature Genetics</i> , 2008 , 40, 994-8	36.3	116
124	Common variants in CASP3 confer susceptibility to Kawasaki disease. <i>Human Molecular Genetics</i> , 2010 , 19, 2898-906	5.6	114
123	Small intestinal stem cell identity is maintained with functional Paneth cells in heterotopically grafted epithelium onto the colon. <i>Genes and Development</i> , 2014 , 28, 1752-7	12.6	113
122	Prediction of sensitivity of esophageal tumors to adjuvant chemotherapy by cDNA microarray analysis of gene-expression profiles. <i>Cancer Research</i> , 2001 , 61, 6474-9	10.1	109
121	Novel genetic markers associate with atrial fibrillation risk in Europeans and Japanese. <i>Journal of the American College of Cardiology</i> , 2014 , 63, 1200-1210	15.1	102
120	Multiple loci are associated with white blood cell phenotypes. <i>PLoS Genetics</i> , 2011 , 7, e1002113	6	92
119	The Id2 gene is a novel target of transcriptional activation by EWS-ETS fusion proteins in Ewing family tumors. <i>Oncogene</i> , 2002 , 21, 8302-9	9.2	92
118	A functional SNP in PSMA6 confers risk of myocardial infarction in the Japanese population. <i>Nature Genetics</i> , 2006 , 38, 921-5	36.3	91
117	Association analyses of East Asian individuals and trans-ancestry analyses with European individuals reveal new loci associated with cholesterol and triglyceride levels. <i>Human Molecular Genetics</i> , 2017 , 26, 1770-1784	5.6	90
116	Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. <i>Nature Genetics</i> , 2017 , 49, 953-958	36.3	89
115	Genotype-Phenotype Correlation of Mutation for the Clinical and Electrocardiographic Characteristics of ProbandsWith Brugada Syndrome: A Japanese Multicenter Registry. <i>Circulation</i> , 2017 , 135, 2255-2270	16.7	88
114	GWAS for executive function and processing speed suggests involvement of the CADM2 gene. <i>Molecular Psychiatry</i> , 2016 , 21, 189-197	15.1	85
113	SNPs in BRAP associated with risk of myocardial infarction in Asian populations. <i>Nature Genetics</i> , 2009 , 41, 329-33	36.3	83

112	A functional polymorphism in THBS2 that affects alternative splicing and MMP binding is associated with lumbar-disc herniation. <i>American Journal of Human Genetics</i> , 2008 , 82, 1122-9	11	82
111	Novel mechanism of HERG current suppression in LQT2: shift in voltage dependence of HERG inactivation. <i>Circulation Research</i> , 1998 , 83, 415-22	15.7	82
110	Identification of 187 single nucleotide polymorphisms (SNPs) among 41 candidate genes for ischemic heart disease in the Japanese population. <i>Human Genetics</i> , 2000 , 106, 288-292	6.3	81
109	Expression of the gene for a membrane-bound fatty acid receptor in the pancreas and islet cell tumours in humans: evidence for GPR40 expression in pancreatic beta cells and implications for insulin secretion. <i>Diabetologia</i> , 2006 , 49, 962-8	10.3	77
108	Four novel KVLQT1 and four novel HERG mutations in familial long-QT syndrome. <i>Circulation</i> , 1997 , 95, 565-7	16.7	76
107	Genomic organization and mutational analysis of HERG, a gene responsible for familial long QT syndrome. <i>Human Genetics</i> , 1998 , 102, 435-9	6.3	75
106	Association between single-nucleotide polymorphisms in selectin genes and immunoglobulin A nephropathy. <i>American Journal of Human Genetics</i> , 2002 , 70, 781-6	11	73
105	Association between a single-nucleotide polymorphism in the promoter of the human interleukin-3 gene and rheumatoid arthritis in Japanese patients, and maximum-likelihood estimation of combinatorial effect that two genetic loci have on susceptibility to the disease. <i>American Journal of Human Genetics</i> , 2001 , 68, 671-85	11	72
104	Meta-analysis of genome-wide association studies of adult height in East Asians identifies 17 novel loci. <i>Human Molecular Genetics</i> , 2015 , 24, 1791-800	5.6	71
103	Up-regulation of the ectodermal-neural cortex 1 (ENC1) gene, a downstream target of the beta-catenin/T-cell factor complex, in colorectal carcinomas. <i>Cancer Research</i> , 2001 , 61, 7722-6	10.1	71
102	Isolation of a novel human gene, MARKL1, homologous to MARK3 and its involvement in hepatocellular carcinogenesis. <i>Neoplasia</i> , 2001 , 3, 4-9	6.4	69
101	ITPKC and CASP3 polymorphisms and risks for IVIG unresponsiveness and coronary artery lesion formation in Kawasaki disease. <i>Pharmacogenomics Journal</i> , 2013 , 13, 52-9	3.5	68
100	Genome-wide association study for C-reactive protein levels identified pleiotropic associations in the IL6 locus. <i>Human Molecular Genetics</i> , 2011 , 20, 1224-31	5.6	68
99	Identification of nine novel loci associated with white blood cell subtypes in a Japanese population. <i>PLoS Genetics</i> , 2011 , 7, e1002067	6	61
98	Growth and gene expression profile analyses of endometrial cancer cells expressing exogenous PTEN. <i>Cancer Research</i> , 2001 , 61, 3741-9	10.1	61
97	Unique activation status of peripheral blood mononuclear cells at acute phase of Kawasaki disease. <i>Clinical and Experimental Immunology</i> , 2010 , 160, 246-55	6.2	59
96	Identification of 187 single nucleotide polymorphisms (SNPs) among 41 candidate genes for ischemic heart disease in the Japanese population. <i>Human Genetics</i> , 2000 , 106, 288-92	6.3	59
95	Discovery and fine mapping of serum protein loci through transethnic meta-analysis. <i>American Journal of Human Genetics</i> , 2012 , 91, 744-53	11	58

94	Twenty single nucleotide polymorphisms (SNPs) and their allelic frequencies in four genes that are responsible for familial long QT syndrome in the Japanese population. <i>Journal of Human Genetics</i> , 2000 , 45, 182-3	4.3	56
93	Identification of AXUD1, a novel human gene induced by AXIN1 and its reduced expression in human carcinomas of the lung, liver, colon and kidney. <i>Oncogene</i> , 2001 , 20, 5062-6	9.2	52
92	Association of single-nucleotide polymorphisms in the polymeric immunoglobulin receptor gene with immunoglobulin A nephropathy (IgAN) in Japanese patients. <i>Journal of Human Genetics</i> , 2003 , 48, 293-299	4.3	51
91	Empirical estimation of genome-wide significance thresholds based on the 1000 Genomes Project data set. <i>Journal of Human Genetics</i> , 2016 , 61, 861-866	4.3	49
90	Genome wide analysis of drug-induced torsades de pointes: lack of common variants with large effect sizes. <i>PLoS ONE</i> , 2013 , 8, e78511	3.7	48
89	Molecular genetics of coronary artery disease. <i>Journal of Human Genetics</i> , 2016 , 61, 71-7	4.3	47
88	Multiple nonglycemic genomic loci are newly associated with blood level of glycated hemoglobin in East Asians. <i>Diabetes</i> , 2014 , 63, 2551-62	0.9	46
87	Trans-ethnic meta-analysis of white blood cell phenotypes. <i>Human Molecular Genetics</i> , 2014 , 23, 6944-605.6	4.5	45
86	Variation of gene-based SNPs and linkage disequilibrium patterns in the human genome. <i>Human Molecular Genetics</i> , 2004 , 13, 1623-32	5.6	44
85	Genomic structure and multiple single-nucleotide polymorphisms (SNPs) of the thiopurine S-methyltransferase (TPMT) gene. <i>Journal of Human Genetics</i> , 2000 , 45, 299-302	4.3	44
84	A functional variant in ZNF512B is associated with susceptibility to amyotrophic lateral sclerosis in Japanese. <i>Human Molecular Genetics</i> , 2011 , 20, 3684-92	5.6	42
83	Haplotypes with copy number and single nucleotide polymorphisms in CYP2A6 locus are associated with smoking quantity in a Japanese population. <i>PLoS ONE</i> , 2012 , 7, e44507	3.7	42
82	Overview of BioBank Japan follow-up data in 32 diseases. <i>Journal of Epidemiology</i> , 2017 , 27, S22-S28	3.4	41
81	Genome-wide association study to identify SNPs conferring risk of myocardial infarction and their functional analyses. <i>Cellular and Molecular Life Sciences</i> , 2005 , 62, 1804-13	10.3	41
80	Prediction model for knee osteoarthritis based on genetic and clinical information. <i>Arthritis Research and Therapy</i> , 2010 , 12, R187	5.7	40
79	A genome-wide association study identifies PLCL2 and AP3D1-DOT1L-SF3A2 as new susceptibility loci for myocardial infarction in Japanese. <i>European Journal of Human Genetics</i> , 2015 , 23, 374-80	5.3	39
78	Homozygosity mapping of a gene responsible for gelatinous drop-like corneal dystrophy to chromosome 1p. <i>American Journal of Human Genetics</i> , 1998 , 63, 1073-7	11	39
77	A functional SNP in EDG2 increases susceptibility to knee osteoarthritis in Japanese. <i>Human Molecular Genetics</i> , 2008 , 17, 1790-7	5.6	37

76	Voltage-shift of the current activation in HERG S4 mutation (R534C) in LQT2. <i>Cardiovascular Research</i> , 1999 , 44, 283-93	9.9	36
75	Significant impact of miRNA-target gene networks on genetics of human complex traits. <i>Scientific Reports</i> , 2016 , 6, 22223	4.9	36
74	Mapping of a gene for May-Hegglin anomaly to chromosome 22q. <i>Human Genetics</i> , 1999 , 105, 379-383	6.3	35
73	Genetic Variation in the SLC8A1 Calcium Signaling Pathway Is Associated With Susceptibility to Kawasaki Disease and Coronary Artery Abnormalities. <i>Circulation: Cardiovascular Genetics</i> , 2016 , 9, 559-568	33	
72	BRAP Activates Inflammatory Cascades and Increases the Risk for Carotid Atherosclerosis. <i>Molecular Medicine</i> , 2011 , 17, 1065-74	6.2	33
71	Induction of tenascin-C by tumor-specific EWS-ETS fusion genes. <i>Genes Chromosomes and Cancer</i> , 2003 , 36, 224-32	5	33
70	Genomic organization and mutational analysis of KVLQT1, a gene responsible for familial long QT syndrome. <i>Human Genetics</i> , 1998 , 103, 290-4	6.3	32
69	Inhibition of experimental intimal thickening in mice lacking a novel G-protein-coupled receptor. <i>Circulation</i> , 2003 , 107, 313-9	16.7	31
68	Impact of atherosclerosis-related gene polymorphisms on mortality and recurrent events after myocardial infarction. <i>Atherosclerosis</i> , 2006 , 185, 400-5	3.1	30
67	Single-nucleotide polymorphisms in the class II region of the major histocompatibility complex in Japanese patients with immunoglobulin A nephropathy. <i>Journal of Human Genetics</i> , 2002 , 47, 532-8	4.3	30
66	Variations in ORAI1 Gene Associated with Kawasaki Disease. <i>PLoS ONE</i> , 2016 , 11, e0145486	3.7	30
65	Identification of AF17 as a downstream gene of the beta-catenin/T-cell factor pathway and its involvement in colorectal carcinogenesis. <i>Cancer Research</i> , 2001 , 61, 6345-9	10.1	30
64	Variants at HLA-A, HLA-C, and HLA-DQB1 Confer Risk of Psoriasis Vulgaris in Japanese. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 542-548	4.3	29
63	PTPRD gene associated with blood pressure response to atenolol and resistant hypertension. <i>Journal of Hypertension</i> , 2015 , 33, 2278-85	1.9	29
62	Isolation of HELAD1, a novel human helicase gene up-regulated in colorectal carcinomas. <i>Oncogene</i> , 2002 , 21, 6387-94	9.2	29
61	Isolation and chromosomal mapping of the human homolog of perilipin (PLIN), a rat adipose tissue-specific gene, by differential display method. <i>Genomics</i> , 1998 , 48, 254-7	4.3	29
60	Mapping of a gene for May-Hegglin anomaly to chromosome 22q. <i>Human Genetics</i> , 1999 , 105, 379-83	6.3	29
59	Common variants in a novel gene, FONG on chromosome 2q33.1 confer risk of osteoporosis in Japanese. <i>PLoS ONE</i> , 2011 , 6, e19641	3.7	27

58	A functional SNP in ITIH3 is associated with susceptibility to myocardial infarction. <i>Journal of Human Genetics</i> , 2007 , 52, 220-229	4.3	27
57	Inflammation as a risk factor for myocardial infarction. <i>Journal of Human Genetics</i> , 2006 , 51, 595-604	4.3	27
56	Identification of 142 single nucleotide polymorphisms in 41 candidate genes for rheumatoid arthritis in the Japanese population. <i>Human Genetics</i> , 2000 , 106, 293-297	6.3	27
55	SNPs on chromosome 5p15.3 associated with myocardial infarction in Japanese population. <i>Journal of Human Genetics</i> , 2011 , 56, 47-51	4.3	26
54	Regulatory polymorphism in transcription factor KLF5 at the MEF2 element alters the response to angiotensin II and is associated with human hypertension. <i>FASEB Journal</i> , 2010 , 24, 1780-8	0.9	24
53	Genome-Wide Association Study of Peripheral Arterial Disease in a Japanese Population. <i>PLoS ONE</i> , 2015 , 10, e0139262	3.7	24
52	Association of a single-nucleotide polymorphism in the immunoglobulin mu-binding protein 2 gene with immunoglobulin A nephropathy. <i>Journal of Human Genetics</i> , 2005 , 50, 30-35	4.3	23
51	Genetic variations in five genes involved in the excitement of cardiomyocytes. <i>Journal of Human Genetics</i> , 2001 , 46, 549-52	4.3	20
50	Identification of 142 single nucleotide polymorphisms in 41 candidate genes for rheumatoid arthritis in the Japanese population. <i>Human Genetics</i> , 2000 , 106, 293-7	6.3	20
49	Construction of a normalized directionally cloned cDNA library from adult heart and analysis of 3040 clones by partial sequencing. <i>Genomics</i> , 1996 , 35, 231-5	4.3	20
48	Exome Analyses of Long QT Syndrome Reveal Candidate Pathogenic Mutations in Calmodulin-Interacting Genes. <i>PLoS ONE</i> , 2015 , 10, e0130329	3.7	20
47	Correlation of genetic etiology with response to beta-adrenergic blockade among symptomatic patients with familial long-QT syndrome. <i>Journal of Human Genetics</i> , 2001 , 46, 38-40	4.3	19
46	Variants in the SCN5A Promoter Associated With Various Arrhythmia Phenotypes. <i>Journal of the American Heart Association</i> , 2016 , 5,	6	18
45	Localization of the gene responsible for Peutz-Jeghers syndrome within a 6-cM region of chromosome 19p13.3. <i>Human Genetics</i> , 1998 , 102, 203-6	6.3	17
44	Multiple single-nucleotide polymorphisms (SNPs) in the Japanese population in six candidate genes for long QT syndrome. <i>Journal of Human Genetics</i> , 2001 , 46, 158-62	4.3	16
43	Characterization of S818L mutation in HERG C-terminus in LQT2. Modification of activation-deactivation gating properties. <i>FEBS Letters</i> , 2000 , 481, 197-203	3.8	15
42	Identification of a novel gene (ECM2) encoding a putative extracellular matrix protein expressed predominantly in adipose and female-specific tissues and its chromosomal localization to 9q22.3. <i>Genomics</i> , 1998 , 52, 378-81	4.3	15
41	Regional evaluation of childhood acute lymphoblastic leukemia genetic susceptibility loci among Japanese. <i>Scientific Reports</i> , 2018 , 8, 789	4.9	14

40	Gene expression patterns as marker for 5-year postoperative prognosis of primary breast cancers. <i>Journal of Cancer Research and Clinical Oncology</i> , 2004 , 130, 537-45	4.9	14
39	Molecular cloning and mapping of a human cDNA for cytosolic malate dehydrogenase (MDH1). <i>Genomics</i> , 1996 , 32, 128-30	4.3	14
38	Phenotypic Variability of ANK2 Mutations in Patients With Inherited Primary Arrhythmia Syndromes. <i>Circulation Journal</i> , 2016 , 80, 2435-2442	2.9	12
37	A genome-wide association study to identify genomic modulators of rate control therapy in patients with atrial fibrillation. <i>American Journal of Cardiology</i> , 2014 , 114, 593-600	3	11
36	Genetic Variants Associated With Susceptibility to Atrial Fibrillation in a Japanese Population. <i>Canadian Journal of Cardiology</i> , 2017 , 33, 443-449	3.8	10
35	Fine-scale SNP map of an 11-kb genomic region at 22q13.1 containing the galectin-1 gene. <i>Journal of Human Genetics</i> , 2005 , 50, 42-45	4.3	10
34	Refined mapping of caltractin in human Xq28 and in the homologous region of the mouse X chromosome places the gene within the bare patches (Bpa) and striated (Str) critical regions. <i>Mammalian Genome</i> , 1995 , 6, 802-4	3.2	10
33	Association between maternal education and malocclusion in Mongolian adolescents: a cross-sectional study. <i>BMJ Open</i> , 2016 , 6, e012283	3	10
32	Submicroscopic deletions at 13q32.1 cause congenital microcoria. <i>American Journal of Human Genetics</i> , 2015 , 96, 631-9	11	9
31	Recombination rates of genes expressed in human tissues. <i>Human Molecular Genetics</i> , 2008 , 17, 577-86	5.6	9
30	High-density SNP map of human ITR, a gene associated with vascular remodeling. <i>Journal of Human Genetics</i> , 2003 , 48, 170-2	4.3	9
29	Molecular cloning of a human cDNA encoding putative cysteine protease (PRSC1) and its chromosome assignment to 14q32.1. <i>Cytogenetic and Genome Research</i> , 1996 , 74, 120-3	1.9	9
28	Association of an IGHV3-66 gene variant with Kawasaki disease. <i>Journal of Human Genetics</i> , 2021 , 66, 475-489	4.3	9
27	A functional SNP in FLT1 increases risk of coronary artery disease in a Japanese population. <i>Journal of Human Genetics</i> , 2016 , 61, 435-41	4.3	8
26	Reduced risk of recurrent myocardial infarction in homozygous carriers of the chromosome 9p21 rs1333049 C risk allele in the contemporary percutaneous coronary intervention era: a prospective observational study. <i>BMJ Open</i> , 2014 , 4, e005438	3	8
25	Identification by differential display of eight known genes induced during in vivo intimal hyperplasia. <i>Journal of Human Genetics</i> , 1998 , 43, 9-13	4.3	8
24	Identification of 46 novel SNPs in the 130-kb region containing a myocardial infarction susceptibility gene on chromosomal band 6p21. <i>Journal of Human Genetics</i> , 2003 , 48, 476-479	4.3	8
23	Renin-angiotensin-aldosterone system polymorphisms and 5-year mortality in survivors of acute myocardial infarction: a report from the Osaka Acute Coronary Insufficiency Study. <i>International Heart Journal</i> , 2014 , 55, 190-6	1.8	7

22	Linkage disequilibrium of evolutionarily conserved regions in the human genome. <i>BMC Genomics</i> , 2006 , 7, 326	4.5	7
21	High-density single-nucleotide polymorphism (SNP) map in the 96-kb region containing the entire human DiGeorge syndrome critical region 2 (DGCR2) gene at 22q11.2. <i>Journal of Human Genetics</i> , 2001 , 46, 604-8	4.3	7
20	Genome-wide association analysis of common genetic variants of resistant hypertension. <i>Pharmacogenomics Journal</i> , 2019 , 19, 295-304	3.5	7
19	Investigation of novel variations of ORAI1 gene and their association with Kawasaki disease. <i>Journal of Human Genetics</i> , 2019 , 64, 511-519	4.3	6
18	Heterozygosities and allelic frequencies of 358 dinucleotide-repeat marker loci in the Japanese population. <i>Journal of Human Genetics</i> , 1998 , 43, 165-8	4.3	6
17	Genetic linkage analyses of Romano-Ward syndrome (RWS) in 13 Japanese families. <i>Human Genetics</i> , 1994 , 94, 380-4	6.3	6
16	Lymphotoxin-alpha3 mediates monocyte-endothelial interaction by TNFR I/NF-kappaB signaling. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 379, 374-8	3.4	5
15	Twenty single-nucleotide polymorphisms in four genes encoding cardiac ion channels. <i>Journal of Human Genetics</i> , 2002 , 47, 208-12	4.3	5
14	Genome-wide association study to identify single-nucleotide polymorphisms conferring risk of myocardial infarction. <i>Methods in Molecular Medicine</i> , 2006 , 128, 173-80		5
13	HLA-C variants associated with amino acid substitutions in the peptide binding groove influence susceptibility to Kawasaki disease. <i>Human Immunology</i> , 2019 , 80, 731-738	2.3	4
12	Assignment of the human caltractin gene (CALT) to Xq28 by fluorescence in situ hybridization. <i>Genomics</i> , 1994 , 24, 609-10	4.3	4
11	Clinical utility and functional analysis of variants in atrial fibrillation-associated locus 4q25. <i>Journal of Cardiology</i> , 2017 , 70, 366-373	3	3
10	Genome-wide association study identifies pharmacogenomic loci linked with specific antihypertensive drug treatment and new-onset diabetes. <i>Pharmacogenomics Journal</i> , 2018 , 18, 106-112 ^{3.5}		3
9	Decreased mortality associated with statin treatment in patients with acute myocardial infarction and lymphotoxin-alpha C804A polymorphism. <i>Atherosclerosis</i> , 2013 , 227, 373-9	3.1	3
8	Pathway analysis with genome-wide association study (GWAS) data detected the association of atrial fibrillation with the mTOR signaling pathway. <i>IJC Heart and Vasculature</i> , 2019 , 24, 100383	2.4	2
7	Identification of OPN3 as associated with non-syndromic oligodontia in a Japanese population. <i>Journal of Human Genetics</i> , 2021 , 66, 769-775	4.3	0
6	Identification of myocardial infarction-susceptible genes and their functional analyses ⁷⁹⁻⁸⁸		
5	Genetic backgrounds of myocardial infarction. <i>Current Cardiovascular Risk Reports</i> , 2007 , 1, 427-431	0.9	

LIST OF PUBLICATIONS

- | | | |
|---|--|-----|
| 4 | Chapter 7 Mutational Analysis of Familial Long QT Syndrome in Japan. <i>Current Topics in Membranes</i> ,
1999 , 46, 103-116 | 2.2 |
| 3 | Genetics of Coronary Disease 2019 , 21-36 | |
| 2 | Genetic Background of Myocardial Infarction 2011 , 113-120 | |
| 1 | Squamous Cell Carcinoma at the Site of Cutaneous Lymphoid Hyperplasia.. <i>Annals of Dermatology</i> ,
2022 , 34, 146-148 | 0.4 |