Maria Moutzouri

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5752906/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The effects of the Mulligan Sustained Natural Apophyseal Glide (SNAG) mobilisation in the lumbar flexion range of asymptomatic subjects as measured by the Zebris CMS20 3-D motion analysis system. BMC Musculoskeletal Disorders, 2008, 9, 131.	1.9	27
2	Investigation of the Effects of a Centrally Applied Lumbar Sustained Natural Apophyseal Glide Mobilization on Lower Limb Sympathetic Nervous System Activity in Asymptomatic Subjects. Journal of Manipulative and Physiological Therapeutics, 2012, 35, 286-294.	0.9	24
3	Cross-cultural translation and validation of the Greek version of the Knee Injury and Osteoarthritis Outcome Score (KOOS) in patients with total knee replacement. Disability and Rehabilitation, 2015, 37, 1477-1483.	1.8	24
4	What is the effect of sensori-motor training on functional outcome and balance performance of patients' undergoing TKR? A systematic review. Physiotherapy, 2016, 102, 136-144.	0.4	23
5	Early self-managed focal sensorimotor rehabilitative training enhances functional mobility and sensorimotor function in patients following total knee replacement: a controlled clinical trial. Clinical Rehabilitation, 2018, 32, 888-898.	2.2	8
6	"Greek KOOS-Child: a valid, disease specific, diagnostically accurate and responsive PROM in children with knee-related pathology― Knee Surgery, Sports Traumatology, Arthroscopy, 2021, 29, 1841-1849.	4.2	5
7	Greek Physiotherapists' Perspectives on Rehabilitation Following Total Knee Replacement: a Descriptive Survey. Physiotherapy Research International, 2017, 22, e1671.	1.5	4
8	Early initiation of home-based sensori-motor training improves muscle strength, activation and size in patients after knee replacement: a secondary analysis of a controlled clinical trial. BMC Musculoskeletal Disorders, 2019, 20, 231.	1.9	4
9	How effective is a blended web-based rehabilitation for improving pain, physical activity, and knee function of patients with knee osteoarthritis? Study protocol for a randomized control trial. PLoS ONE, 2022, 17, e0268652.	2.5	2