

Erik J Veneklaas

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5760362/erik-j-veneklaas-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

119
papers

11,668
citations

40
h-index

107
g-index

123
ext. papers

13,916
ext. citations

5.4
avg, IF

5.73
L-index

#	Paper	IF	Citations
119	Thermal imagery of woodland tree canopies provides new insights into drought-induced tree mortality.. <i>Science of the Total Environment</i> , 2022 , 155395	10.2	
118	How does spatial micro-environmental heterogeneity influence seedling recruitment in ironstone outcrops?. <i>Journal of Vegetation Science</i> , 2021 , 32, e13010	3.1	
117	Patterns and drivers of structure, diversity, and composition in species-rich shrublands restored after mining. <i>Restoration Ecology</i> , 2021 , 29, e13360	3.1	1
116	Leaf manganese concentrations as a tool to assess belowground plant functioning in phosphorus-impooverished environments. <i>Plant and Soil</i> , 2021 , 461, 43-61	4.2	23
115	Root positioning and trait shifts in <i>Hibbertia racemosa</i> as dependent on its neighbour's nutrient-acquisition strategy. <i>Plant, Cell and Environment</i> , 2021 , 44, 1257-1267	8.4	4
114	A threatened ecological community: research advances and priorities for <i>Banksia</i> woodlands. <i>Australian Journal of Botany</i> , 2021 , 69, 53	1.2	6
113	Corrigendum to: A threatened ecological community: research advances and priorities for <i>Banksia</i> woodlands. <i>Australian Journal of Botany</i> , 2021 , 69, 111	1.2	4
112	No evidence of regulation in root-mediated iron reduction in two Strategy I cluster-rooted <i>Banksia</i> species (Proteaceae). <i>Plant and Soil</i> , 2021 , 461, 203-218	4.2	1
111	Root length is proxy for high-throughput screening of waterlogging tolerance in <i>Urochloa</i> spp. grasses. <i>Functional Plant Biology</i> , 2021 , 48, 411-421	2.7	2
110	The barrier to radial oxygen loss impedes the apoplastic entry of iron into the roots of <i>Urochloa humidicola</i> . <i>Journal of Experimental Botany</i> , 2021 , 72, 3279-3293	7	4
109	Native plant diversity is a stronger driver for soil quality than inorganic amendments in semi-arid post-mining rehabilitation. <i>Geoderma</i> , 2021 , 394, 115001	6.7	0
108	Individual tree growth in jarrah (<i>Eucalyptus marginata</i>) forest is explained by size and distance of neighbouring trees in thinned and non-thinned plots. <i>Forest Ecology and Management</i> , 2021 , 494, 119364	2.9	4
107	AusTraits, a curated plant trait database for the Australian flora. <i>Scientific Data</i> , 2021 , 8, 254	8.2	6
106	Investigating the effect of neighbour competition on individual tree growth in thinned and unthinned eucalypt forests. <i>Forest Ecology and Management</i> , 2021 , 499, 119637	3.9	2
105	TRY plant trait database - enhanced coverage and open access. <i>Global Change Biology</i> , 2020 , 26, 119-188	11.4	399
104	Phosphate fertiliser alters carboxylates and bacterial communities in sweet potato (<i>Ipomoea batatas</i> (L.) Lam.) rhizosheaths. <i>Plant and Soil</i> , 2020 , 454, 245-260	4.2	2
103	Potassium Fertilisation Is Required to Sustain Cassava Yield and Soil Fertility. <i>Agronomy</i> , 2020 , 10, 1103	3.6	1

102	The potential for phosphorus benefits through root placement in the rhizosphere of phosphorus-mobilising neighbours. <i>Oecologia</i> , 2020 , 193, 843-855	2.9	4
101	Trait-based formal definition of plant functional types and functional communities in the multi-species and multi-traits context. <i>Ecological Complexity</i> , 2019 , 40, 100787	2.6	3
100	Root-zone hypoxia reduces growth of the tropical forage grass <i>Urochloa humidicola</i> in high-nutrient but not low-nutrient conditions. <i>Annals of Botany</i> , 2019 , 124, 1019-1032	4.1	10
99	Composition and ecological drivers of the kwongan scrub and woodlands in the northern Swan Coastal Plain, Western Australia. <i>Austral Ecology</i> , 2019 , 44, 906-916	1.5	3
98	Water availability drives the effectiveness of inorganic amendments to increase plant growth and substrate quality. <i>Catena</i> , 2019 , 182, 104116	5.8	3
97	Two sides to every leaf: water and CO transport in hypostomatous and amphistomatous leaves. <i>New Phytologist</i> , 2019 , 222, 1179-1187	9.8	37
96	Tree host-pathogen interactions as influenced by drought timing: linking physiological performance, biochemical defence and disease severity. <i>Tree Physiology</i> , 2019 , 39, 6-18	4.2	20
95	Root dynamics and survival in a nutrient-poor and species-rich woodland under a drying climate. <i>Plant and Soil</i> , 2018 , 424, 91-102	4.2	3
94	Shallow soils negatively affect water relations and photosynthesis in two semi-arid <i>Eucalyptus</i> species. <i>Environmental and Experimental Botany</i> , 2018 , 155, 239-250	5.9	5
93	Community patterns and environmental drivers in hyper-diverse kwongan scrub vegetation of Western Australia. <i>Applied Vegetation Science</i> , 2018 , 21, 694-722	3.3	10
92	Hydraulic redistribution: limitations for plants in saline soils. <i>Plant, Cell and Environment</i> , 2017 , 40, 2437-2446	8.4	8
91	Spectral detection of stress-related pigments in salt-lake succulent halophytic shrubs. <i>International Journal of Applied Earth Observation and Geoinformation</i> , 2016 , 52, 457-463	7.3	2
90	Apparent Overinvestment in Leaf Venation Relaxes Leaf Morphological Constraints on Photosynthesis in Arid Habitats. <i>Plant Physiology</i> , 2016 , 172, 2286-2299	6.6	27
89	Salinity tolerances of three succulent halophytes (<i>Tecticornia</i> spp.) differentially distributed along a salinity gradient. <i>Functional Plant Biology</i> , 2016 , 43, 739-750	2.7	12
88	Informing arid region mine-site restoration through comparative ecophysiology of <i>Acacia</i> species under drought. <i>Journal of Arid Environments</i> , 2016 , 133, 73-84	2.5	3
87	Links between soil texture and root architecture of <i>Eucalyptus</i> species may limit distribution ranges under future climates. <i>Plant and Soil</i> , 2016 , 403, 217-229	4.2	12
86	Genetic delineation of local provenance defines seed collection zones along a climate gradient. <i>AoB PLANTS</i> , 2016 , 8,	2.9	5
85	Osmotic potential at full turgor: an easily measurable trait to help breeders select for drought tolerance in wheat. <i>Plant Breeding</i> , 2016 , 135, 279-285	2.4	16

84	Optimal allocation of leaf epidermal area for gas exchange. <i>New Phytologist</i> , 2016 , 210, 1219-28	9.8	88
83	A climate change context for the decline of a foundation tree species in south-western Australia: insights from phylogeography and species distribution modelling. <i>Annals of Botany</i> , 2015 , 116, 941-52	4.1	16
82	Interactions among cluster-root investment, leaf phosphorus concentration, and relative growth rate in two <i>Lupinus</i> species. <i>American Journal of Botany</i> , 2015 , 102, 1529-37	2.7	2
81	Isometric partitioning of hydraulic conductance between leaves and stems: balancing safety and efficiency in different growth forms and habitats. <i>Plant, Cell and Environment</i> , 2015 , 38, 1628-36	8.4	11
80	Shallow environmental gradients put inland species at risk: Insights and implications from predicting future distributions of <i>Eucalyptus</i> species in South Western Australia. <i>Austral Ecology</i> , 2015 , 40, 923-932	1.5	7
79	A Critical Evaluation of Interventions to Progress Transdisciplinary Research. <i>Society and Natural Resources</i> , 2015 , 28, 670-681	2.4	13
78	Contrasting submergence tolerance in two species of stem-succulent halophytes is not determined by differences in stem internal oxygen dynamics. <i>Annals of Botany</i> , 2015 , 115, 409-18	4.1	6
77	Is nitrogen transfer among plants enhanced by contrasting nutrient-acquisition strategies?. <i>Plant, Cell and Environment</i> , 2015 , 38, 50-60	8.4	18
76	Global variability in leaf respiration in relation to climate, plant functional types and leaf traits. <i>New Phytologist</i> , 2015 , 206, 614-36	9.8	244
75	Spatio-temporal water dynamics in mature <i>Banksia menziesii</i> trees during drought. <i>Physiologia Plantarum</i> , 2014 , 152, 301-15	4.6	16
74	Low levels of ribosomal RNA partly account for the very high photosynthetic phosphorus-use efficiency of Proteaceae species. <i>Plant, Cell and Environment</i> , 2014 , 37, 1276-98	8.4	87
73	Drought tolerances of three stem-succulent halophyte species of an inland semiarid salt lake system. <i>Functional Plant Biology</i> , 2014 , 41, 1230-1238	2.7	10
72	Complementary plant nutrient-acquisition strategies promote growth of neighbour species. <i>Functional Ecology</i> , 2014 , 28, 819-828	5.6	48
71	Physiological and ecological significance of biomineralization in plants. <i>Trends in Plant Science</i> , 2014 , 19, 166-74	13.1	111
70	Are leaf functional traits invariant with plant size and what is invariance anyway?. <i>Functional Ecology</i> , 2014 , 28, 1330-1343	5.6	34
69	Transpiration and water relations of evergreen shrub species on an artificial landform for mine waste storage versus an adjacent natural site in semi-arid Western Australia. <i>Ecohydrology</i> , 2014 , 7, 965-981	2.5	17
68	Does cluster-root activity benefit nutrient uptake and growth of co-existing species?. <i>Oecologia</i> , 2014 , 174, 23-31	2.9	62
67	Species distribution and crown decline are associated with contrasting water relations in four common sympatric eucalypt species in southwestern Australia. <i>Plant and Soil</i> , 2013 , 364, 409-423	4.2	31

66	Transpiration and plant water relations of evergreen woody vegetation on a recently constructed artificial ecosystem under seasonally dry conditions in Western Australia. <i>Hydrological Processes</i> , 2012 , 26, 3281-3292	3.3	12
65	Morphologies and elemental compositions of calcium crystals in phyllodes and branchlets of <i>Acacia robeorum</i> (Leguminosae: Mimosoideae). <i>Annals of Botany</i> , 2012 , 109, 887-96	4.1	50
64	Opportunities for improving phosphorus-use efficiency in crop plants. <i>New Phytologist</i> , 2012 , 195, 306-308	3.8	479
63	Inbreeding and outbreeding depression in <i>Stylidium hispidum</i> : implications for mixing seed sources for ecological restoration. <i>Ecology and Evolution</i> , 2012 , 2, 2262-73	2.8	39
62	Tropical cyclones and the ecohydrology of Australia's recent continental-scale drought. <i>Geophysical Research Letters</i> , 2012 , 39, n/a-n/a	4.9	48
61	Arid-zone <i>Acacia</i> species can access poorly soluble iron phosphate but show limited growth response. <i>Plant and Soil</i> , 2012 , 358, 119-130	4.2	8
60	Precipitation of calcium, magnesium, strontium and barium in tissues of four <i>Acacia</i> species (Leguminosae: Mimosoideae). <i>PLoS ONE</i> , 2012 , 7, e41563	3.7	24
59	Nature and nurture: the importance of seed phosphorus content. <i>Plant and Soil</i> , 2012 , 357, 1-8	4.2	129
58	Dinitrogen-fixing <i>Acacia</i> species from phosphorus-impooverished soils resorb leaf phosphorus efficiently. <i>Plant, Cell and Environment</i> , 2011 , 34, 2060-70	8.4	21
57	Drying the surface soil reduces the nitrogen content of faba bean (<i>Vicia faba</i> L.) through a reduction in nitrogen fixation. <i>Plant and Soil</i> , 2011 , 339, 351-362	4.2	13
56	Spatial analysis of fine root distribution on a recently constructed ecosystem in a water-limited environment. <i>Plant and Soil</i> , 2011 , 344, 255-272	4.2	36
55	Strategies and agronomic interventions to improve the phosphorus-use efficiency of farming systems. <i>Plant and Soil</i> , 2011 , 349, 89-120	4.2	262
54	Spatial analysis of fine root distribution on a recently constructed ecosystem in a water-limited environment. <i>Plant and Soil</i> , 2011 , 348, 471-489	4.2	7
53	Plant and microbial strategies to improve the phosphorus efficiency of agriculture. <i>Plant and Soil</i> , 2011 , 349, 121-156	4.2	532
52	Update on phosphorus nutrition in Proteaceae. Phosphorus nutrition of proteaceae in severely phosphorus-impooverished soils: are there lessons to be learned for future crops?. <i>Plant Physiology</i> , 2011 , 156, 1058-66	6.6	146
51	The fate of hydraulically redistributed water in a semi-arid zone eucalyptus species. <i>Tree Physiology</i> , 2011 , 31, 649-58	4.2	19
50	Contrasting physiological responses of two co-occurring eucalypts to seasonal drought at restored bauxite mine sites. <i>Tree Physiology</i> , 2011 , 31, 1052-66	4.2	18
49	Photosynthesis at an extreme end of the leaf trait spectrum: how does it relate to high leaf dry mass per area and associated structural parameters?. <i>Journal of Experimental Botany</i> , 2010 , 61, 3015-28	7	54

48	Effects of phosphorus supply on growth, phosphate concentration and cluster-root formation in three <i>Lupinus</i> species. <i>Annals of Botany</i> , 2010 , 105, 365-74	4.1	40
47	Changes in water relations for <i>Acacia ancistrocarpa</i> on natural and mine-rehabilitation sites in response to an experimental wetting pulse in the Great Sandy Desert. <i>Plant and Soil</i> , 2010 , 326, 75-96	4.2	13
46	Influence of leaf dry mass per area, CO ₂ , and irradiance on mesophyll conductance in sclerophylls. <i>Journal of Experimental Botany</i> , 2009 , 60, 2303-14	7	129
45	Stomatal crypts have small effects on transpiration: a numerical model analysis. <i>Plant Physiology</i> , 2009 , 151, 2018-27	6.6	51
44	Stomatal crypts may facilitate diffusion of CO ₂ to adaxial mesophyll cells in thick sclerophylls. <i>Plant, Cell and Environment</i> , 2009 , 32, 1596-611	8.4	58
43	Soil seed bank compositional change constrains biodiversity in an invaded species-rich woodland. <i>Biological Conservation</i> , 2009 , 142, 256-269	6.2	54
42	Altered vegetation structure and composition linked to fire frequency and plant invasion in a biodiverse woodland. <i>Biological Conservation</i> , 2009 , 142, 2270-2281	6.2	70
41	Population Size Effects on Progeny Performance in <i>Banksia ilicifolia</i> R. Br. (Proteaceae). <i>HAYATI Journal of Biosciences</i> , 2009 , 16, 43-48	1.2	1
40	Leaf water relations during summer water deficit: differential responses in turgor maintenance and variation in leaf structure among different plant communities in south-western Australia. <i>Plant, Cell and Environment</i> , 2008 , 31, 1791-802	8.4	104
39	Water relations and mineral nutrition of closely related woody plant species on desert dunes and interdunes. <i>Australian Journal of Botany</i> , 2008 , 56, 27	1.2	40
38	Water relations and mineral nutrition of <i>Triodia</i> grasses on desert dunes and interdunes. <i>Australian Journal of Botany</i> , 2008 , 56, 408	1.2	22
37	Using multiple trait associations to define hydraulic functional types in plant communities of south-western Australia. <i>Oecologia</i> , 2008 , 158, 385-97	2.9	62
36	Rhizosphere processes do not explain variation in P acquisition from sparingly soluble forms among <i>Lupinus albus</i> accessions. <i>Australian Journal of Agricultural Research</i> , 2008 , 59, 616		7
35	Carboxylate composition of root exudates does not relate consistently to a crop species' ability to use phosphorus from aluminium, iron or calcium phosphate sources. <i>New Phytologist</i> , 2007 , 173, 181-90	9.8	136
34	<i>Banksia</i> species (Proteaceae) from severely phosphorus-impooverished soils exhibit extreme efficiency in the use and re-mobilization of phosphorus. <i>Plant, Cell and Environment</i> , 2007 , 30, 1557-65	8.4	115
33	Root Architecture of Jarrah (<i>Eucalyptus marginata</i>) Trees in Relation to Post-Mining Deep Ripping in Western Australia. <i>Restoration Ecology</i> , 2007 , 15, S65-S73	3.1	25
32	Does phenotypic plasticity in carboxylate exudation differ among rare and widespread <i>Banksia</i> species (Proteaceae)? <i>New Phytologist</i> , 2007 , 173, 592-599	9.8	27
31	Root structure and functioning for efficient acquisition of phosphorus: Matching morphological and physiological traits. <i>Annals of Botany</i> , 2006 , 98, 693-713	4.1	800

30	Contrasting water relations of three coastal tree species with different exposure to salinity. <i>Physiologia Plantarum</i> , 2006 , 127, 360-373	4.6	18
29	Variable tolerance of wetland tree species to combined salinity and waterlogging is related to regulation of ion uptake and production of organic solutes. <i>New Phytologist</i> , 2006 , 169, 123-33	9.8	64
28	Triticum aestivum shows a greater biomass response to a supply of aluminium phosphate than Lupinus albus, despite releasing fewer carboxylates into the rhizosphere. <i>New Phytologist</i> , 2006 , 169, 515-24	9.8	62
27	Distribution of Carboxylates and Acid Phosphatase and Depletion of Different Phosphorus Fractions in the Rhizosphere of a Cereal and Three Grain Legumes. <i>Plant and Soil</i> , 2006 , 281, 109-120	4.2	142
26	Enhanced soil and leaf nutrient status of a Western Australian Banksia woodland community invaded by Ehrharta calycina and Pelargonium capitatum. <i>Plant and Soil</i> , 2006 , 284, 253-264	4.2	51
25	Carboxylate release of wheat, canola and 11 grain legume species as affected by phosphorus status. <i>Plant and Soil</i> , 2006 , 288, 127-139	4.2	145
24	Preferential outcrossing in Banksia ilicifolia (Proteaceae). <i>Australian Journal of Botany</i> , 2005 , 53, 163	1.2	15
23	Gallery forest types and their environmental correlates in a Colombian savanna landscape. <i>Ecography</i> , 2005 , 28, 236-252	6.5	33
22	Dynamics of non-structural carbohydrates in two Ficus species after transfer to deep shade. <i>Environmental and Experimental Botany</i> , 2005 , 54, 148-154	5.9	18
21	Phosphorus benefits of different legume crops to subsequent wheat grown in different soils of Western Australia. <i>Plant and Soil</i> , 2005 , 271, 175-187	4.2	139
20	Plant phosphorus status has a limited influence on the concentration of phosphorus-mobilising carboxylates in the rhizosphere of chickpea. <i>Functional Plant Biology</i> , 2005 , 32, 153-159	2.7	40
19	Phosphorus uptake by grain legumes and subsequently grown wheat at different levels of residual phosphorus fertiliser. <i>Australian Journal of Agricultural Research</i> , 2005 , 56, 1041		74
18	The worldwide leaf economics spectrum. <i>Nature</i> , 2004 , 428, 821-7	50.4	4915
17	Carboxylate concentrations in the rhizosphere of lateral roots of chickpea (Cicer arietinum) increase during plant development, but are not correlated with phosphorus status of soil or plants. <i>New Phytologist</i> , 2004 , 162, 745-753	9.8	59
16	Rhizosphere carboxylate concentrations of chickpea are affected by genotype and soil type. <i>Plant and Soil</i> , 2004 , 261, 1-10	4.2	35
15	Short Communication: Leaf trait relationships in Australian plant species. <i>Functional Plant Biology</i> , 2004 , 31, 551-558	2.7	103
14	Seasonal patterns in water use and leaf turnover of different plant functional types in a species-rich woodland, south-western Australia. <i>Plant and Soil</i> , 2003 , 257, 295-304	4.2	40
13	Chickpea and white lupin rhizosphere carboxylates vary with soil properties and enhance phosphorus uptake. <i>Plant and Soil</i> , 2003 , 248, 187-197	4.2	209

12	Chickpea and white lupin rhizosphere carboxylates vary with soil properties and enhance phosphorus uptake 2003 , 187-197		0
11	The pattern of carboxylate exudation in <i>Banksia grandis</i> (Proteaceae) is affected by the form of phosphate added to the soil. <i>Plant and Soil</i> , 2002 , 238, 111-122	4.2	109
10	Relative growth rate and biomass allocation in 20 <i>Aegilops</i> (Poaceae) species. <i>New Phytologist</i> , 1998 , 140, 425-437	9.8	41
9	Relative growth rate and biomass allocation in 20 <i>Aegilops</i> (Poaceae) species. <i>New Phytologist</i> , 1998 , 140, 425-437	9.8	39
8	Yield and water use of wheat (<i>Triticum aestivum</i>) in a Mediterranean environment: Cultivar differences and sowing density effects. <i>Plant and Soil</i> , 1996 , 181, 251-262	4.2	26
7	Relative growth rate, biomass allocation pattern and water use efficiency of three wheat cultivars during early ontogeny as dependent on water availability. <i>Physiologia Plantarum</i> , 1996 , 98, 493-504	4.6	11
6	Relative growth rate, biomass allocation pattern and water use efficiency of three wheat cultivars during early ontogeny as dependent on water availability. <i>Physiologia Plantarum</i> , 1996 , 98, 493-504	4.6	23
5	Effects of global environmental change on carbon partitioning in vegetative plants of <i>Triticum aestivum</i> and closely related <i>Aegilops</i> species. <i>Global Change Biology</i> , 1995 , 1, 397-406	11.4	23
4	Litterfall and nutrient fluxes in two montane tropical rain forests, Colombia. <i>Journal of Tropical Ecology</i> , 1991 , 7, 319-336	1.3	81
3	Stockpiling disrupts the biological integrity of topsoil for ecological restoration. <i>Plant and Soil</i> , 1991 , 135, 1-10	4.2	0
2	Microbial inoculation to improve plant performance in mine-waste substrates: A test using pigeon pea (<i>Cajanus cajan</i>). <i>Land Degradation and Development</i> , 1991 , 2, 1-10	4.4	3
1	Restoration ecophysiology: an ecophysiological approach to improve restoration strategies and outcomes in severely disturbed landscapes. <i>Restoration Ecology</i> , 1991 , 1, 135-141	3.1	1