
## Naoko Tachibana

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5766592/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Polygenic burden of Parkinson's disease risk stratifies the prognosis of isolated rapid-eye-movement disorder: A preliminary observational study. Parkinsonism and Related Disorders, 2022, 96, 52-56.	2.2	2
2	Potential patient screening for late-onset Pompe disease in suspected sleep apnea: a rationale and study design for a Prospective Multicenter Observational Cohort Study in Japan (PSSAP-J Study). Sleep and Breathing, 2021, 25, 695-704.	1.7	1
3	Appropriate assessment method of 123I-MIBG myocardial scintigraphy for the diagnosis of Lewy body diseases and idiopathic REM sleep behavior disorder. Journal of Neurology, 2020, 267, 3248-3257.	3.6	5
4	Rate of augmentation and risk factors with long-term follow-up in Japanese patients with restless legs syndrome. Neurological Sciences, 2018, 39, 1559-1564.	1.9	3
5	Living with Restless Legs Syndrome/Willis-Ekbom Disease. Sleep Medicine Clinics, 2015, 10, 359-367.	2.6	7
6	History of Japanese Clinical Sleep Medicine. , 2015, , 129-131.		0
7	Decrease in myocardial 123I-MIBG radioactivity in REM sleep behavior disorder: Two patients with different clinical progression. Sleep Medicine, 2008, 9, 583-585.	1.6	20
8	Why do we continue to use Epworth sleepiness scale?. Sleep Medicine, 2007, 8, 541-542.	1.6	18
9	Decreased myocardial <sup>123</sup> lâ€MIBG uptake and impaired facial expression recognition in a patient with REM sleep behavior disorder. Movement Disorders, 2007, 22, 746-747.	3.9	13
10	REM sleep behaviour disorder in Parkinson's disease. Journal of Neurology, 2007, 254, IV8-IV14.	3.6	2
11	Hypersomnolence and increased REM sleep with low cerebrospinal fluid hypocretin level in a patient after removal of craniopharyngioma. Sleep Medicine, 2005, 6, 567-569.	1.6	33
12	Imbalance between the Reality of Sleep Specialists and the Demands of Society in Japan. Industrial Health, 2005, 43, 49-52.	1.0	4
13	How much further should we go to develop and establish sleep medicine in Japan? (Through a) Tj ETQq1 1 0.784	314 rgBT / 1.0	Overlock 1
14	Longitudinal change in REM sleep components in a patient with multiple system atrophy associated with REM sleep behavior disorder: paradoxical improvement of nocturnal behaviors in a progressive neurodegenerative disease. Sleep Medicine, 2004, 5, 155-158.	1.6	31
15	Japanese versus USA clinical services for sleep medicine. Sleep and Biological Rhythms, 2003, 1, 215-220.	1.0	12
16	What prevents Japan from establishing a clinical service of sleep medicine: Experience gained at a local hospital. Psychiatry and Clinical Neurosciences, 1999, 53, 263-265.	1.8	2
17	Subclinical REM Sleep Behavior Disorder in a Patient With Corticobasal Degeneration. Sleep, 1997, 20, 891-894.	1.1	92