

# Qi Long Wu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5767175/publications.pdf>

Version: 2024-02-01

11  
papers

482  
citations

1163117

8  
h-index

1281871

11  
g-index

11  
all docs

11  
docs citations

11  
times ranked

835  
citing authors

#	ARTICLE	IF	CITATIONS
1	A CO <sub>2</sub> adsorption dominated carbon defect-based electrocatalyst for efficient carbon dioxide reduction. <i>Journal of Materials Chemistry A</i> , 2020, 8, 1205-1211.	10.3	75
2	Novel Bi-Doped Amorphous SnO <sub>x</sub> Nanoshells for Efficient Electrochemical CO <sub>2</sub> Reduction into Formate at Low Overpotentials. <i>Advanced Materials</i> , 2020, 32, e2002822.	21.0	104
3	Trimetallic Synergy in Intermetallic PtSnBi Nanoplates Boosts Formic Acid Oxidation. <i>Advanced Materials</i> , 2019, 31, e1903683.	21.0	112
4	<i>In situ</i> aluminium ions regulation for quantum efficiency and light stability promotion in white light emitting material. <i>RSC Advances</i> , 2019, 9, 15265-15268.	3.6	1
5	Heterobimetallic metal-organic framework nanocages as highly efficient catalysts for CO <sub>2</sub> conversion under mild conditions. <i>Journal of Materials Chemistry A</i> , 2018, 6, 2964-2973.	10.3	73
6	Gas-solid aldol condensation reaction in confined space of metal organic framework for formaldehyde detection. <i>Nanoscale</i> , 2018, 10, 19286-19289.	5.6	3
7	Individual High-Quality N-Doped Carbon Nanotubes Embedded with Nonprecious Metal Nanoparticles toward Electrochemical Reaction. <i>ACS Applied Materials &amp; Interfaces</i> , 2018, 10, 39757-39767.	8.0	33
8	Carbon Defect-Induced Reversible Carbon-Oxygen Interfaces for Efficient Oxygen Reduction. <i>ACS Applied Materials &amp; Interfaces</i> , 2018, 10, 39735-39744.	8.0	45
9	Photocatalytic hydrogen evolution from formate and aldehyde over molecular iridium complexes stabilized by bipyridine-bridging organosilica nanotubes. <i>Applied Catalysis B: Environmental</i> , 2018, 236, 466-474.	20.2	22
10	Slow magnetic relaxations in azide or formate bridged chains based on dicubane-like 3d-4f clusters. <i>Journal of Molecular Structure</i> , 2018, 1171, 551-555.	3.6	1
11	Divalent metal ions modulated strong frustrated M( $\mu_2$ -Fe( $\mu_3$ )-O <sub>3</sub> ) (M = Fe, Mn, Mg) chains with metamagnetism only in a mixed valence iron complex. <i>Chemical Communications</i> , 2015, 51, 15336-15339.	4.1	13