

Harm van Marwijk

List of Publications by Citations

Source: <https://exaly.com/author-pdf/577450/harm-van-marwijk-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

271
papers

9,780
citations

51
h-index

88
g-index

299
ext. papers

11,431
ext. citations

3.7
avg, IF

5.98
L-index

#	Paper	IF	Citations
271	The Netherlands Study of Depression and Anxiety (NESDA): rationale, objectives and methods. <i>International Journal of Methods in Psychiatric Research</i> , 2008 , 17, 121-40	4.3	694
270	Depression and the risk for cardiovascular diseases: systematic review and meta analysis. <i>International Journal of Geriatric Psychiatry</i> , 2007 , 22, 613-26	3.9	642
269	Controlled Interventions to Reduce Burnout in Physicians: A Systematic Review and Meta-analysis. <i>JAMA Internal Medicine</i> , 2017 , 177, 195-205	11.5	584
268	Patients' preferences in the treatment of depressive disorder in primary care. <i>General Hospital Psychiatry</i> , 2004 , 26, 184-9	5.6	297
267	The Four-Dimensional Symptom Questionnaire (4DSQ): a validation study of a multidimensional self-report questionnaire to assess distress, depression, anxiety and somatization. <i>BMC Psychiatry</i> , 2006 , 6, 34	4.2	293
266	Two-year course of depressive and anxiety disorders: results from the Netherlands Study of Depression and Anxiety (NESDA). <i>Journal of Affective Disorders</i> , 2011 , 133, 76-85	6.6	223
265	Stepped-care prevention of anxiety and depression in late life: a randomized controlled trial. <i>Archives of General Psychiatry</i> , 2009 , 66, 297-304		212
264	The association of depression and anxiety with pain: a study from NESDA. <i>PLoS ONE</i> , 2014 , 9, e106907	3.7	164
263	Pain and the onset of depressive and anxiety disorders. <i>Pain</i> , 2014 , 155, 53-59	8	135
262	Impact of pain on the course of depressive and anxiety disorders. <i>Pain</i> , 2012 , 153, 429-436	8	126
261	A patient-doctor relationship questionnaire (PDRQ-9) in primary care: development and psychometric evaluation. <i>General Hospital Psychiatry</i> , 2004 , 26, 115-20	5.6	118
260	The European General Practice Research Network presents a comprehensive definition of multimorbidity in family medicine and long term care, following a systematic review of relevant literature. <i>Journal of the American Medical Directors Association</i> , 2013 , 14, 319-25	5.9	117
259	The Netherlands study of depression in older persons (NESDO); a prospective cohort study. <i>BMC Research Notes</i> , 2011 , 4, 524	2.3	113
258	Incidence of depression and anxiety in the spouses of patients with dementia: a naturalistic cohort study of recorded morbidity with a 6-year follow-up. <i>American Journal of Geriatric Psychiatry</i> , 2010 , 18, 146-53	6.5	110
257	Effectiveness of psychological interventions in preventing recurrence of depressive disorder: meta-analysis and meta-regression. <i>Journal of Affective Disorders</i> , 2015 , 174, 400-10	6.6	106
256	The prognosis of depression in older patients in general practice and the community. A systematic review. <i>Family Practice</i> , 2007 , 24, 168-80	1.9	102
255	Non-pharmacological interventions for somatoform disorders and medically unexplained physical symptoms (MUPS) in adults. <i>The Cochrane Library</i> , 2014 , CD011142	5.2	97

254	Longitudinal association between pain, and depression and anxiety over four years. <i>Journal of Psychosomatic Research</i> , 2015 , 78, 64-70	4.1	92
253	Anxiety and the risk of death in older men and women. <i>British Journal of Psychiatry</i> , 2004 , 185, 399-404	5.4	92
252	The two-year incidence of depression and anxiety disorders in spousal caregivers of persons with dementia: who is at the greatest risk?. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 293-303	6.5	89
251	Cost-effectiveness of a stepped care intervention to prevent depression and anxiety in late life: randomised trial. <i>British Journal of Psychiatry</i> , 2010 , 196, 319-25	5.4	81
250	Non-pharmacological interventions for somatoform disorders and medically-unexplained physical symptoms (MUPS) in adults 2014 ,		79
249	Associations of financial strain and income with depressive and anxiety disorders. <i>Journal of Epidemiology and Community Health</i> , 2015 , 69, 660-5	5.1	76
248	Pharmacological interventions for somatoform disorders in adults. <i>The Cochrane Library</i> , 2014 , CD010628	5.2	76
247	Detecting depressive and anxiety disorders in distressed patients in primary care; comparative diagnostic accuracy of the Four-Dimensional Symptom Questionnaire (4DSQ) and the Hospital Anxiety and Depression Scale (HADS). <i>BMC Family Practice</i> , 2009 , 10, 58	2.6	76
246	The course of untreated anxiety and depression, and determinants of poor one-year outcome: a one-year cohort study. <i>BMC Psychiatry</i> , 2010 , 10, 86	4.2	74
245	The two-year course of late-life depression; results from the Netherlands study of depression in older persons. <i>BMC Psychiatry</i> , 2015 , 15, 20	4.2	73
244	Depression in old age (75+), the PIKO study. <i>Journal of Affective Disorders</i> , 2008 , 106, 295-9	6.6	72
243	Is the Beck Anxiety Inventory a good tool to assess the severity of anxiety? A primary care study in the Netherlands Study of Depression and Anxiety (NESDA). <i>BMC Family Practice</i> , 2011 , 12, 66	2.6	71
242	Outcome of depression in later life in primary care: longitudinal cohort study with three years' follow-up. <i>BMJ, The</i> , 2009 , 338, a3079	5.9	71
241	Loneliness is associated with poor prognosis in late-life depression: Longitudinal analysis of the Netherlands study of depression in older persons. <i>Journal of Affective Disorders</i> , 2015 , 185, 1-7	6.6	70
240	Depression, anxiety, and arterial stiffness. <i>Biological Psychiatry</i> , 2011 , 69, 795-803	7.9	70
239	Interpersonal psychotherapy for elderly patients in primary care. <i>American Journal of Geriatric Psychiatry</i> , 2006 , 14, 777-86	6.5	70
238	Prevention of late-life anxiety and depression has sustained effects over 24 months: a pragmatic randomized trial. <i>American Journal of Geriatric Psychiatry</i> , 2011 , 19, 230-9	6.5	67
237	The comorbidity burden of type 2 diabetes mellitus: patterns, clusters and predictions from a large English primary care cohort. <i>BMC Medicine</i> , 2019 , 17, 145	11.4	65

236	Differences in heart rate variability between depressed and non-depressed elderly. <i>International Journal of Geriatric Psychiatry</i> , 2006 , 21, 147-50	3.9	65
235	Does a family meetings intervention prevent depression and anxiety in family caregivers of dementia patients? A randomized trial. <i>PLoS ONE</i> , 2012 , 7, e30936	3.7	62
234	Big Five personality and depression diagnosis, severity and age of onset in older adults. <i>Journal of Affective Disorders</i> , 2013 , 151, 178-85	6.6	59
233	Pain, not chronic disease, is associated with the recurrence of depressive and anxiety disorders. <i>BMC Psychiatry</i> , 2014 , 14, 187	4.2	59
232	The accuracy of family physicians' dementia diagnoses at different stages of dementia: a systematic review. <i>International Journal of Geriatric Psychiatry</i> , 2012 , 27, 342-54	3.9	59
231	Effect of psychiatric consultation models in primary care. A systematic review and meta-analysis of randomized clinical trials. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 521-33	4.1	57
230	An Exploration of Formal and Informal Mindfulness Practice and Associations with Wellbeing. <i>Mindfulness</i> , 2019 , 10, 89-99	2.9	56
229	Adherence to evidence-based guidelines for depression and anxiety disorders is associated with recording of the diagnosis. <i>General Hospital Psychiatry</i> , 2009 , 31, 460-9	5.6	56
228	The center for epidemiological studies depression scale (CES-D) is an adequate screening instrument for depressive and anxiety disorders in a very old population living in residential homes. <i>International Journal of Geriatric Psychiatry</i> , 2011 , 26, 239-46	3.9	54
227	Impact of euthanasia on primary care physicians in the Netherlands. <i>Palliative Medicine</i> , 2007 , 21, 609-14	5.5	54
226	Receiving treatment for common mental disorders. <i>General Hospital Psychiatry</i> , 2009 , 31, 46-55	5.6	53
225	Which physician and practice characteristics are associated with adherence to evidence-based guidelines for depressive and anxiety disorders?. <i>Medical Care</i> , 2010 , 48, 240-8	3.1	52
224	Associations between life stress and subclinical cardiovascular disease are partly mediated by depressive and anxiety symptoms. <i>Journal of Psychosomatic Research</i> , 2015 , 78, 332-9	4.1	51
223	Preferences regarding disclosure of a diagnosis of dementia: a systematic review. <i>International Psychogeriatrics</i> , 2014 , 26, 1603-18	3.4	51
222	Effectiveness of case management among older adults with early symptoms of dementia and their primary informal caregivers: a randomized clinical trial. <i>International Journal of Nursing Studies</i> , 2011 , 48, 933-43	5.8	51
221	Stepped care treatment for depression and anxiety in primary care. a randomized controlled trial. <i>Trials</i> , 2011 , 12, 171	2.8	51
220	Prevention of adverse health trajectories in a vulnerable elderly population through nurse home visits: a randomized controlled trial [ISRCTN05358495]. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2010 , 65, 734-42	6.4	51
219	Prevalence of depression in older patients consulting their general practitioner in The Netherlands. <i>International Journal of Geriatric Psychiatry</i> , 2005 , 20, 1013-9	3.9	51

218	The challenge of ageing populations and patient frailty: can primary care adapt?. <i>BMJ, The</i> , 2018 , 362, k3349	5.9	50
217	Cost-effectiveness of collaborative care for the treatment of major depressive disorder in primary care. A systematic review. <i>BMC Health Services Research</i> , 2010 , 10, 19	2.9	50
216	A cluster-randomised trial evaluating an intervention for patients with stress-related mental disorders and sick leave in primary care. <i>PLOS Clinical Trials</i> , 2007 , 2, e26		50
215	The prognosis of undetected depression in older general practice patients. A one year follow-up study. <i>Journal of Affective Disorders</i> , 2009 , 114, 310-5	6.6	48
214	A target-driven collaborative care model for Major Depressive Disorder is effective in primary care in the Netherlands. A randomized clinical trial from the depression initiative. <i>Journal of Affective Disorders</i> , 2013 , 146, 328-37	6.6	46
213	Do GPs' medical records demonstrate a good recognition of depression? A new perspective on case extraction. <i>Journal of Affective Disorders</i> , 2011 , 133, 522-7	6.6	46
212	Test-retest reliability of the PRIME-MD: limitations in diagnosing mental disorders in primary care. <i>European Journal of Public Health</i> , 2009 , 19, 303-7	2.1	43
211	Patient factors associated with guideline-concordant treatment of anxiety and depression in primary care. <i>Journal of General Internal Medicine</i> , 2010 , 25, 648-55	4	43
210	Pain as a risk factor for common mental disorders. Results from the Netherlands Mental Health Survey and Incidence Study-2: a longitudinal, population-based study. <i>Pain</i> , 2018 , 159, 712-718	8	42
209	Depression, disability and somatic diseases among elderly. <i>Journal of Affective Disorders</i> , 2014 , 167, 187-96	6.6	42
208	The incidence of depression and its risk factors in Dutch nursing homes and residential care homes. <i>American Journal of Geriatric Psychiatry</i> , 2012 , 20, 932-42	6.5	41
207	Effectiveness of collaborative stepped care for anxiety disorders in primary care: a pragmatic cluster randomised controlled trial. <i>Psychotherapy and Psychosomatics</i> , 2014 , 83, 37-44	9.4	40
206	The effects of integrative reminiscence on depressive symptomatology and mastery of older adults. <i>Community Mental Health Journal</i> , 2009 , 45, 476-84	2.1	40
205	Correlates of alcohol abstinence and at-risk alcohol consumption in older adults with depression: the NESDO study. <i>American Journal of Geriatric Psychiatry</i> , 2014 , 22, 866-74	6.5	37
204	Medically unexplained and explained physical symptoms in the general population: association with prevalent and incident mental disorders. <i>PLoS ONE</i> , 2015 , 10, e0123274	3.7	36
203	Collaborative care for anxiety disorders in primary care: a systematic review and meta-analysis. <i>BMC Family Practice</i> , 2016 , 17, 62	2.6	36
202	The effect of patient feedback on physicians' consultation skills: a systematic review. <i>Academic Medicine</i> , 2011 , 86, 1426-36	3.9	35
201	The association between somatization and disability in primary care patients. <i>Journal of Psychosomatic Research</i> , 2015 , 79, 117-22	4.1	34

200	Contradictory effects for prevention of depression and anxiety in residents in homes for the elderly: a pragmatic randomized controlled trial. <i>International Psychogeriatrics</i> , 2012 , 24, 1242-51	3.4	34
199	Sense of competence questionnaire among informal caregivers of older adults with dementia symptoms: a psychometric evaluation. <i>Clinical Practice and Epidemiology in Mental Health</i> , 2007 , 3, 11	3.2	34
198	Effectiveness of disease management programmes for recognition, diagnosis and treatment of depression in primary care. <i>European Journal of General Practice</i> , 2004 , 10, 6-12	2.8	34
197	Effects of Two-Year Vitamin B and Folic Acid Supplementation on Depressive Symptoms and Quality of Life in Older Adults with Elevated Homocysteine Concentrations: Additional Results from the B-PROOF Study, an RCT. <i>Nutrients</i> , 2016 , 8,	6.7	34
196	Antidepressant prescription for depression in general practice in The Netherlands. <i>International Journal of Clinical Pharmacy</i> , 2001 , 23, 46-9		33
195	Long-term antidepressant use: a qualitative study on perspectives of patients and GPs in primary care. <i>British Journal of General Practice</i> , 2016 , 66, e708-19	1.6	33
194	The impact of chronic somatic diseases on the course of depressive and anxiety disorders. <i>Psychotherapy and Psychosomatics</i> , 2013 , 82, 64-6	9.4	32
193	Cost-effectiveness of collaborative care including PST and an antidepressant treatment algorithm for the treatment of major depressive disorder in primary care; a randomised clinical trial. <i>BMC Health Services Research</i> , 2007 , 7, 34	2.9	32
192	De Vierdimensionale Klachtenlijst (4DKL) spoort psychische problemen op. <i>Huisarts En Wetenschap</i> , 2008 , 51, 251-255	0.1	32
191	Depression in older persons with versus without vascular disease in the open population: similar depressive symptom patterns, more disability. <i>Journal of Affective Disorders</i> , 2004 , 83, 155-60	6.6	32
190	Negative association of concomitant physical symptoms with the course of major depressive disorder: a systematic review. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 511-9	4.1	31
189	Prevalence of depressive symptoms and depressive disorder in primary care patients over 65 years of age. <i>Family Practice</i> , 1994 , 11, 80-4	1.9	31
188	"Being all alone makes me sad": loneliness in older adults with depressive symptoms. <i>International Psychogeriatrics</i> , 2014 , 1-11	3.4	30
187	Lithium augmentation in geriatric depression. <i>Journal of Affective Disorders</i> , 1990 , 20, 217-23	6.6	30
186	The European general practice research network presents the translations of its comprehensive definition of multimorbidity in family medicine in ten European languages. <i>PLoS ONE</i> , 2015 , 10, e0115796	3.7	30
185	Depressive and anxiety disorders and risk of subclinical atherosclerosis Findings from the Netherlands Study of Depression and Anxiety (NESDA). <i>Journal of Psychosomatic Research</i> , 2010 , 69, 203-10	4.1	29
184	Stepped care for depression and anxiety: from primary care to specialized mental health care: a randomised controlled trial testing the effectiveness of a stepped care program among primary care patients with mood or anxiety disorders. <i>BMC Health Services Research</i> , 2009 , 9, 90	2.9	29
183	Vitamin D supplementation for the prevention of depression and poor physical function in older persons: the D-Vitaal study, a randomized clinical trial. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 1119-1130	7	28

182	Sensitivity to depression or anxiety and subclinical cardiovascular disease. <i>Journal of Affective Disorders</i> , 2013 , 146, 126-31	6.6	28
181	Clinical effectiveness of usual care with or without antidepressant medication for primary care patients with minor or mild-major depression: a randomized equivalence trial. <i>BMC Medicine</i> , 2007 , 5, 36	11.4	28
180	The Three-Month Effect of Mobile Internet-Based Cognitive Therapy on the Course of Depressive Symptoms in Remitted Recurrently Depressed Patients: Results of a Randomized Controlled Trial. <i>Psychotherapy and Psychosomatics</i> , 2015 , 84, 90-99	9.4	27
179	Cost utility analysis of a collaborative stepped care intervention for panic and generalized anxiety disorders in primary care. <i>Journal of Psychosomatic Research</i> , 2014 , 77, 57-63	4.1	27
178	The UPBEAT nurse-delivered personalized care intervention for people with coronary heart disease who report current chest pain and depression: a randomised controlled pilot study. <i>PLoS ONE</i> , 2014 , 9, e98704	3.7	27
177	Identifying target groups for the prevention of depression among caregivers of dementia patients. <i>International Psychogeriatrics</i> , 2012 , 24, 298-306	3.4	27
176	Geographical epidemiology of health and overall deprivation in England, its changes and persistence from 2004 to 2015: a longitudinal spatial population study. <i>Journal of Epidemiology and Community Health</i> , 2018 , 72, 140-147	5.1	26
175	Determinants of (non-)recognition of depression by general practitioners: results of the Netherlands Study of Depression and Anxiety. <i>Journal of Affective Disorders</i> , 2012 , 138, 397-404	6.6	26
174	Multidisciplinary collaborative care for depressive disorder in the occupational health setting: design of a randomised controlled trial and cost-effectiveness study. <i>BMC Health Services Research</i> , 2008 , 8, 99	2.9	26
173	Cost-effectiveness of collaborative care for chronically ill patients with comorbid depressive disorder in the general hospital setting, a randomised controlled trial. <i>BMC Health Services Research</i> , 2007 , 7, 28	2.9	26
172	Cost-effectiveness of a disease management program for major depression in elderly primary care patients. <i>Journal of General Internal Medicine</i> , 2006 , 21, 1020-6	4	26
171	Does patient feedback improve the consultation skills of general practice trainees? A controlled trial. <i>Medical Education</i> , 2010 , 44, 156-64	3.7	25
170	Carotid atherosclerosis in depression and anxiety: associations for age of depression onset. <i>World Journal of Biological Psychiatry</i> , 2011 , 12, 549-58	3.8	25
169	(Cost)-effectiveness of family meetings on indicated prevention of anxiety and depressive symptoms and disorders of primary family caregivers of patients with dementia: design of a randomized controlled trial. <i>BMC Geriatrics</i> , 2008 , 8, 2	4.1	25
168	(Cost)-effectiveness of case-management by district nurses among primary informal caregivers of older adults with dementia symptoms and the older adults who receive informal care: design of a randomized controlled trial [ISRCTN83135728]. <i>BMC Public Health</i> , 2005 , 5, 133	4.1	25
167	Which DSM validated tools for diagnosing depression are usable in primary care research? A systematic literature review. <i>European Psychiatry</i> , 2017 , 39, 99-105	6	24
166	Illness burden and physical outcomes associated with collaborative care in patients with comorbid depressive disorder in chronic medical conditions: A systematic review and meta-analysis. <i>General Hospital Psychiatry</i> , 2018 , 50, 1-14	5.6	24
165	The association between medically unexplained physical symptoms and health care use over two years and the influence of depressive and anxiety disorders and personality traits: a longitudinal study. <i>BMC Health Services Research</i> , 2016 , 16, 100	2.9	24

164	The associations between personality characteristics and absenteeism: a cross-sectional study in workers with and without depressive and anxiety disorders. <i>Journal of Occupational Rehabilitation</i> , 2013 , 23, 309-17	3.6	24
163	Prevention of anxiety and depression in the age group of 75 years and over: a randomised controlled trial testing the feasibility and effectiveness of a generic stepped care programme among elderly community residents at high risk of developing anxiety and depression versus usual care [ISRCTN126474556]. <i>BMC Public Health</i> , 2006 , 6, 186	4.1	24
162	Exploring dementia management attitudes in primary care: a key informant survey to primary care physicians in 25 European countries. <i>International Psychogeriatrics</i> , 2017 , 29, 1413-1423	3.4	23
161	Vitamin D, PTH and the risk of overall and disease-specific mortality: Results of the Longitudinal Aging Study Amsterdam. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2016 , 164, 386-394	5.1	23
160	Psychological treatment of anxiety in primary care: a meta-analysis. <i>Psychological Medicine</i> , 2013 , 43, 351-61	6.9	23
159	How effective is bibliotherapy for very old adults with subthreshold depression? A randomized controlled trial. <i>American Journal of Geriatric Psychiatry</i> , 2011 , 19, 256-65	6.5	23
158	Disrupting the rhythm of depression using Mobile Cognitive Therapy for recurrent depression: randomized controlled trial design and protocol. <i>BMC Psychiatry</i> , 2011 , 11, 12	4.2	23
157	Adverse effects of multiple physical symptoms on the course of depressive and anxiety symptoms in primary care. <i>Psychotherapy and Psychosomatics</i> , 2010 , 79, 389-91	9.4	23
156	Collaborative stepped care for anxiety disorders in primary care: aims and design of a randomized controlled trial. <i>BMC Health Services Research</i> , 2009 , 9, 159	2.9	23
155	The guideline "consultation psychiatry" of the Netherlands Psychiatric Association. <i>Journal of Psychosomatic Research</i> , 2009 , 66, 531-5	4.1	23
154	Primary care patients with mental health problems: outcome of a randomised clinical trial. <i>British Journal of General Practice</i> , 2007 , 57, 886-91	1.6	23
153	Depression increases the onset of cardiovascular disease over and above other determinants in older primary care patients, a cohort study. <i>BMC Cardiovascular Disorders</i> , 2015 , 15, 40	2.3	22
152	Cost-effectiveness of a stepped-care intervention to prevent major depression in patients with type 2 diabetes mellitus and/or coronary heart disease and subthreshold depression: design of a cluster-randomized controlled trial. <i>BMC Psychiatry</i> , 2013 , 13, 128	4.2	22
151	Primary care management of major depression in patients aged > or =55 years: outcome of a randomised clinical trial. <i>British Journal of General Practice</i> , 2008 , 58, 680-6, I-II; discussion 687	1.6	22
150	Serious adverse events reported in placebo randomised controlled trials of oral naltrexone: a systematic review and meta-analysis. <i>BMC Medicine</i> , 2019 , 17, 10	11.4	22
149	The cost-effectiveness of a family meetings intervention to prevent depression and anxiety in family caregivers of patients with dementia: a randomized trial. <i>Trials</i> , 2013 , 14, 305	2.8	21
148	Vitamin D supplementation to prevent depression and poor physical function in older adults: Study protocol of the D-Vitaal study, a randomized placebo-controlled clinical trial. <i>BMC Geriatrics</i> , 2015 , 15, 151	4.1	21
147	Cost-effectiveness of a stepped care programme to prevent depression and anxiety in residents in homes for the older people: a randomised controlled trial. <i>International Journal of Geriatric Psychiatry</i> , 2014 , 29, 182-90	3.9	21

146	Effectiveness of guided self-help for depression and anxiety disorders in primary care: a pragmatic randomized controlled trial. <i>Psychiatry Research</i> , 2011 , 187, 113-20	9.9	21
145	Outcomes for depression and anxiety in primary care and details of treatment: a naturalistic longitudinal study. <i>BMC Psychiatry</i> , 2011 , 11, 180	4.2	21
144	Self-reports on the IQCODE in older adults: a psychometric evaluation. <i>Journal of Geriatric Psychiatry and Neurology</i> , 2008 , 21, 83-92	3.8	21
143	Measurement properties of depression questionnaires in patients with diabetes: a systematic review. <i>Quality of Life Research</i> , 2018 , 27, 1415-1430	3.7	20
142	Systematic review of measurement properties of questionnaires measuring somatization in primary care patients. <i>Journal of Psychosomatic Research</i> , 2017 , 103, 42-62	4.1	20
141	To what extent does the anxiety scale of the Four-Dimensional Symptom Questionnaire (4DSQ) detect specific types of anxiety disorder in primary care? A psychometric study. <i>BMC Psychiatry</i> , 2014 , 14, 121	4.2	19
140	Recognition of anxiety disorders by family physicians after rigorous medical record case extraction: results of the Netherlands Study of Depression and Anxiety. <i>General Hospital Psychiatry</i> , 2012 , 34, 460-7	5.6	19
139	Diagnose depressie niet zo betrouwbaar te stellen. <i>Huisarts En Wetenschap</i> , 2004 , 47, 894-899	0.1	19
138	Diagnostic accuracy of the Patient Health Questionnaire-9 for assessment of depression in type II diabetes mellitus and/or coronary heart disease in primary care. <i>Journal of Affective Disorders</i> , 2016 , 190, 68-74	6.6	18
137	Accelerometer- and Pedometer-Based Physical Activity Interventions Among Adults With Cardiometabolic Conditions: A Systematic Review and Meta-analysis. <i>JAMA Network Open</i> , 2019 , 2, e1912895	10.4	18
136	Cost-effectiveness of a minimal intervention for stress-related sick leave in general practice: results of an economic evaluation alongside a pragmatic randomised control trial. <i>Journal of Affective Disorders</i> , 2010 , 120, 177-87	6.6	18
135	A postmarketing study of flunarizine in migraine and vertigo. <i>International Journal of Clinical Pharmacy</i> , 1997 , 19, 269-74		17
134	Effectiveness of a Minimal Intervention for Stress-related mental disorders with Sick leave (MISS); study protocol of a cluster randomised controlled trial in general practice [ISRCTN43779641]. <i>BMC Public Health</i> , 2006 , 6, 124	4.1	17
133	Utility scores for different health states related to depression: individual participant data analysis. <i>Quality of Life Research</i> , 2017 , 26, 1649-1658	3.7	16
132	A high physical symptom count reduces the effectiveness of treatment for depression, independently of chronic medical conditions. <i>Journal of Psychosomatic Research</i> , 2013 , 74, 179-85	4.1	16
131	Effectiveness of Supported Self-Help in Recurrent Depression: A Randomized Controlled Trial in Primary Care. <i>Psychotherapy and Psychosomatics</i> , 2017 , 86, 220-230	9.4	16
130	High incidence of clinically relevant depressive symptoms in vulnerable persons of 75 years or older living in the community. <i>Aging and Mental Health</i> , 2010 , 14, 828-33	3.5	16
129	Validity aspects of the patient feedback questionnaire on consultation skills (PFC), a promising learning instrument in medical education. <i>Patient Education and Counseling</i> , 2009 , 76, 202-6	3.1	16

128	Effectiveness of family meetings for family caregivers on delaying time to nursing home placement of dementia patients: a randomized trial. <i>PLoS ONE</i> , 2012 , 7, e42145	3.7	16
127	Selective serotonin reuptake inhibitor and selective serotonin and norepinephrine reuptake inhibitor use and risk of fractures in adults: A systematic review and meta-analysis. <i>International Journal of Geriatric Psychiatry</i> , 2018 , 33, 1688-1708	3.9	16
126	Reliability of consultation skills assessments using standardised versus real patients. <i>Medical Education</i> , 2011 , 45, 578-84	3.7	15
125	Frequent attenders in general practice: problem solving treatment provided by nurses [ISRCTN51021015]. <i>BMC Family Practice</i> , 2005 , 6, 42	2.6	15
124	Effectiveness of a stepped-care intervention to prevent major depression in patients with type 2 diabetes mellitus and/or coronary heart disease and subthreshold depression: A pragmatic cluster randomized controlled trial. <i>PLoS ONE</i> , 2017 , 12, e0181023	3.7	15
123	Identifying undetected dementia in UK primary care patients: a retrospective case-control study comparing machine-learning and standard epidemiological approaches. <i>BMC Medical Informatics and Decision Making</i> , 2019 , 19, 248	3.6	15
122	Chronic morbidity, deprivation and primary medical care spending in England in 2015-16: a cross-sectional spatial analysis. <i>BMC Medicine</i> , 2018 , 16, 19	11.4	14
121	A research group from the European General Practice Research Network (EGPRN) explores the concept of multimorbidity for further research into long term care. <i>Journal of the American Medical Directors Association</i> , 2013 , 14, 132-3	5.9	14
120	Feasibility and effectiveness of activity-scheduling as a guided self-help intervention for the prevention of depression and anxiety in residents in homes for the elderly: a pragmatic randomized controlled trial. <i>International Psychogeriatrics</i> , 2011 , 23, 969-78	3.4	14
119	Training GP's to use a minimal intervention for stress-related mental disorders with sick leave (MISS): Effects on performance: Results of the MISS project; a cluster-randomised controlled trial [ISRCTN43779641]. <i>Patient Education and Counseling</i> , 2010 , 78, 206-11	3.1	14
118	Design and pilot results of a single blind randomized controlled trial of systematic demand-led home visits by nurses to frail elderly persons in primary care [ISRCTN05358495]. <i>BMC Geriatrics</i> , 2005 , 5, 11	4.1	14
117	Interpersonal psychotherapy (IPT) for late-life depression in general practice: uptake and satisfaction by patients, therapists and physicians. <i>BMC Family Practice</i> , 2007 , 8, 52	2.6	13
116	Depression and anxiety, an indicated prevention (DIP) protocol in homes for the elderly: feasibility and (cost) effectiveness of a stepped care programme. <i>BMC Geriatrics</i> , 2007 , 7, 6	4.1	13
115	Cost-effectiveness of interpersonal psychotherapy for elderly primary care patients with major depression. <i>International Journal of Technology Assessment in Health Care</i> , 2007 , 23, 480-7	1.8	13
114	Self-report depression scales for elderly patients in primary care: a preliminary study. <i>Family Practice</i> , 1993 , 10, 63-5	1.9	13
113	Cost-utility of collaborative care for major depressive disorder in primary care in the Netherlands. <i>Journal of Psychosomatic Research</i> , 2015 , 79, 316-23	4.1	12
112	Determinants of receiving mental health care for depression in older adults. <i>Journal of Affective Disorders</i> , 2012 , 143, 69-74	6.6	12
111	Cost-effectiveness of problem-solving treatment in comparison with usual care for primary care patients with mental health problems: a randomized trial. <i>BMC Family Practice</i> , 2012 , 13, 98	2.6	12

110	Depression in Dutch homes for the elderly: under-diagnosis in demented residents?. <i>International Journal of Geriatric Psychiatry</i> , 2010 , 25, 712-8	3.9	12
109	Alprazolam for depression. <i>The Cochrane Library</i> , 2012 , CD007139	5.2	12
108	Development and feasibility of a patient feedback programme to improve consultation skills in general practice training. <i>Patient Education and Counseling</i> , 2008 , 72, 12-9	3.1	12
107	Reliability and validity of the assessment of depression in general practice: the Short Depression Interview (SDI). <i>General Hospital Psychiatry</i> , 2002 , 24, 396-405	5.6	12
106	Diagnostic accuracy of depression questionnaires in adult patients with diabetes: A systematic review and meta-analysis. <i>PLoS ONE</i> , 2019 , 14, e0218512	3.7	11
105	Somatisation as a risk factor for incident depression and anxiety. <i>Journal of Psychosomatic Research</i> , 2015 , 79, 614-9	4.1	11
104	Low-dose naltrexone as a treatment for chronic fatigue syndrome. <i>BMJ Case Reports</i> , 2020 , 13,	0.9	11
103	The role of somatic health problems in the recognition of depressive and anxiety disorders by general practitioners. <i>Journal of Affective Disorders</i> , 2013 , 151, 1025-32	6.6	11
102	Longitudinal associations of multiple physical symptoms with recurrence of depressive and anxiety disorders. <i>Journal of Psychosomatic Research</i> , 2017 , 97, 96-101	4.1	10
101	An implementation science study to enhance cardiovascular disease prevention in Mukono and Buikwe districts in Uganda: a stepped-wedge design. <i>BMC Health Services Research</i> , 2019 , 19, 253	2.9	10
100	Does an outreaching stepped care program reduce depressive symptoms in community-dwelling older adults? A randomized implementation trial. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 807-17	6.5	10
99	Predictors of Societal Costs in Dementia Patients and Their Informal Caregivers: A Two-Year Prospective Cohort Study. <i>American Journal of Geriatric Psychiatry</i> , 2015 , 23, 1193-203	6.5	10
98	Management of patients with persistent medically unexplained symptoms: a descriptive study. <i>BMC Family Practice</i> , 2018 , 19, 88	2.6	10
97	One consensual depression diagnosis tool to serve many countries: a challenge! A RAND/UCLA methodology. <i>BMC Research Notes</i> , 2018 , 11, 4	2.3	10
96	Enhanced care for depression. <i>Current Opinion in Psychiatry</i> , 2013 , 26, 7-12	4.9	10
95	Yield of a new method to detect cognitive impairment in general practice. <i>International Journal of Geriatric Psychiatry</i> , 2007 , 22, 590-7	3.9	10
94	Case Finding of Mild Cognitive Impairment and Dementia and Subsequent Care; Results of a Cluster RCT in Primary Care. <i>PLoS ONE</i> , 2016 , 11, e0156958	3.7	10
93	Predictors of Persistent Medically Unexplained Physical Symptoms: Findings From a General Population Study. <i>Frontiers in Psychiatry</i> , 2018 , 9, 613	5	10

92	Behavioural activation by mental health nurses for late-life depression in primary care: a randomized controlled trial. <i>BMC Psychiatry</i> , 2017 , 17, 230	4.2	9
91	What research agenda could be generated from the European General Practice Research Network concept of Multimorbidity in Family Practice?. <i>BMC Family Practice</i> , 2015 , 16, 125	2.6	9
90	Prevention of anxiety disorders in primary care: a feasibility study. <i>BMC Psychiatry</i> , 2012 , 12, 206	4.2	9
89	Cost-effectiveness of nurse-led self-help for recurrent depression in the primary care setting: design of a pragmatic randomised controlled trial. <i>BMC Psychiatry</i> , 2012 , 12, 59	4.2	9
88	Osteoarthritis pain assessment in family practice. <i>Arthritis and Rheumatism</i> , 1994 , 7, 40-5		9
87	Using electronic health records to quantify and stratify the severity of type 2 diabetes in primary care in England: rationale and cohort study design. <i>BMJ Open</i> , 2018 , 8, e020926	3	9
86	Dutch family physicians' awareness of cognitive impairment among the elderly. <i>BMC Geriatrics</i> , 2015 , 15, 105	4.1	8
85	Process evaluation of a stepped-care program to prevent depression in primary care: patients' and practice nurses' experiences. <i>BMC Family Practice</i> , 2017 , 18, 26	2.6	8
84	Case-finding of dementia in general practice and effects of subsequent collaborative care; design of a cluster RCT. <i>BMC Public Health</i> , 2012 , 12, 609	4.1	8
83	The costs of guideline-concordant care and of care according to patients' needs in anxiety and depression. <i>Journal of Evaluation in Clinical Practice</i> , 2011 , 17, 537-46	2.5	8
82	A randomized controlled trial to improve the recognition, diagnosis and treatment of major depression in elderly people in general practice: design, first results and feasibility of the West Friesland Study. <i>Primary Care and Community Psychiatry</i> , 2003 , 8, 135-140		8
81	Chest pain, depression and anxiety in coronary heart disease: Consequence or cause? A prospective clinical study in primary care. <i>Journal of Psychosomatic Research</i> , 2020 , 129, 109891	4.1	8
80	Mental health care use in medically unexplained and explained physical symptoms: findings from a general population study. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 2063-72	3.1	8
79	Patients' and practice nurses' perceptions of depression in patients with type 2 diabetes and/or coronary heart disease screened for subthreshold depression. <i>BMC Family Practice</i> , 2018 , 19, 202	2.6	8
78	The French version of the HSCL-25 has now been validated for use in primary care. <i>PLoS ONE</i> , 2019 , 14, e0214804	3.7	7
77	Cardiovascular risk factor mapping and distribution among adults in Mukono and Buikwe districts in Uganda: small area analysis. <i>BMC Cardiovascular Disorders</i> , 2020 , 20, 284	2.3	7
76	Effectiveness and cost-effectiveness of transmural collaborative care with consultation letter (TCCCL) and duloxetine for major depressive disorder (MDD) and (sub)chronic pain in collaboration with primary care: design of a randomized placebo-controlled multi-Centre trial: TCC:PAINDIP. <i>BMC Psychiatry</i> , 2013 , 13, 147	4.2	7
75	The prognosis of depressive disorder in elderly primary care patients. An exploratory observational study: course at 6 and 12 months. <i>Scandinavian Journal of Primary Health Care</i> , 1998 , 16, 107-11	2.7	7

74	Interventions Using Wearable Physical Activity Trackers Among Adults With Cardiometabolic Conditions: A Systematic Review and Meta-analysis. <i>JAMA Network Open</i> , 2021 , 4, e2116382	10.4	7
73	How do general practitioners recognize the definition of multimorbidity? A European qualitative study. <i>European Journal of General Practice</i> , 2016 , 22, 159-68	2.8	6
72	Assessing the Impact of EQ-5D Country-specific Value Sets on Cost-utility Outcomes. <i>Medical Care</i> , 2021 , 59, 82-90	3.1	6
71	Vitamin D Status and Depressive Symptoms in Older Adults: A Role for Physical Functioning?. <i>American Journal of Geriatric Psychiatry</i> , 2018 , 26, 1131-1143	6.5	5
70	Non-fatal disease burden for subtypes of depressive disorder: population-based epidemiological study. <i>BMC Psychiatry</i> , 2016 , 16, 139	4.2	5
69	Implementing an outreaching, preference-led stepped care intervention programme to reduce late life depressive symptoms: results of a mixed-methods study. <i>Implementation Science</i> , 2014 , 9, 107	8.4	5
68	Preventing depression in homes for older adults: are effects sustained over 2 years?. <i>International Journal of Geriatric Psychiatry</i> , 2014 , 29, 191-7	3.9	5
67	Waarde van de Vierdimensionale Klachtenlijst (4DKL) voor het detecteren van depressieve stoornissen. <i>Huisarts En Wetenschap</i> , 2007 , 50, 494-501	0.1	5
66	How to improve mental health competency in general practice training?--a SWOT analysis. <i>European Journal of General Practice</i> , 2004 , 10, 61-5	2.8	5
65	Management of depression in elderly general practice patients. <i>Scandinavian Journal of Primary Health Care</i> , 1994 , 12, 162-8	2.7	5
64	Barriers and facilitators to the adoption of electronic clinical decision support systems: a qualitative interview study with UK general practitioners. <i>BMC Medical Informatics and Decision Making</i> , 2021 , 21, 193	3.6	5
63	Pain as a risk factor for suicidal ideation. A population-based longitudinal cohort study. <i>General Hospital Psychiatry</i> , 2020 , 63, 54-61	5.6	5
62	Pulling out all the stops: what motivates 65+ year olds with depressive symptoms to participate in an outreaching preference-led intervention programme?. <i>Aging and Mental Health</i> , 2015 , 19, 453-63	3.5	4
61	The CIPRUS study, a nurse-led psychological treatment for patients with undifferentiated somatoform disorder in primary care: study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 206	2.8	4
60	Cross-cultural validation of the Turkish Four-Dimensional Symptom Questionnaire (4DSQ) using differential item and test functioning (DIF and DTF) analysis. <i>BMC Family Practice</i> , 2016 , 17, 53	2.6	4
59	Screening high-risk patients and assisting in diagnosing anxiety in primary care: the Patient Health Questionnaire evaluated. <i>BMC Psychiatry</i> , 2013 , 13, 192	4.2	4
58	Mean clinical challenge rate and level of recognition of depression remain unchanged after two years of vocational training. <i>Family Practice</i> , 2001 , 18, 590-1	1.9	4
57	Identification and management of frailty in English primary care: a qualitative study of national policy. <i>BJGP Open</i> , 2020 , 4,	3.1	4

56	Unburdening dementia - a basic social process grounded theory based on a primary care physician survey from 25 countries. <i>Scandinavian Journal of Primary Health Care</i> , 2020 , 38, 253-264	2.7	4
55	What is meant by "frailty" in undergraduate medical education? A national survey of UK medical schools. <i>European Geriatric Medicine</i> , 2021 , 12, 355-362	3	4
54	Assessing the severity of Type 2 diabetes using clinical data-based measures: a systematic review. <i>Diabetic Medicine</i> , 2019 , 36, 688-701	3.5	3
53	Increasing socioeconomic gap between the young and old: temporal trends in health and overall deprivation in England by age, sex, urbanity and ethnicity, 2004-2015. <i>Journal of Epidemiology and Community Health</i> , 2018 , 72, 636-644	5.1	3
52	Comparative Effect of Collaborative Care, Pain Medication, and Duloxetine in the Treatment of Major Depressive Disorder and Comorbid (Sub)Chronic Pain: Results of an Exploratory Randomized, Placebo-Controlled, Multicenter Trial (CC:PAINDIP). <i>Frontiers in Psychiatry</i> , 2018 , 9, 118	5	3
51	Wat heeft vijf jaar kwaliteitsbeleid ggz voor de huisarts opgeleverd?. <i>Huisarts En Wetenschap</i> , 2006 , 49, 365-369	0.1	3
50	Overuse of antipsychotic medication in elderly people with dementia? A view from general practice. <i>Mental Health in Family Medicine</i> , 2009 , 6, 191-3		3
49	Understanding barriers to women seeking and receiving help for perinatal mental health problems in UK general practice: development of a questionnaire. <i>Primary Health Care Research and Development</i> , 2019 , 20, e156	1.6	3
48	Two-year effectiveness of a stepped-care depression prevention intervention and predictors of incident depression in primary care patients with diabetes type 2 and/or coronary heart disease and subthreshold depression: data from the Step-Dep cluster randomised controlled trial. <i>BMJ Open</i> , 2018 , 8, e020412	3	3
47	Notice of Retraction. Hodkinson et al. Accelerometer- and Pedometer-Based Physical Activity Interventions Among Adults With Cardiometabolic Conditions: A Systematic Review and Meta-analysis. <i>JAMA Netw Open</i> . 2019;2(10):e1912895. <i>JAMA Network Open</i> , 2020 , 3, e2032700	10.4	3
46	Development and validation of the Diabetes Severity SCOrE (DISSCO) in 139 626 individuals with type 2 diabetes: a retrospective cohort study. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	2
45	A protocol paper: community engagement interventions for cardiovascular disease prevention in socially disadvantaged populations in the UK: an implementation research study. <i>Global Health Research and Policy</i> , 2020 , 5, 12	7.6	2
44	Using patient and public involvement to improve the research design and funding application for a project aimed at fostering a more collaborative approach to the NHS health check: the CaVIAR project (better Care Via Improved Access to Records). <i>Research Involvement and Engagement</i> , 2018 , 4, 18	4.4	2
43	Effectiveness of a cognitive behavioural intervention for patients with undifferentiated somatoform disorder: Results from the CIPRUS cluster randomized controlled trial in primary care. <i>Journal of Psychosomatic Research</i> , 2019 , 127, 109745	4.1	2
42	Collaborative care voor depressieve patiënten. <i>Huisarts En Wetenschap</i> , 2013 , 56, 210-213	0.1	2
41	PMH39 Cost Effectiveness of A Collaboratieve Care Stepped intervention for Anxiety Disorders in the Primary Care Setting. <i>Value in Health</i> , 2011 , 14, A293	3.3	2
40	Betere detectie van dementie. <i>Huisarts En Wetenschap</i> , 2008 , 51, 115-119	0.1	2
39	Could dementia be detected from UK primary care patients' records by simple automated methods earlier than by the treating physician? A retrospective case-control study. <i>Wellcome Open Research</i> , 2020 , 5, 120	4.8	2

38	Prevention of dementia using mobile phone applications (PRODEMOS): protocol for an international randomised controlled trial. <i>BMJ Open</i> , 2021 , 11, e049762	3	2
37	A supported self-help for recurrent depression in primary care; An economic evaluation alongside a multi-center randomised controlled trial. <i>PLoS ONE</i> , 2018 , 13, e0208570	3.7	2
36	Cross-Cultural Validation of the Definition of Multimorbidity in the Bulgarian Language. <i>Folia Medica</i> , 2015 , 57, 127-32	0.5	1
35	Collaborative stepped care is effectief bij angststoornissen. <i>Huisarts En Wetenschap</i> , 2014 , 57, 566-570	0.1	1
34	Stapsgewijze preventie van angst en depressie bij ouderen. <i>Huisarts En Wetenschap</i> , 2009 , 52, 668-673	0.1	1
33	Commentary: Good primary mental health care is simple. <i>British Journal of General Practice</i> , 2008 , 58, 409	1.6	1
32	Marketing beïnvloedt huisartsen te veel Exit TCAE?. <i>Huisarts En Wetenschap</i> , 2003 , 46, 802-803	0.1	1
31	UPBEAT-UK: a programme of research into the relationship between coronary heart disease and depression in primary care patients. <i>Programme Grants for Applied Research</i> , 2016 , 4, 1-172	1.5	1
30	A brief cognitive behavioural intervention is cost-effective for primary care patients with medically unexplained physical symptoms compared to usual care. <i>Journal of Psychosomatic Research</i> , 2020 , 138, 110217	4.1	0
29	Supported self-help to prevent relapse or recurrence of depression: Who benefits most?. <i>Journal of Affective Disorders</i> , 2019 , 257, 180-186	6.6	0
28	Preventieve huisbezoeken bij kwetsbare ouderen. <i>Huisarts En Wetenschap</i> , 2011 , 54, 366-374	0.1	0
27	Feasibility of cognitive assessments depends on the population. <i>International Journal of Geriatric Psychiatry</i> , 2007 , 22, 1167-8	3.9	0
26	Nine Forward-Backward Translations of the Hopkins Symptom Checklist-25 With Cultural Checks. <i>Frontiers in Psychiatry</i> , 2021 , 12, 688154	5	0
25	Design and Development of a Mobile Health (mHealth) Platform for Dementia Prevention in the Prevention of Dementia by Mobile Phone Applications (PRODEMOS) Project.. <i>Frontiers in Neurology</i> , 2021 , 12, 733878	4.1	0
24	Cognitieve gedragsinterventie door de poh-ggz bij SOLK. <i>Huisarts En Wetenschap</i> , 2020 , 63, 31-38	0.1	
23	Recente ontwikkelingen in de psychotherapie. <i>Huisarts En Wetenschap</i> , 2014 , 57, 260-264	0.1	
22	Paniek aanval. <i>Huisarts En Wetenschap</i> , 2013 , 56, 247-247	0.1	
21	Preventie van depressie en angst in verzorgingshuizen. <i>Huisarts En Wetenschap</i> , 2013 , 56, 154-158	0.1	

20	Familiegesprekken voor mantelzorgers van mensen met dementie. <i>Huisarts En Wetenschap</i> , 2013 , 56, 160-164	0.1
19	Zelfhulp voor depressie en angststoornissen. <i>Huisarts En Wetenschap</i> , 2011 , 54, 488-490	0.1
18	The complex power of the gut. <i>European Journal of General Practice</i> , 2010 , 16, 70-1	2.8
17	Disease management voor depressieve patiënten van 55+. <i>Huisarts En Wetenschap</i> , 2009 , 52, 281-286	0.1
16	Study findings hard to interpret. <i>Archives of Internal Medicine</i> , 2012 , 172, 1429; author reply 1429-30	
15	Patient Feedback Via a National Registry Could Improve Physicians' CME. <i>Academic Medicine</i> , 2012 , 87, 549	3.9
14	Depressie en angst bij partners van dementiepatiënten. <i>Huisarts En Wetenschap</i> , 2010 , 53, 420-424	0.1
13	Verbetert feedback van patiënten de consultvaardigheden van huisartsen-in-opleiding?. <i>Huisarts En Wetenschap</i> , 2010 , 53, 556-561	0.1
12	Preventie van psychische problemen vanuit het perspectief van de huisartspraktijk. <i>TSG: Tijdschrift Voor Gezondheidswetenschappen</i> , 2007 , 85, 62-63	0.2
11	Interpersoonlijke psychotherapie voor depressieve ouderen. <i>Huisarts En Wetenschap</i> , 2007 , 50, 797-801	0.1
10	Helpen antidepressiva bij depressieve klachten of lichte depressieve stoornissen?. <i>Huisarts En Wetenschap</i> , 2008 , 51, 361-369	0.1
9	De impact van euthanasie op Nederlandse huisartsen. <i>Huisarts En Wetenschap</i> , 2008 , 51, 588-593	0.1
8	De toepasbaarheid van interpersoonlijke psychotherapie bij depressieve ouderen. <i>Dth</i> , 2003 , 23, 158-164	
7	Sombere stemming. <i>Huisarts En Wetenschap</i> , 2003 , 46, 444-448	0.1
6	Prevention of dementia using mobile phone applications (PRODEMOS): A randomized controlled trial in progress. <i>Alzheimer's and Dementia</i> , 2020 , 16, e043039	1.2
5	Zelfhulp om terugval in een depressie te voorkomen. <i>Huisarts En Wetenschap</i> , 2018 , 61, 20-23	0.1
4	Identification and management of frail patients in English primary care: an analysis of the General Medical Services 2018/2019 contract dataset. <i>BMJ Open</i> , 2021 , 11, e041091	3
3	Validiteit van de 4DKL bij mensen met een migratieachtergrond. <i>Huisarts En Wetenschap</i> , 2021 , 64, 33-36	0.1

- 2 'Antidepressants unlimited'. Are antidepressants the best option for treating depression in general practice? Debating session WONCA 2004. *European Journal of General Practice*, **2005**, 11, 119-21 2.8
- 1 Understanding the implementation of interventions to improve the management of frailty in primary care: a rapid realist review. *BMJ Open*, **2022**, 12, e054780 3