
Hua Ren

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5780841/publications.pdf Version: 2024-02-01



HUA DEN

#	Article	IF	CITATIONS
1	Plateletâ€toâ€lymphocyte ratio is associated with prognosis in patients with coronavirus diseaseâ€19. Journal of Medical Virology, 2020, 92, 1533-1541.	2.5	418
2	Hypofractionated versus conventional fractionated postmastectomy radiotherapy for patients with high-risk breast cancer: a randomised, non-inferiority, open-label, phase 3 trial. Lancet Oncology, The, 2019, 20, 352-360.	5.1	258
3	Switching on prodrugs using radiotherapy. Nature Chemistry, 2021, 13, 805-810.	6.6	91
4	Hypofractionated Versus Conventional Fractionated Radiotherapy After Breast-Conserving Surgery in the Modern Treatment Era: A Multicenter, Randomized Controlled Trial From China. Journal of Clinical Oncology, 2020, 38, 3604-3614.	0.8	58
5	Mapping Patterns of Ipsilateral Supraclavicular Nodal Metastases in Breast Cancer: Rethinking the Clinical Target Volume for High-risk Patients. International Journal of Radiation Oncology Biology Physics, 2015, 93, 268-276.	0.4	51
6	Circulating serum microRNA-345 correlates with unfavorable pathological response to preoperative chemoradiotherapy in locally advanced rectal cancer. Oncotarget, 2016, 7, 64233-64243.	0.8	39
7	Radiation-Induced Lymphopenia Predicts Poorer Prognosis in Patients With Breast Cancer: A Post Hoc Analysis of a Randomized Controlled Trial of Postmastectomy Hypofractionated Radiation Therapy. International Journal of Radiation Oncology Biology Physics, 2020, 108, 277-285.	0.4	33
8	Comparison of Treatment Outcomes With Breast-conserving Surgery Plus Radiotherapy Versus Mastectomy for Patients With Stage I Breast Cancer: A Propensity Score-matched Analysis. Clinical Breast Cancer, 2018, 18, e975-e984.	1.1	21
9	Patterns of Primary Tumor Invasion and Regional Lymph Node Spread Based on Magnetic Resonance Imaging in Early-Stage Nasal NK/T-cell Lymphoma: Implications for Clinical Target Volume Definition and Prognostic Significance. International Journal of Radiation Oncology Biology Physics, 2017, 97, 50-59.	0.4	19
10	Phaseâ€lâ€study of postoperative radiotherapy combined with capecitabine for gastric cancer. World Journal of Gastroenterology, 2014, 20, 1067.	1.4	10
11	Nomogram predicting survival as a selection criterion for postmastectomy radiotherapy in patients with T1 to T2 breast cancer with 1 to 3 positive lymph nodes. Cancer, 2020, 126, 3857-3866.	2.0	10
12	Chidamide and Radiotherapy Synergistically Induce Cell Apoptosis and Suppress Tumor Growth and Cancer Stemness by Regulating the MiR-375-EIF4G3 Axis in Lung Squamous Cell Carcinomas. Journal of Oncology, 2021, 2021, 1-15.	0.6	10
13	Interim analysis of postoperative chemoradiotherapy with capecitabine and oxaliplatin versus capecitabine alone for pathological stage II and III rectal cancer: a randomized multicenter phase III trial. Oncotarget, 2016, 7, 25576-25584.	0.8	10
14	Intensity Modulated Radiation Therapy for Early-Stage Primary Gastric Diffuse Large B-Cell Lymphoma: Dosimetric Analysis, Clinical Outcome, and Quality of Life. International Journal of Radiation Oncology Biology Physics, 2016, 95, 712-720.	0.4	8
15	Trastuzumab Provides a Comparable Prognosis in Patients With HER2-Positive Breast Cancer to Those With HER2-Negative Breast Cancer: Post Hoc Analyses of a Randomized Controlled Trial of Post-Mastectomy Hypofractionated Radiotherapy. Frontiers in Oncology, 2020, 10, 605750.	1.3	7
16	PD-1 restrains IL-17A production from γĨ´T cells to modulate acute radiation-induced lung injury. Translational Lung Cancer Research, 2021, 10, 685-698.	1.3	7
17	Dosimetric and Clinical Outcomes With Intensity Modulated Radiation Therapy After Chemotherapy for Patients With Early-Stage Diffuse Large B-cell Lymphoma of Waldeyer Ring. International Journal of Radiation Oncology Biology Physics, 2016, 96, 379-386.	0.4	6
18	Radiotherapy plays an important role in improving the survival outcome in patients with T1–2N1M0 breast cancer – a joint analysis of 4262 real world cases from two institutions. BMC Cancer, 2020, 20, 1155.	1.1	6

Hua Ren

#	Article	IF	CITATIONS
19	ATF2-Induced Overexpression of IncRNA LINC00882, as a Novel Therapeutic Target, Accelerates Hepatocellular Carcinoma Progression via Sponging miR-214-3p to Upregulate CENPM. Frontiers in Oncology, 2021, 11, 714264.	1.3	6
20	Down-staging depth score to predict outcomes in locally advanced rectal cancer achieving ypl stage after neoadjuvant chemo-radiotherapy versus de novo stage pl cohort: A propensity score-matched analysis. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2018, 30, 373-381.	0.7	6
21	Locoregional recurrence patterns in women with breast cancer who have not undergone post-mastectomy radiotherapy. Radiation Oncology, 2020, 15, 212.	1.2	4
22	Prognosis and Prophylactic Regional Nodal Irradiation in Breast Cancer Patients With the First Isolated Chest Wall Recurrence After Mastectomy. Frontiers in Oncology, 2020, 10, 600525.	1.3	4
23	Adjuvant treatment may benefit patients with high-risk upper rectal cancer: A nomogram and recursive partitioning analysis of 547 patients. Oncotarget, 2016, 7, 66160-66169.	0.8	4
24	Tomotherapy as an adjuvant treatment for gastroesophageal junction and stomach cancer may reduce bowel and bone marrow toxicity compared to intensity-modulated radiotherapy and volumetric-modulated arc therapy. Oncotarget, 2017, 8, 39727-39735.	0.8	4
25	Postoperative Capecitabine with Concurrent Intensity-Modulated Radiotherapy or Three-Dimensional Conformal Radiotherapy for Patients with Stage II and III Rectal Cancer. PLoS ONE, 2015, 10, e0124601.	1.1	3
26	Associations of Genetic Variations in MicroRNA Seed Regions With Acute Adverse Events and Survival in Patients With Rectal Cancer Receiving Postoperative Chemoradiation Therapy. International Journal of Radiation Oncology Biology Physics, 2018, 100, 1026-1033.	0.4	3
27	Survey on the Use of Radiotherapy to Treat Early Breast Cancer following Breast-conserving Surgery in China. Tumori, 2014, 100, 512-517.	0.6	2
28	Timing of Chemotherapy and Radiotherapy Following Breast-Conserving Surgery for Early-Stage Breast Cancer: A Retrospective Analysis. Frontiers in Oncology, 2020, 10, 571390.	1.3	2
29	Postoperative Chemoradiotherapy With Capecitabine and Oxaliplatin vs Capecitabine for Stage II to III Rectal Cancer. JAMA Network Open, 2021, 4, e2136116.	2.8	2
30	Surgical treatment of primary tracheobronchial malignant tumors. Chinese-German Journal of Clinical Oncology, 2010, 9, 97-100.	0.1	1
31	Patients with pathological stage N2 rectal cancer treated with early adjuvant chemotherapy have a lower treatment failure rate. BMC Cancer, 2017, 17, 182.	1.1	1
32	Application of the national early warning score (NEWS) in patients with acute aortic dissection: A case–control study. Journal of Clinical Nursing, 2021, , .	1.4	1
33	Down-staging depth score could be a survival predictor for locally advanced gastric cancer patients after preoperative chemoradiotherapy. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2021, 33, 447-456.	0.7	1
34	Possible contribution of IMRT in postoperative radiochemotherapy for rectal cancer: analysis on 1798 patients by prediction model. Oncotarget, 2016, 7, 46536-46544.	0.8	1