

Toshihiko Matsumoto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5782263/publications.pdf>

Version: 2024-02-01

39
papers

476
citations

933264

10
h-index

752573

20
g-index

58
all docs

58
docs citations

58
times ranked

673
citing authors

#	ARTICLE	IF	CITATIONS
1	Variables associated with methamphetamine use within the past year and sex differences among patients with methamphetamine use disorder: A cross-sectional study in Japan. <i>American Journal on Addictions</i> , 2022, , .	1.3	2
2	Ifenprodil for the treatment of methamphetamine use disorder: An exploratory, randomized, double-blind, placebo-controlled trial. <i>Neuropsychopharmacology Reports</i> , 2022, , .	1.1	3
3	Gender Differences in Triggers of Stimulant Use Based on the National Survey of Prisoners in Japan. <i>Substance Use and Misuse</i> , 2021, 56, 54-60.	0.7	1
4	The relationship between severity of drug problems and perceived interdependence of drug use and sexual intercourse among adult males in drug addiction rehabilitation centers in Japan. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2021, 16, 5.	1.0	3
5	Does cognitive behavioral therapy for anxiety disorders assist the discontinuation of benzodiazepines among patients with anxiety disorders? A systematic review and meta-analysis. <i>Psychiatry and Clinical Neurosciences</i> , 2021, 75, 119-127.	1.0	11
6	Gender Differences in the Relationship between Methamphetamine Use and High-risk Sexual Behavior among Prisoners: A Nationwide, Cross-sectional Survey in Japan. <i>Journal of Psychoactive Drugs</i> , 2021, , 1-9.	1.0	1
7	Relapse prevention group therapy via video-conferencing for substance use disorder: protocol for a multicentre randomised controlled trial in Indonesia. <i>BMJ Open</i> , 2021, 11, e050259.	0.8	2
8	Effect of a web-based relapse prevention program on abstinence among Japanese drug users: A pilot randomized controlled trial. <i>Journal of Substance Abuse Treatment</i> , 2020, 111, 37-46.	1.5	13
9	Risk factors for the onset of dependence and chronic psychosis due to cannabis use: Survey of patients with cannabis-related psychiatric disorders. <i>Neuropsychopharmacology Reports</i> , 2020, 40, 332-341.	1.1	5
10	Parental drinking according to parental composition and adolescent binge drinking: findings from a nationwide high school survey in Japan. <i>BMC Public Health</i> , 2020, 20, 1878.	1.2	5
11	Changes of clinical symptoms in patients with new psychoactive substance (NPS)-related disorders from fiscal year 2012 to 2014: A study in hospitals specializing in the treatment of addiction. <i>Neuropsychopharmacology Reports</i> , 2019, 39, 119-129.	1.1	7
12	Study of effects of ifenprodil in patients with methamphetamine dependence: Protocol for an exploratory, randomized, double-blind, placebo-controlled trial. <i>Neuropsychopharmacology Reports</i> , 2019, 39, 90-99.	1.1	12
13	Factors influencing nursing home placement of patients with dementia: a retrospective, single-centre study in Japan. <i>Psychogeriatrics</i> , 2019, 19, 111-116.	0.6	1
14	The influence of tightening regulations on patients with new psychoactive substance-related disorders in Japan. <i>Neuropsychopharmacology Reports</i> , 2018, 38, 189-196.	1.1	5
15	The Efficacy of a Web-Based Screening and Brief Intervention for Reducing Alcohol Consumption Among Japanese Problem Drinkers: Protocol for a Single-Blind Randomized Controlled Trial. <i>JMIR Research Protocols</i> , 2018, 7, e10650.	0.5	2
16	Exploration of knowledge and skills that contribute to support of suicide attempts by clinical psychologists. <i>The Proceedings of the Annual Convention of the Japanese Psychological Association</i> , 2018, 82, 1AM-030-1AM-030.	0.0	0
17	Exploring suicide risk factors among Japanese individuals: The largest case-control psychological autopsy study in Japan. <i>Asian Journal of Psychiatry</i> , 2017, 27, 123-126.	0.9	10
18	Female suicides: Psychosocial and psychiatric characteristics identified by a psychological autopsy study in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2017, 71, 271-279.	1.0	11

#	ARTICLE	IF	CITATIONS
19	Characteristics of cancer patients who died by suicide in the Tokyo metropolitan area. Japanese Journal of Clinical Oncology, 2017, 47, 458-462.	0.6	2
20	Web-based cognitive behavioral relapse prevention program with tailored feedback for people with methamphetamine and other drug use problems: protocol for a multicenter randomized controlled trial in Japan. BMC Psychiatry, 2016, 16, 87.	1.1	8
21	Prevalence, prescribed quantities, and trajectory of multiple prescriber episodes for benzodiazepines: A 2-year cohort study. Drug and Alcohol Dependence, 2016, 158, 118-125.	1.6	23
22	Current Situation and Clinical Characteristics of Sedative-Related Disorder Patients in Japan: Comparison with Methamphetamine-Related Disorder Patients. Activa Nervosa Superior, 2015, 57, 12-28.	0.4	2
23	A Study of Therapeutic Attitudes Towards Working With Drug Abusers: Reliability and Validity of the Japanese Version of the Drug and Drug Problems Perception Questionnaire. Archives of Psychiatric Nursing, 2015, 29, 302-308.	0.7	5
24	Suicide prevention strategies in Japan: A 15-year review (1998-2013). Journal of Public Health Policy, 2015, 36, 52-66.	1.0	19
25	Clinical features of patients with designer drug-related disorder in Japan: A comparison with patients with methamphetamine and hypnotic/anxiolytic-related disorders. Psychiatry and Clinical Neurosciences, 2014, 68, 374-382.	1.0	9
26	Suicide risk among individuals with sleep disturbances in Japan: a case-control psychological autopsy study. Sleep Medicine, 2014, 15, 430-435.	0.8	39
27	Prevalence of problematic drinking among outpatients attending general hospitals in Tokyo. Japanese Journal of Alcohol Studies and Drug Dependence, 2013, 48, 300-13.	0.4	1
28	Risky sexual behaviors, mental health, and history of childhood abuse among adolescents. Asian Journal of Psychiatry, 2012, 5, 48-52.	0.9	13
29	Sex differences in risk factors for suicidality among Japanese substance use disorder patients: Association with age, types of abused substances, and depression. Psychiatry and Clinical Neurosciences, 2012, 66, 390-396.	1.0	8
30	Comparative study of suicide risk in depressive disorder patients with and without problem drinking. Psychiatry and Clinical Neurosciences, 2011, 65, 529-532.	1.0	3
31	Possible effectiveness of intervention using a self-teaching workbook in adolescent drug abusers detained in a juvenile classification home. Psychiatry and Clinical Neurosciences, 2011, 65, 576-583.	1.0	7
32	Depression and suicide risk of outpatients at specialized hospitals for substance use disorder: comparison with depressive disorder patients at general psychiatric clinics. Japanese Journal of Alcohol Studies and Drug Dependence, 2011, 46, 554-9.	0.4	1
33	Comparative study of the prevalence of suicidal behavior and sexual abuse history in delinquent and non-delinquent adolescents. Psychiatry and Clinical Neurosciences, 2009, 63, 238-240.	1.0	11
34	Self-injury in Japanese junior and senior high school students: Prevalence and association with substance use. Psychiatry and Clinical Neurosciences, 2008, 62, 123-125.	1.0	22
35	Profiles associated with treatment retention in Japanese patients with methamphetamine use disorder: Preliminary survey. Psychiatry and Clinical Neurosciences, 2008, 62, 526-532.	1.0	8
36	Association between childhood attention-deficit/hyperactivity symptoms and adulthood dissociation in male inmates: Preliminary report. Psychiatry and Clinical Neurosciences, 2007, 61, 444-446.	1.0	18

#	ARTICLE	IF	CITATIONS
37	Drug preferences in illicit drug abusers with a childhood tendency of attention deficit/hyperactivity disorder: A study using the Wender Utah Rating Scale in a Japanese prison. <i>Psychiatry and Clinical Neurosciences</i> , 2005, 59, 311-318.	1.0	26
38	Characteristics of self-cutters among male inmates: Association with bulimia and dissociation. <i>Psychiatry and Clinical Neurosciences</i> , 2005, 59, 319-326.	1.0	42
39	Methamphetamine in Japan: the consequences of methamphetamine abuse as a function of route of administration. <i>Addiction</i> , 2002, 97, 809-817.	1.7	90