

# SÃœleyman Bayram

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5787861/publications.pdf>

Version: 2024-02-01

32  
papers

642  
citations

566801

15  
h-index

580395

25  
g-index

32  
all docs

32  
docs citations

32  
times ranked

871  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Effect of Five Single Nucleotide Polymorphisms on Hb F Variation of $\beta^2$ -Thalassemia Traits and Hematologically Normal Individuals in Southeast Turkey. <i>Hemoglobin</i> , 2020, 44, 231-239.	0.4	1
2	Genetic polymorphisms in human telomerase reverse transcriptase (hTERT) gene polymorphisms do not associated with breast cancer in patients in a Turkish population: hospital-based case-control study. <i>Cellular and Molecular Biology</i> , 2018, 64, 108-115.	0.3	5
3	The relationship between the variations in G $\alpha$ and A $\alpha$ promoters and the Hereditary Persistence of Fetal Hemoglobin (HPFH). <i>Cellular and Molecular Biology</i> , 2018, 64, 32-39.	0.3	3
4	Investigation of XRCC1 Arg399Gln, Arg280His, and Arg194Trp polymorphisms effects on the induction of micronucleus by Aflatoxin B1 in in vitro. <i>Gene Reports</i> , 2017, 9, 1-6.	0.4	1
5	The prevalence and genotypes of alpha-thalassemia in Adana: two rare alpha variants. <i>Turkish Journal of Biochemistry</i> , 2016, 41, 467-472.	0.3	0
6	Genotoxicity and cytotoxicity of copper oxychloride in cultured human lymphocytes using cytogenetic and molecular tests. <i>Cytotechnology</i> , 2016, 68, 2027-2036.	0.7	6
7	Polymorphisms in human telomerase reverse transcriptase (h TERT ) gene and susceptibility to gastric cancer in a Turkish population: Hospital-based case-control study. <i>Gene</i> , 2016, 585, 84-92.	1.0	19
8	Effect of p53 Arg72Pro polymorphism on the induction of micronucleus by aflatoxin B1 in in vitro in human blood lymphocytes. <i>Drug and Chemical Toxicology</i> , 2016, 39, 331-337.	1.2	6
9	A functional HOTAIR rs12826786 C>T polymorphism is associated with breast cancer susceptibility and poor clinicopathological characteristics in a Turkish population: a hospital-based case-control study. <i>Tumor Biology</i> , 2016, 37, 5577-5584.	0.8	45
10	A functional HOTAIR rs920778 polymorphism does not contributes to gastric cancer in a Turkish population: a case-control study. <i>Familial Cancer</i> , 2015, 14, 561-567.	0.9	46
11	Effect of HOTAIR rs920778 polymorphism on breast cancer susceptibility and clinicopathologic features in a Turkish population. <i>Tumor Biology</i> , 2015, 36, 3863-3870.	0.8	73
12	MicroRNA 211 expression is upregulated and associated with poor prognosis in colorectal cancer: a case-control study. <i>Tumor Biology</i> , 2015, 36, 9703-9709.	0.8	17
13	The Exonuclease 1 Glu589Lys Gene Polymorphism and Cancer Susceptibility: Evidence Based on a Meta-analysis. <i>Asian Pacific Journal of Cancer Prevention</i> , 2014, 15, 2571-2576.	0.5	3
14	Association between RASSF1A Ala133Ser Polymorphism and Cancer Susceptibility: A Meta-Analysis Involving 8,892 Subjects. <i>Asian Pacific Journal of Cancer Prevention</i> , 2014, 15, 3691-3698.	0.5	5
15	The role of interleukin 28B gene polymorphism in Turkish patients with hepatocellular carcinoma. <i>Annals of Hepatology</i> , 2014, 13, 788-95.	0.6	7
16	Effect of PON1 gene polymorphisms in Turkish patients with hepatocellular carcinoma. <i>Meta Gene</i> , 2013, 1, 93-101.	0.3	13
17	CHK2 1100delC, IVS2+1G>A and I157T mutations are not present in hepatocellular cancer cases from a Turkish population. <i>Gene</i> , 2013, 512, 232-236.	1.0	3
18	RASSF1A Ala133Ser polymorphism is associated with increased susceptibility to hepatocellular carcinoma in a Turkish population. <i>Gene</i> , 2012, 498, 264-269.	1.0	11

#	ARTICLE	IF	CITATIONS
19	Prevalence of Beta-Thalassemia Trait and Abnormal Hemoglobins in the Province of AdÄ±yaman, Turkey. <i>Pediatric Hematology and Oncology</i> , 2012, 29, 620-623.	0.3	6
20	CHEK2 1100delC, IVS2+1G>A and I157T mutations are not present in colorectal cancer cases from Turkish population. <i>Cancer Epidemiology</i> , 2012, 36, 453-457.	0.8	10
21	Lack of an association of programmed cell death-1 PD1.3 polymorphism with risk of hepatocellular carcinoma susceptibility in Turkish population: A caseâ€“control study. <i>Gene</i> , 2012, 511, 308-313.	1.0	23
22	The significance of Exonuclease 1 K589E polymorphism on hepatocellular carcinoma susceptibility in the Turkish population: a caseâ€“control study. <i>Molecular Biology Reports</i> , 2012, 39, 5943-5951.	1.0	31
23	p53 codon 72 polymorphism is associated with susceptibility to hepatocellular carcinoma in the Turkish population: a caseâ€“control study. <i>Molecular Biology Reports</i> , 2012, 39, 1639-1647.	1.0	30
24	No association of pre-microRNA-146a rs2910164 polymorphism and risk of hepatocellular carcinoma development in Turkish population: A case-control study. <i>Gene</i> , 2011, 486, 104-109.	1.0	71
25	The association between the survivin âˆ“31G/C promoter polymorphism and hepatocellular carcinoma risk in a Turkish population. <i>Cancer Epidemiology</i> , 2011, 35, 555-559.	0.8	22
26	Functional polymorphisms of cyclooxygenase-2 gene and risk for hepatocellular carcinoma. <i>Molecular and Cellular Biochemistry</i> , 2011, 347, 201-208.	1.4	32
27	No association of the exonuclease 1 T439M polymorphism and risk of hepatocellular carcinoma development in the Turkish population: a case-control study. <i>Asian Pacific Journal of Cancer Prevention</i> , 2011, 12, 2455-60.	0.5	2
28	MDM2 promoter polymorphism is associated with increased susceptibility to hepatocellular carcinoma in Turkish population. <i>Cancer Epidemiology</i> , 2010, 34, 448-452.	0.8	24
29	The significance of E266K polymorphism in the NOD1 gene on Helicobacter Pylori infection: an effective force on pathogenesis?. <i>Clinical and Experimental Medicine</i> , 2010, 10, 107-112.	1.9	29
30	Cyclin D1 G870A polymorphism is associated with an increased risk of hepatocellular carcinoma in the Turkish population: Caseâ€“control study. <i>Cancer Epidemiology</i> , 2010, 34, 298-302.	0.8	31
31	G-308A TNF-Î± polymorphism is associated with an increased risk of hepatocellular carcinoma in the Turkish population: Case-control study. <i>Cancer Epidemiology</i> , 2009, 33, 261-264.	0.8	54
32	Confirmation of the chromosome damaging effects of lamivudine in in vitro human peripheral blood lymphocytes. <i>Environmental and Molecular Mutagenesis</i> , 2008, 49, 328-333.	0.9	13