

Jongmin Sim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5796614/publications.pdf>

Version: 2024-02-01

29
papers

304
citations

840776

11
h-index

940533

16
g-index

29
all docs

29
docs citations

29
times ranked

604
citing authors

#	ARTICLE	IF	CITATIONS
1	Loss of microRNA-200a expression correlates with tumor progression in breast cancer. <i>Translational Research</i> , 2014, 163, 242-251.	5.0	32
2	FOXO3a expression is associated with lymph node metastasis and poor disease-free survival in triple-negative breast cancer. <i>Journal of Clinical Pathology</i> , 2018, 71, 806-813.	2.0	23
3	High MicroRNA-370 Expression Correlates with Tumor Progression and Poor Prognosis in Breast Cancer. <i>Journal of Breast Cancer</i> , 2015, 18, 323.	1.9	22
4	Increased Expression of Forkhead Box M1 Is Associated with Aggressive Phenotype and Poor Prognosis in Estrogen Receptor-Positive Breast Cancer. <i>Journal of Korean Medical Science</i> , 2015, 30, 390.	2.5	22
5	Overexpression of Forkhead Box O3a and Its Association With Aggressive Phenotypes and Poor Prognosis in Human Hepatocellular Carcinoma. <i>American Journal of Clinical Pathology</i> , 2018, 149, 117-127.	0.7	22
6	Splenic hamartoma: A case report and review of the literature. <i>World Journal of Clinical Cases</i> , 2013, 1, 217.	0.8	17
7	Clinicopathological Significance of Dual-Specificity Protein Phosphatase 4 Expression in Invasive Ductal Carcinoma of the Breast. <i>Journal of Breast Cancer</i> , 2015, 18, 1.	1.9	17
8	Clinicopathological significance of CADM4 expression, and its correlation with expression of E-cadherin and Ki-67 in colorectal adenocarcinomas. <i>Journal of Clinical Pathology</i> , 2012, 65, 902-906.	2.0	14
9	Comparison of 1p and 19q status of glioblastoma by whole exome sequencing, array-comparative genomic hybridization, and fluorescence in situ hybridization. <i>Medical Oncology</i> , 2018, 35, 60.	2.5	14
10	Immunohistochemical Expression of Dual-Specificity Protein Phosphatase 4 in Patients with Colorectal Adenocarcinoma. <i>Gastroenterology Research and Practice</i> , 2015, 2015, 1-8.	1.5	13
11	Anaplastic Transformation of Papillary Thyroid Carcinoma Only Seen in Pleural Metastasis: A Case Report with Review of the Literature. <i>Head and Neck Pathology</i> , 2017, 11, 162-167.	2.6	11
12	Identification of recurrence-associated microRNAs in stage I lung adenocarcinoma. <i>Medicine (United States)</i> , 2018, 97, 1071-1078.	1.8	11
13	MET is overexpressed in microsatellite instability-high gastric carcinoma. <i>Pathology Research and Practice</i> , 2019, 215, 433-438.	2.3	10
14	Clinicopathological significance of CADM4 expression in invasive ductal carcinoma of the breast. <i>Journal of Clinical Pathology</i> , 2013, 66, 681-686.	2.0	8
15	Single-stranded DNA binding protein 2 expression is associated with patient survival in hepatocellular carcinoma. <i>BMC Cancer</i> , 2018, 18, 1244.	2.6	7
16	Comparison of the Lymph2Cx Assay and Hans Algorithm in Determining the Cell-of-Origin of Diffuse Large B-Cell Lymphomas, Not Otherwise Specified. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2020, 28, 731-740.	1.2	7
17	Pulmonary Nodular Lymphoid Hyperplasia with Mass-Formation: Clinicopathologic Characteristics of Nine Cases and Review of the Literature. <i>Journal of Pathology and Translational Medicine</i> , 2018, 52, 211-218.	1.1	7
18	Lymphoepithelioma-like Carcinoma of the Renal Pelvis: A Case Report and Review of the Literature. <i>Korean Journal of Pathology</i> , 2014, 48, 458-461.	1.3	6

#	ARTICLE	IF	CITATIONS
19	Loss of MTUS1 Expression Is Associated With Poor Prognosis in Patients With Gallbladder Carcinoma. <i>In Vivo</i> , 2020, 34, 125-132.	1.3	6
20	Low Expression of Single-stranded DNA Binding Protein 2 (SSBP2) Predicts Unfavourable Postoperative Outcomes in Patients With Clear Cell Renal Cell Carcinoma. <i>In Vivo</i> , 2020, 34, 101-107.	1.3	5
21	Clinicopathological Significance of MTUS1 Expression in Patients With Renal Cell Carcinoma. <i>Anticancer Research</i> , 2020, 40, 2961-2967.	1.1	5
22	Anaplastic lymphoma kinase (ALK)-expressing Lung Adenocarcinoma with Combined Neuroendocrine Component or Neuroendocrine Transformation: Implications for Neuroendocrine Transformation and Response to ALK-tyrosine Kinase Inhibitors. <i>Journal of Korean Medical Science</i> , 2018, 33, e123.	2.5	5
23	Clinicopathologic Correlations of E-cadherin and Prrx-1 Expression Loss in Hepatocellular Carcinoma. <i>Journal of Pathology and Translational Medicine</i> , 2016, 50, 327-336.	1.1	5
24	Correlation of CD47 Expression with Adverse Clinicopathologic Features and an Unfavorable Prognosis in Colorectal Adenocarcinoma. <i>Diagnostics</i> , 2021, 11, 668.	2.6	4
25	Low-Level Expression of MTUS1 Is Associated with Poor Survival in Patients with Lung Adenocarcinoma. <i>Diagnostics</i> , 2021, 11, 1250.	2.6	4
26	MicroRNA-374a Expression as a Prognostic Biomarker in Lung Adenocarcinoma. <i>Journal of Pathology and Translational Medicine</i> , 2019, 53, 354-360.	1.1	4
27	The clinicopathologic significance of extranodal tumor extension in locally advanced (pT3) colorectal adenocarcinoma and its association with the loss of E-cadherin expression. <i>International Journal of Clinical and Experimental Pathology</i> , 2019, 12, 3417-3425.	0.5	1
28	Loss of Wnt7a expression correlates with tumor progression and poor prognosis in colorectal carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 4967-4976.	0.5	1
29	Decreased Expression of Cell Adhesion Molecule 4 in Gastric Adenocarcinoma and Its Prognostic Implications. <i>Diagnostics</i> , 2022, 12, 941.	2.6	1