

Tadashi Yamasaki

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/5802743/tadashi-yamasaki-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

27
papers

743
citations

13
h-index

27
g-index

28
ext. papers

817
ext. citations

4.1
avg, IF

4.18
L-index

#	Paper	IF	Citations
27	Variable inflation rate of a magmatic deformation source beneath Aira caldera after the 1914 eruption of Sakurajima volcano: Inferences from a linear Maxwell viscoelastic model constrained by geodetic data. <i>Journal of Volcanology and Geothermal Research</i> , 2021 , 421, 107446	2.8	
26	The influence of elastic thickness non-uniformity on viscoelastic crustal response to magma emplacement: application to the Kutcharo caldera, eastern Hokkaido, Japan. <i>Geophysical Journal International</i> , 2020 , 224, 701-718	2.6	2
25	Unexpected large eruptions from buoyant magma bodies within viscoelastic crust. <i>Nature Communications</i> , 2020 , 11, 2403	17.4	16
24	Magmatic-hydrothermal system of Aso Volcano, Japan, inferred from electrical resistivity structures. <i>Earth, Planets and Space</i> , 2020 , 72,	2.9	5
23	Viscoelastic crustal response to magma supply and discharge in the upper crust: Implications for the uplift of the Aira caldera before and after the 1914 eruption of the Sakurajima volcano. <i>Earth and Planetary Science Letters</i> , 2020 , 531, 115981	5.3	4
22	Viscoelastic crustal deformation by magmatic intrusion: A case study in the Kutcharo caldera, eastern Hokkaido, Japan. <i>Journal of Volcanology and Geothermal Research</i> , 2018 , 349, 128-145	2.8	18
21	Crustal Deformation Infers a Magma Chamber. <i>Journal of Geography (Chigaku Zasshi)</i> , 2018 , 127, 111-138	0.5	3
20	Imaging a low viscosity zone beneath the Kutcharo caldera, eastern Hokkaido, Japan, using geodetic data. <i>Earth and Planetary Science Letters</i> , 2018 , 504, 1-12	5.3	5
19	Analysis of the spatial viscosity variation in the crust beneath the western North Anatolian Fault. <i>Journal of Geodynamics</i> , 2015 , 88, 80-89	2.2	3
18	Crustal and basin evolution of the southwestern Barents Sea: From Caledonian orogeny to continental breakup. <i>Tectonics</i> , 2014 , 33, 347-373	4.3	81
17	Weak ductile shear zone beneath a major strike-slip fault: Inferences from earthquake cycle model constrained by geodetic observations of the western North Anatolian Fault Zone. <i>Journal of Geophysical Research: Solid Earth</i> , 2014 , 119, 3678-3699	3.6	45
16	The signature of depth-dependent viscosity structure in post-seismic deformation. <i>Geophysical Journal International</i> , 2012 , 190, 769-784	2.6	14
15	The Norway Basin revisited: From continental breakup to spreading ridge extinction. <i>Marine and Petroleum Geology</i> , 2012 , 35, 1-19	4.7	63
14	The crustal viscosity gradient measured from post-seismic deformation: A case study of the 1997 Manyi (Tibet) earthquake. <i>Earth and Planetary Science Letters</i> , 2012 , 351-352, 105-114	5.3	37
13	Back-arc rifting initiated with a hot and wet continental lithosphere. <i>Earth and Planetary Science Letters</i> , 2011 , 302, 172-184	5.3	13
12	Redistribution of the lithosphere deformation by the emplacement of underplated mafic bodies: implications for microcontinent formation. <i>Journal of the Geological Society</i> , 2010 , 167, 961-971	2.7	18
11	Change in tectonic force inferred from basin subsidence: Implications for the dynamical aspects of back-arc rifting in the western Mediterranean. <i>Earth and Planetary Science Letters</i> , 2009 , 277, 174-183	5.3	8

10	Styles of lithospheric extension controlled by underplated mafic bodies. <i>Tectonophysics</i> , 2009 , 468, 169-184	3.1	28
9	Potential role of strain hardening in the cessation of rifting at constant tectonic force. <i>Journal of Geodynamics</i> , 2009 , 47, 47-62	2.2	8
8	Numerical modelling study on the flexural uplift of the Transantarctic Mountains. <i>Geophysical Journal International</i> , 2008 , 174, 377-390	2.6	12
7	A rheological weak zone intensified by post-rift thermal relaxation as a possible origin of simple shear deformation associated with reactivation of rifting. <i>Earth and Planetary Science Letters</i> , 2006 , 248, 134-146	5.3	9
6	High strain rate zone in central Honshu resulting from the viscosity heterogeneities in the crust and mantle. <i>Earth and Planetary Science Letters</i> , 2005 , 232, 13-27	5.3	34
5	Localized rheological weakening by grain-size reduction during lithospheric extension. <i>Tectonophysics</i> , 2004 , 386, 117-145	3.1	9
4	Double seismic zone and dehydration embrittlement of the subducting slab. <i>Journal of Geophysical Research</i> , 2003 , 108,		197
3	Low-frequency tremors, intraslab and interplate earthquakes in Southwest Japan from a viewpoint of slab dehydration. <i>Geophysical Research Letters</i> , 2003 , 30,	4.9	89
2	The effects of the spinel-garnet phase transition on the formation of rifted sedimentary basins. <i>Geophysical Journal International</i> , 1997 , 130, 681-692	2.6	18
1	Effects of the Quaternary sea-level change on the subsidence of a sedimentary basin: a case study of the Osaka Bay sedimentary basin, Japan. <i>Tectonophysics</i> , 1996 , 267, 229-238	3.1	4