

Jurrien Dean

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5806540/publications.pdf>

Version: 2024-02-01

32
papers

3,034
citations

304368

22
h-index

414034

32
g-index

38
all docs

38
docs citations

38
times ranked

2860
citing authors

#	ARTICLE	IF	CITATIONS
1	Sperm acrosome overgrowth and infertility in mice lacking chromosome 18 pachytene piRNA. PLoS Genetics, 2021, 17, e1009485.	1.5	39
2	Maternal factors regulating preimplantation development in mice. Current Topics in Developmental Biology, 2020, 140, 317-340.	1.0	22
3	EXOSC10 sculpts the transcriptome during the growth-to-maturation transition in mouse oocytes. Nucleic Acids Research, 2020, 48, 5349-5365.	6.5	28
4	Anchoring cortical granules in the cortex ensures trafficking to the plasma membrane for post-fertilization exocytosis. Nature Communications, 2019, 10, 2271.	5.8	19
5	Sertoli cell-only phenotype and scRNA-seq define PRAMEF12 as a factor essential for spermatogenesis in mice. Nature Communications, 2019, 10, 5196.	5.8	35
6	The Zona Pellucida Facilitates Fertilization, Blocks Polyspermy and Protects Pre-Implantation Embryos. , 2018, , 294-299.		0
7	Glycan-Independent Gamete Recognition Triggers Egg Zinc Sparks and ZP2 Cleavage to Prevent Polyspermy. Developmental Cell, 2018, 46, 627-640.e5.	3.1	61
8	BTBD18 Regulates a Subset of piRNA-Generating Loci through Transcription Elongation in Mice. Developmental Cell, 2017, 40, 453-466.e5.	3.1	30
9	Cytoplasmic cleavage of DPPA3 is required for intracellular trafficking and cleavage-stage development in mice. Nature Communications, 2017, 8, 1643.	5.8	16
10	A Unique Egg Cortical Granule Localization Motif Is Required for Ovastacin Sequestration to Prevent Premature ZP2 Cleavage and Ensure Female Fertility in Mice. PLoS Genetics, 2017, 13, e1006580.	1.5	40
11	Genetic mosaics and time-lapse imaging identify functions of H3.3 residues in mouse oocytes and embryos. Development (Cambridge), 2016, 144, 519-528.	1.2	8
12	BTG4, a maternal mRNA cleaner. Journal of Molecular Cell Biology, 2016, 8, 369-370.	1.5	15
13	ZP2 peptide beads select human sperm in vitro, decoy mouse sperm in vivo, and provide reversible contraception. Science Translational Medicine, 2016, 8, 336ra60.	5.8	39
14	Exact Requirements for Development of the Egg. New England Journal of Medicine, 2016, 374, 279-280.	13.9	14
15	Reprogramming the genome to totipotency in mouse embryos. Trends in Cell Biology, 2015, 25, 82-91.	3.6	89
16	Figla-Cre Transgenic Mice Expressing Myristoylated EGFP in Germ Cells Provide a Model for Investigating Perinatal Oocyte Dynamics. PLoS ONE, 2014, 9, e84477.	1.1	12
17	The subcortical maternal complex controls symmetric division of mouse zygotes by regulating F-actin dynamics. Nature Communications, 2014, 5, 4887.	5.8	102
18	A single domain of the ZP2 zona pellucida protein mediates gamete recognition in mice and humans. Journal of Cell Biology, 2014, 205, 801-809.	2.3	146

#	ARTICLE	IF	CITATIONS
19	Ovastacin, a cortical granule protease, cleaves ZP2 in the zona pellucida to prevent polyspermy. <i>Journal of Cell Biology</i> , 2012, 197, 37-44.	2.3	241
20	Human sperm bind to the N-terminal domain of ZP2 in humanized zonae pellucidae in transgenic mice. <i>Journal of Cell Biology</i> , 2012, 197, 897-905.	2.3	89
21	Gamete Recognition in Mice Depends on the Cleavage Status of an Egg's Zona Pellucida Protein. <i>Science</i> , 2010, 329, 216-219.	6.0	163
22	Role of <i>Filia</i> , a maternal effect gene, in maintaining euploidy during cleavage-stage mouse embryogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 7473-7478.	3.3	134
23	A Subcortical Maternal Complex Essential for Preimplantation Mouse Embryogenesis. <i>Developmental Cell</i> , 2008, 15, 416-425.	3.1	242
24	Maternally derived FILIA-MATER complex localizes asymmetrically in cleavage-stage mouse embryos. <i>Development (Cambridge)</i> , 2008, 135, 259-269.	1.2	102
25	Ovarian gene expression in the absence of FIGLA, an oocyte-specific transcription factor. <i>BMC Developmental Biology</i> , 2007, 7, 67.	2.1	102
26	The enigma of sperm-egg recognition in mice. <i>Society of Reproduction and Fertility Supplement</i> , 2007, 63, 359-65.	0.2	8
27	Reassessing the molecular biology of sperm-egg recognition with mouse genetics. <i>BioEssays</i> , 2004, 26, 29-38.	1.2	80
28	Fertility and Taxon-Specific Sperm Binding Persist after Replacement of Mouse Sperm Receptors with Human Homologs. <i>Developmental Cell</i> , 2003, 5, 33-43.	3.1	155
29	Oocyte-specific genes regulate follicle formation, fertility and early mouse development. <i>Journal of Reproductive Immunology</i> , 2002, 53, 171-180.	0.8	117
30	Expression of Cre recombinase in mouse oocytes: A means to study maternal effect genes. <i>Genesis</i> , 2000, 26, 110-112.	0.8	329
31	Mater, a maternal effect gene required for early embryonic development in mice. <i>Nature Genetics</i> , 2000, 26, 267-268.	9.4	487
32	Mater encodes a maternal protein in mice with a leucine-rich repeat domain homologous to porcine ribonuclease inhibitor. <i>Mammalian Genome</i> , 2000, 11, 281-287.	1.0	65