

# Annie Palstam

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5806602/publications.pdf>

Version: 2024-02-01

29  
papers

802  
citations

471509

17  
h-index

526287

27  
g-index

32  
all docs

32  
docs citations

32  
times ranked

913  
citing authors

#	ARTICLE	IF	CITATIONS
1	A register-based study on associations between pre-stroke physical activity and cognition early after stroke (part of PAPSIGOT). <i>Scientific Reports</i> , 2022, 12, 5779.	3.3	2
2	Self-Assessed Aspects of Health 3 Months after COVID-19 Hospitalization—A Swedish Cross-Sectional Study. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 8020.	2.6	7
3	Patterns and predictors of sick leave after Covid-19 and long Covid in a national Swedish cohort. <i>BMC Public Health</i> , 2021, 21, 1023.	2.9	73
4	Women with Fibromyalgia Prefer Resistance Exercise with Heavy Loads—A Randomized Crossover Pilot Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6276.	2.6	3
5	Recurrent sick leave after COVID-19: investigating the first wave of the pandemic in a comprehensive Swedish registry-based study. <i>BMC Public Health</i> , 2021, 21, 1914.	2.9	11
6	A Call to Include a Perspective of Sustainable Development in Physical Therapy Research. <i>Physical Therapy</i> , 2021, 101, .	2.4	10
7	Physical Function, Cognitive Function, and Daily Activities in Patients Hospitalized Due to COVID-19: A Descriptive Cross-Sectional Study in Sweden. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 11600.	2.6	13
8	Differences in self-perceived general health, pain, and depression 1 to 5 years post-stroke related to work status at 1 year. <i>Scientific Reports</i> , 2020, 10, 13251.	3.3	2
9	Transforming self-reported outcomes from a stroke register to the modified Rankin Scale: a cross-sectional, explorative study. <i>Scientific Reports</i> , 2020, 10, 17215.	3.3	1
10	Participation and autonomy five years after stroke: A longitudinal observational study. <i>PLoS ONE</i> , 2019, 14, e0219513.	2.5	45
11	Very Early MoCA Can Predict Functional Dependence at 3 Months After Stroke: A Longitudinal, Cohort Study. <i>Frontiers in Neurology</i> , 2019, 10, 1051.	2.4	26
12	“My life after stroke through a camera lens”—A photovoice study on participation in Sweden. <i>PLoS ONE</i> , 2019, 14, e0222099.	2.5	10
13	Work-related predictors for return to work after stroke. <i>Acta Neurologica Scandinavica</i> , 2019, 139, 382-388.	2.1	18
14	Aspects of exercise with person-centred guidance influencing the transition to independent exercise: a qualitative interview study among older adults with rheumatoid arthritis. <i>European Review of Aging and Physical Activity</i> , 2019, 16, 4.	2.9	18
15	Controlled, cross-sectional, multi-center study of physical capacity and associated factors in women with fibromyalgia. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 121.	1.9	23
16	Experiences of returning to work and maintaining work 7 to 8 years after a stroke: a qualitative interview study in Sweden. <i>BMJ Open</i> , 2018, 8, e021182.	1.9	26
17	Patient-reported outcome measures in osteoarthritis: a systematic search and review of their use and psychometric properties. <i>RMD Open</i> , 2018, 4, e000715.	3.8	22
18	Prestroke physical activity could influence acute stroke severity (part of PAPSIGOT). <i>Neurology</i> , 2018, 91, e1461-e1467.	1.1	65

#	ARTICLE	IF	CITATIONS
19	Gene-to-gene interactions regulate endogenous pain modulation in fibromyalgia patients and healthy controlsâ€”antagonistic effects between opioid and serotonin-related genes. <i>Pain</i> , 2017, 158, 1194-1203.	4.2	54
20	Benefits of resistance exercise in lean women with fibromyalgia: involvement of IGF-1 and leptin. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 106.	1.9	19
21	Work Ability in Fibromyalgia: An Update in the 21st Century. <i>Current Rheumatology Reviews</i> , 2017, 13, 180-187.	0.8	30
22	Decrease of fear avoidance beliefs following person-centered progressive resistance exercise contributes to reduced pain disability in women with fibromyalgia: secondary exploratory analyses from a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2016, 18, 116.	3.5	28
23	Effects of 15 weeks of resistance exercise on pro-inflammatory cytokine levels in the vastus lateralis muscle of patients with fibromyalgia. <i>Arthritis Research and Therapy</i> , 2016, 18, 137.	3.5	22
24	Resistance exercise improves physical fatigue in women with fibromyalgia: a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2016, 18, 176.	3.5	52
25	Comparison of the Levels of Pro-Inflammatory Cytokines Released in the Vastus Lateralis Muscle of Patients with Fibromyalgia and Healthy Controls during Contractions of the Quadriceps Muscle â€” A Microdialysis Study. <i>PLoS ONE</i> , 2015, 10, e0143856.	2.5	32
26	Resistance exercise improves muscle strength, health status and pain intensity in fibromyalgiaâ€”a randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2015, 17, 161.	3.5	122
27	Perceived exertion at work in women with fibromyalgia: Explanatory factors and comparison with healthy women. <i>Journal of Rehabilitation Medicine</i> , 2014, 46, 773-780.	1.1	18
28	Factors promoting sustainable work in women with fibromyalgia. <i>Disability and Rehabilitation</i> , 2013, 35, 1622-1629.	1.8	30
29	Which aspects of health differ between working and nonworking women with fibromyalgia? A cross-sectional study of work status and health. <i>BMC Public Health</i> , 2012, 12, 1076.	2.9	20