

# Masayoshi Kobayashi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5809455/publications.pdf>

Version: 2024-02-01

15  
papers

314  
citations

1040056

9  
h-index

996975

15  
g-index

16  
all docs

16  
docs citations

16  
times ranked

380  
citing authors

#	ARTICLE	IF	CITATIONS
1	Is the aortic depressor nerve involved in arterial chemoreflexes in rats?. Journal of the Autonomic Nervous System, 1999, 78, 38-48.	1.9	67
2	Olfactory Nerve Recovery Following Mild and Severe Injury and the Efficacy of Dexamethasone Treatment. Chemical Senses, 2009, 34, 573-580.	2.0	56
3	Cross-Cultural Comparison of Data Using the Odor Stick Identification Test for Japanese (OSIT-J). Chemical Senses, 2006, 31, 335-342.	2.0	55
4	A New Clinical Olfactory Function Test. JAMA Otolaryngology, 2007, 133, 331.	1.2	26
5	The Odor Stick Identification Test for the Japanese (OSIT-J): Clinical Suitability for Patients Suffering from Olfactory Disturbance. Chemical Senses, 2005, 30, i216-i217.	2.0	22
6	Blockade of interleukin-6 receptor suppresses inflammatory reaction and facilitates functional recovery following olfactory system injury. Neuroscience Research, 2013, 76, 125-132.	1.9	20
7	Tumor necrosis factor- $\alpha$ antagonist suppresses local inflammatory reaction and facilitates olfactory nerve recovery following injury. Auris Nasus Larynx, 2017, 44, 70-78.	1.2	14
8	Anti-high mobility group box 1 antibody suppresses local inflammatory reaction and facilitates olfactory nerve recovery following injury. Journal of Neuroinflammation, 2018, 15, 124.	7.2	14
9	Influence of visual information and test paradigm on clinical olfactory test results. Auris Nasus Larynx, 2008, 35, 53-60.	1.2	8
10	Target site of inhibition of baroreflex vagal bradycardia by nasal stimulation. Brain Research, 2004, 1009, 137-146.	2.2	7
11	Late phase responses after nasal challenges with allergen and histamine in asthmatic children with perennial nasal allergy. Auris Nasus Larynx, 2001, 28, 305-310.	1.2	5
12	A Time Limit for Initiating Anti-Inflammatory Treatment for Improved Olfactory Function after Head Injury. Journal of Neurotrauma, 2018, 35, 652-660.	3.4	4
13	Endoscopic endonasal transmaxillary ligation of a feeding artery and coblation plasma technology enables en bloc resection of advanced juvenile nasopharyngeal angiofibroma without preoperative embolization. Auris Nasus Larynx, 2019, 46, 306-310.	1.2	3
14	Optical coherence tomography for observation of the olfactory epithelium in mice. Auris Nasus Larynx, 2019, 46, 230-237.	1.2	3
15	High-dose IgG suppresses local inflammation and facilitates functional recovery after olfactory system injury. Annals of Clinical and Translational Neurology, 2022, , .	3.7	0