## **Anners Lerdal**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5814900/publications.pdf

Version: 2024-02-01

132	4,192	31 h-index	57
papers	citations		g-index
134	134 docs citations	134	5213
all docs		times ranked	citing authors

#	Article	IF	CITATIONS
1	Unidentified communication challenges in the intensive care unit: AÂqualitative study using multiple triangulations. Australian Critical Care, 2023, 36, 215-222.	0.6	10
2	Many Patients With Persistent Pain 1 Year After TKA Report Improvement by 5 to 7 Years: A Mixed-methods Study. Clinical Orthopaedics and Related Research, 2022, 480, 2075-2088.	0.7	4
3	Personally-Generated Quality of Life Outcomes in Adults with Juvenile Idiopathic Arthritis. Journal of Rheumatology, 2022, , jrheum.211245.	1.0	4
4	Factors Correlated With Physical Function 1 Year After Total Knee Arthroplasty in Patients With Knee Osteoarthritis. JAMA Network Open, 2022, 5, e2219636.	2.8	12
5	Preoperative risk factors associated with chronic pain profiles following total knee arthroplasty. European Journal of Pain, 2021, 25, 680-692.	1.4	23
6	Sleep disruption and duration are associated with variants in genes involved in energy homeostasis in adults with HIV/AIDS. Sleep Medicine, 2021, 82, 84-95.	0.8	2
7	Substance use in the Norwegian general population: prevalence and associations with disease. Journal of Substance Use, 2021, 26, 144-150.	0.3	3
8	Treatment Satisfaction With and Adherence to Diseaseâ€Modifying Antirheumatic Drugs in Adult Patients With Juvenile Idiopathic Arthritis. Arthritis Care and Research, 2021, 73, 221-231.	1.5	9
9	A mouth rinse based on a tea solution of Salvia officinalis for oral discomfort in palliative cancer care: a randomized controlled trial. Supportive Care in Cancer, 2021, 29, 4997-5007.	1.0	11
10	Authors' reply to the comment by Riddle et al. European Journal of Pain, 2021, 25, 1173-1174.	1.4	0
11	The Association between the Alcohol Biomarker Phosphatidylethanol (PEth) and Self-Reported Alcohol Consumption among Russian and Norwegian Medical Patients. Alcohol and Alcoholism, 2021, 56, 726-736.	0.9	7
12	Post-stroke fatigue: an exploratory study with patients and health professionals to develop a patient-reported outcome measure. Journal of Patient-Reported Outcomes, 2021, 5, 35.	0.9	19
13	A cross-sectional study exploring overlap in post-stroke fatigue caseness using three fatigue instruments: Fatigue Severity Scale, Fatigue Questionnaire and the Lynch's Clinical Interview. Journal of Psychosomatic Research, 2021, 150, 110605.	1.2	5
14	Multimorbidity, psychoactive substance use and psychological distress among acute medically ill patients: a cross-sectional study. BMJ Open, 2021, 11, e052428.	0.8	1
15	A literature review of factors associated with fatigue after stroke and a proposal for a framework for clinical utility. Neuropsychological Rehabilitation, 2020, 30, 1449-1476.	1.0	37
16	High levels of preoperative pain and fatigue are red flags for moderateâ€severe pain 12 months after total knee arthroplasty—A longitudinal cohort study. Musculoskeletal Care, 2020, 19, 186-192.	0.6	8
17	Daily use of alcohol in the Norwegian general population: Prevalence and associated factors. Drugs and Alcohol Today, 2020, 20, 109-121.	0.3	2
18	More Severe Radiographic Osteoarthritis Is Associated With Increased Improvement in Patients' Health State Following a Total Knee Arthroplasty. Journal of Arthroplasty, 2020, 35, 3131-3137.	1.5	9

#	Article	IF	Citations
19	Sexual Assault and the Association With Health, Quality of Life, and Self-Efficacy in the General Norwegian Population. Journal of Interpersonal Violence, 2020, , 088626052092630.	1.3	6
20	Prevalence of psychoactive substance use among acutely hospitalised patients in Oslo and Moscow: a cross-sectional, observational study. BMJ Open, 2020, 10, e032572.	0.8	4
21	Predictors of chronic pain and level of physical function in total knee arthroplasty: a protocol for a systematic review and meta-analysis. BMJ Open, 2020, 10, e037674.	0.8	3
22	High pre- and postoperative symptom burden in non-responders to total knee arthroplasty. PLoS ONE, 2020, 15, e0233347.	1.1	4
23	Psychometric properties of a short version of Lee Fatigue Scale used as a generic PROM in persons with stroke or osteoarthritis: assessment using a Rasch analysis approach. Health and Quality of Life Outcomes, 2020, 18, 168.	1.0	5
24	Trajectories of fatigue among stroke patients from the acute phase to 18 months post-injury: A latent class analysis. PLoS ONE, 2020, 15, e0231709.	1.1	22
25	Lack of content overlap and essential dimensions – A review of measures used for post-stroke fatigue. Journal of Psychosomatic Research, 2019, 124, 109759.	1.2	24
26	Perception of illness among patients with heart failure is related to their general health independently of their mood and functional capacity. Journal of Patient-Reported Outcomes, 2019, 3, 55.	0.9	16
27	Serious life events and post-traumatic stress disorder in the Norwegian population. BJPsych Open, 2019, 5, e82.	0.3	35
28	Flight Anxiety Reported from 1986 to 2015. Aerospace Medicine and Human Performance, 2019, 90, 384-388.	0.2	9
29	Optimists report fewer physical and mental health conditions than pessimists in the general Norwegian population. Health Psychology Report, 2019, 7, 9-18.	0.5	4
30	Longitudinal Health Status from Early Disease to Adulthood and Associated Prognostic Factors in Juvenile Idiopathic Arthritis. Journal of Rheumatology, 2019, 46, 1335-1344.	1.0	13
31	Self-evaluated anxiety in the Norwegian population: prevalence and associated factors. Archives of Public Health, 2019, 77, 10.	1.0	18
32	Predicting postoperative fatigue in surgically treated lung cancer patients in Norway: a longitudinal 5-month follow-up study. BMJ Open, 2019, 9, e028192.	0.8	6
33	Self-rated global health in the Norwegian general population. Health and Quality of Life Outcomes, 2019, 17, 188.	1.0	13
34	Effectiveness of nursing interventions for breathlessness in people with chronic obstructive pulmonary disease: AÂsystematic review and metaâ€analysis. Journal of Advanced Nursing, 2019, 75, 927-945.	1.5	4
35	Description of a multifaceted intervention programme for fatigue after acquired brain injury: a pilot study. Neuropsychological Rehabilitation, 2019, 29, 946-968.	1.0	7
36	General self-efficacy in the Norwegian population: Differences and similarities between sociodemographic groups. Scandinavian Journal of Public Health, 2019, 47, 695-704.	1.2	45

#	Article	lF	CITATIONS
37	Associations between patient symptoms and falls in an acute care hospital: A crossâ€sectional study. Journal of Clinical Nursing, 2018, 27, 1826-1835.	1.4	16
38	Physical Functioning, Pain, and Healthâ€Related Quality of Life in Adults With Juvenile Idiopathic Arthritis: A Longitudinal 30â€Year Followup Study. Arthritis Care and Research, 2018, 70, 741-749.	1.5	56
39	Prevalence of pressure ulcer and associated risk factors in middle―and olderâ€aged medical inpatients in Norway. Journal of Clinical Nursing, 2018, 27, e535-e543.	1.4	38
40	Higher physical activity and lower pain levels before surgery predict non-improvement of knee pain 1 Ayear after TKA. Knee Surgery, Sports Traumatology, Arthroscopy, 2018, 26, 1698-1708.	2.3	21
41	Rasch analysis of the Herth Hope Index in cancer patients. Health and Quality of Life Outcomes, 2018, 16, 196.	1.0	23
42	Self-diagnosed depression in the Norwegian general population – associations with neuroticism, extraversion, optimism, and general self-efficacy. BMC Public Health, 2018, 18, 1076.	1.2	35
43	Investigating post-stroke fatigue: An individual participant data meta-analysis. Journal of Psychosomatic Research, 2018, 113, 107-112.	1.2	42
44	The Impact of Demographic, Clinical, Symptom and Psychological Characteristics on the Trajectories of Acute Postoperative Pain After Total Knee Arthroplasty. Pain Medicine, 2017, 18, 124-139.	0.9	25
45	Supplementing the Braden scale for pressure ulcer risk among medical inpatients: the contribution of selfâ€reported symptoms and standard laboratory tests. Journal of Clinical Nursing, 2017, 26, 202-214.	1.4	30
46	Pressure ulcer is associated with malnutrition as assessed by Nutritional Risk Screening (NRS 2002) in a mixed hospital population. Food and Nutrition Research, 2017, 61, 1324230.	1.2	21
47	Acute-Phase Fatigue Predicts Limitations with Activities of Daily Living 18 Months after First-Ever Stroke. Journal of Stroke and Cerebrovascular Diseases, 2017, 26, 523-531.	0.7	27
48	Population-based norms of the Life Orientation Test–Revised (LOT-R). International Journal of Clinical and Health Psychology, 2017, 17, 216-224.	2.7	62
49	The relationship between pain with walking and self-rated health 12Âmonths following total knee arthroplasty: a longitudinal study. BMC Musculoskeletal Disorders, 2017, 18, 75.	0.8	13
50	Predictors of physical and mental health in persons with morbid obesity attending a patient education course – a two-year follow-up study. Health and Quality of Life Outcomes, 2017, 15, 103.	1.0	4
51	Psychometric limitations of the 13-item Sense of Coherence Scale assessed by Rasch analysis. BMC Psychology, 2017, 5, 18.	0.9	13
52	A systematic review of nursing interventions for breathlessness in COPD. , 2017, , .		0
53	Differences and similarities in the trajectories of self-esteem and positive and negative affect in persons with chronic illness: an explorative longitudinal study. Journal of Multidisciplinary Healthcare, 2016, Volume 9, 355-363.	1.1	2
54	Trajectories of physical and mental health among persons with morbid obesity and persons with COPD: a longitudinal comparative study. Journal of Multidisciplinary Healthcare, 2016, 9, 191.	1.1	6

#	Article	IF	CITATIONS
55	Psychometric Limitations of the Center for Epidemiologic Studies-Depression Scale for Assessing Depressive Symptoms among Adults with HIV/AIDS: A Rasch Analysis. Depression Research and Treatment, 2016, 2016, 1-11.	0.7	24
56	A Trend for Increased Risk of Revision Surgery due to Deep Infection following Fast-Track Hip Arthroplasty. Advances in Orthopedics, 2016, 2016, 1-7.	0.4	16
57	Factors Related to Postoperative Pain Trajectories following Total Knee Arthroplasty: A Longitudinal Study of Patients Admitted to a Russian Orthopaedic Clinic. Pain Research and Treatment, 2016, 2016, 1-12.	1.7	24
58	Preoperative Pain, Symptoms, and Psychological Factors related to Higher Acute Pain Trajectories during Hospitalization for Total Knee Arthroplasty. PLoS ONE, 2016, 11, e0161681.	1.1	38
59	Factors that can predict pain with walking, 12 months after total knee arthroplasty. Monthly Notices of the Royal Astronomical Society: Letters, 2016, 87, 600-606.	1.2	41
60	A Rasch Analysis of Assessments of Morning and Evening Fatigue in Oncology Patients Using the Lee Fatigue Scale. Journal of Pain and Symptom Management, 2016, 51, 1002-1012.	0.6	24
61	Sleep among bereaved caregivers of patients admitted to hospice: a 1-year longitudinal pilot study. BMJ Open, 2016, 6, e009345.	0.8	11
62	Pain and concomitant symptoms in medical and elective surgical inpatients: a point prevalence survey. Journal of Clinical Nursing, 2016, 25, 3229-3240.	1.4	8
63	Acute phase factors associated with the course of depression during the first 18 months after first-ever stroke. Disability and Rehabilitation, 2016, 38, 30-35.	0.9	15
64	The experiences of nurses implementing the Modified Early Warning Score and a 24-hour on-call Mobile Intensive Care Nurse: An exploratory study. Intensive and Critical Care Nursing, 2016, 34, 33-41.	1.4	24
65	The last three days of life: a comparison of pain management in the young old and the oldest old hospitalised patients using the <scp>R</scp> esident <scp>A</scp> ssessment <scp>I</scp> nstrument for <scp>P</scp> alliative <scp>C</scp> are. International Journal of Older People Nursing, 2015, 10, 263-272.	0.6	9
66	Morbid obese adults increased their sense of coherence 1 year after a patient education course: a longitudinal study. Journal of Multidisciplinary Healthcare, 2015, 8, 157.	1.1	10
67	Trajectories of illness perceptions in persons with chronic illness: An explorative longitudinal study. Journal of Health Psychology, 2015, 20, 942-953.	1.3	37
68	The Reliability and Validity of the Norwegian Version of the Quality of Life in Late-Stage Dementia Scale. Dementia and Geriatric Cognitive Disorders, 2015, 40, 233-242.	0.7	26
69	Cytokine polymorphisms and plasma levels are associated with sleep onset insomnia in adults living with HIV/AIDS. Brain, Behavior, and Immunity, 2015, 47, 58-65.	2.0	27
70	Factors associated with self-esteem in persons with morbid obesity and in persons with chronic obstructive pulmonary disease: A cross-sectional study. Psychology, Health and Medicine, 2015, 20, 431-442.	1.3	15
71	Fatigue after Stroke – Perspectives and Future Directions. International Journal of Stroke, 2015, 10, 280-281.	2.9	18
72	Circadian regulation gene polymorphisms are associated with sleep disruption and duration, and circadian phase and rhythm in adults with HIV. Chronobiology International, 2015, 32, 1278-1293.	0.9	22

#	Article	IF	CITATIONS
73	A 1-year follow-up study exploring the associations between perception of illness and health-related quality of life in persons with chronic obstructive pulmonary disease. International Journal of COPD, 2014, 9, 41.	0.9	23
74	The relationships of self-efficacy, physical activity, and paid work to health-related quality of life among patients with chronic obstructive pulmonary disease (COPD). Journal of Multidisciplinary Healthcare, 2014, 7, 239.	1.1	30
75	Sleep in family caregivers of patients admitted to hospice: A pilot study. Palliative and Supportive Care, 2014, 12, 439-444.	0.6	8
76	Trajectories of self-efficacy in persons with chronic illness: An explorative longitudinal study. Psychology and Health, 2014, 29, 350-364.	1.2	21
77	Fatigue in the acute phase after first stroke predicts poorer physical health 18 months later. Neurology, 2014, 82, 2255-2255.	1.5	2
78	A Review of the Relationship Between Poststroke Fatigue and Physical Activity. Topics in Geriatric Rehabilitation, 2014, 30, 296-306.	0.2	2
79	Subjective sleep quality in relation to objective sleep estimates: comparison, gender differences and changes between the acute phase and the sixâ€month followâ€up after stroke. Journal of Advanced Nursing, 2014, 70, 639-650.	1.5	19
80	Can the Fatigue Severity Scale 7-item version be used across different patient populations as a generic fatigue measure - a comparative study using a Rasch model approach. Health and Quality of Life Outcomes, 2014, 12, 24.	1.0	58
81	Rasch analysis of the sense of coherence scale in a sample of people with morbid obesity – a cross-sectional study. BMC Psychology, 2014, 2, .	0.9	14
82	Rasch analysis of the Beck Depression Inventory-II in stroke survivors: A cross-sectional study. Journal of Affective Disorders, 2014, 158, 48-52.	2.0	21
83	Cytokine polymorphisms are associated with fatigue in adults living with HIV/AIDS. Brain, Behavior, and Immunity, 2014, 40, 95-103.	2.0	36
84	Last three days of life in the hospital: a comparison of symptoms, signs and treatments in the young old and the oldest old patients using the Resident Assessment Instrument for Palliative Care. International Journal of Older People Nursing, 2013, 8, 199-206.	0.6	10
85	Development of a short version of the Lee Visual Analogue Fatigue Scale in a sample of women with HIV/AIDS: a Rasch analysis application. Quality of Life Research, 2013, 22, 1467-1472.	1.5	18
86	Fatigue in the acute phase after first stroke predicts poorer physical health 18 months later. Neurology, 2013, 81, 1581-1587.	1.5	50
87	Nurses' perceptions of unpleasant symptoms and signs in ventilated and sedated patients. Nursing in Critical Care, 2013, 18, 176-186.	1.1	22
88	Lee Fatigue and Energy Scales: Exploring aspects of validity in a sample of women with HIV using an application of a Rasch model. Psychiatry Research, 2013, 205, 241-246.	1.7	32
89	Lack of Energy: An Important and Distinct Component of HIV-Related Fatigue and Daytime Function. Journal of Pain and Symptom Management, 2013, 45, 191-201.	0.6	29
90	Rasch analysis of the General Self-Efficacy Scale in a sample of persons with morbid obesity. Health and Quality of Life Outcomes, 2013, 11, 202.	1.0	40

#	Article	IF	Citations
91	Depressive symptoms in acute stroke: A cross-sectional study of their association with sociodemographics and clinical factors. Neuropsychological Rehabilitation, 2013, 23, 658-677.	1.0	24
92	Pain characteristics and selfâ€rated health after elective orthopaedic surgery – a crossâ€sectional survey. Journal of Clinical Nursing, 2013, 22, 1242-1253.	1.4	24
93	Prescribed Exercise: A Prospective Study of Health-Related Quality of Life and Physical Fitness Among Participants in an Officially Sponsored Municipal Physical Training Program. Journal of Physical Activity and Health, 2013, 10, 1016-1023.	1.0	23
94	Curvilinear Relationship Between Age and Post-Stroke Fatigue among Patients in the Acute Phase following First-Ever Stroke. International Journal of Physical Medicine & Rehabilitation, 2013, 01, .	0.5	9
95	The Course of Fatigue during the First 18 Months after First-Ever Stroke: A Longitudinal Study. Stroke Research and Treatment, 2012, 2012, 1-8.	0.5	31
96	Relationships between Physical Activity, Symptoms and Quality of Life among Inpatients with Severe Mental Illness. British Journal of Occupational Therapy, 2012, 75, 69-75.	0.5	14
97	Changes in health-related quality of life in people with morbid obesity attending a learning and mastery course. A longitudinal study with 12-months follow-up. Health and Quality of Life Outcomes, 2012, 10, 95.	1.0	14
98	The visitor's regard of their need for support, comfort, information proximity and assurance in the intensive care unit. Intensive and Critical Care Nursing, 2012, 28, 263-268.	1.4	13
99	Using a didactic model to improve patient observation skills in neonatal intensive care nurse trainees $\hat{a} \in A$ pilot study. Intensive and Critical Care Nursing, 2012, 28, 208-214.	1.4	2
100	Agreement in documentation of symptoms, clinical signs, and treatment at the end of life: a comparison of data retrieved from nurse interviews and electronic patient records using the Resident Assessment Instrument for Palliative Care. Journal of Clinical Nursing, 2012, 21, 1416-1424.	1.4	11
101	Stroke patients' functions in personal activities of daily living in relation to sleep and socioâ€demographic and clinical variables in the acute phase after firstâ€time stroke and at six months of followâ€up. Journal of Clinical Nursing, 2012, 21, 1886-1895.	1.4	39
102	Factors associated with selfâ€efficacy in persons with chronic illness. Scandinavian Journal of Psychology, 2012, 53, 333-339.	0.8	60
103	Sleep-Wake Patterns during the Acute Phase after First-Ever Stroke. Stroke Research and Treatment, 2011, 2011, 1-7.	0.5	30
104	Group climate development in cognitive and interpersonal group therapy for social phobia Group Dynamics, 2011, 15, 32-48.	0.7	18
105	Patterns of morning and evening fatigue among adults with HIV/AIDS. Journal of Clinical Nursing, 2011, 20, 2204-2216.	1.4	25
106	Pain control at the end of life: a comparative study of hospitalized cancer and noncancer patients. Scandinavian Journal of Caring Sciences, 2011, 25, 771-779.	1.0	13
107	Psychometric properties of the Fatigue Severity Scaleâ€"Rasch analyses of individual responses in a Norwegian stroke cohort. International Journal of Nursing Studies, 2011, 48, 1258-1265.	2.5	128
108	Personal factors associated with health-related quality of life in persons with morbid obesity on treatment waiting lists in Norway. Quality of Life Research, 2011, 20, 1187-1196.	1.5	42

#	Article	IF	CITATIONS
109	A 7-item version of the fatigue severity scale has better psychometric properties among HIV-infected adults: an application of a Rasch model. Quality of Life Research, 2011, 20, 1447-1456.	1.5	41
110	Physical impairment, depressive symptoms and pre-stroke fatigue are related to fatigue in the acute phase after stroke. Disability and Rehabilitation, 2011, 33, 334-342.	0.9	116
111	A populationâ€based study of associations between current posttraumatic stress symptoms and current fatigue. Journal of Traumatic Stress, 2010, 23, 606-614.	1.0	19
112	Exploring person-centredness: a qualitative meta-synthesis of four studies. Scandinavian Journal of Caring Sciences, 2010, 24, 620-634.	1.0	170
113	Psychometric properties of the Fatigue Severity Scale: Rasch analyses of responses in a Norwegian and a Swedish MS cohort. Multiple Sclerosis Journal, 2010, 16, 733-741.	1.4	44
114	Poststroke Fatigue—A Review. Journal of Pain and Symptom Management, 2009, 38, 928-949.	0.6	199
115	The complexity of the relationship between chronic pain and quality of life: a study of the general Norwegian population. Quality of Life Research, 2009, 18, 971-980.	1.5	50
116	Stages of Change – Continuous Measure (URICA‣2): psychometrics of a Norwegian version. Journal of Advanced Nursing, 2009, 65, 193-202.	1.5	16
117	Perceptions of illness and its development in patients with multiple sclerosis: a prospective cohort study. Journal of Advanced Nursing, 2009, 65, 184-192.	1.5	17
118	Level of satisfaction during mammography screening in relation to discomfort, service provided, level of pain and breast compression. European Journal of Radiography, 2009, 1, 66-72.	0.2	15
119	Illness and injury presenting to a Norwegian travel insurance company's helpline. Travel Medicine and Infectious Disease, 2007, 5, 165-170.	1.5	7
120	A prospective study of patterns of fatigue in multiple sclerosis. European Journal of Neurology, 2007, 14, 1338-1343.	1.7	188
121	Age and the Experience of Chronic Pain. Clinical Journal of Pain, 2005, 21, 513-523.	0.8	177
122	Fatigue in the general population: A translation and test of the psychometric properties of the Norwegian version of the fatigue severity scale. Scandinavian Journal of Public Health, 2005, 33, 123-130.	1.2	294
123	Psychometric Properties of the WHOQOL-BREF Questionnaire for the Norwegian General Population. Journal of Nursing Measurement, 2004, 12, 147-159.	0.2	58
124	Quality of Life in the General Norwegian Population, Measured by the Quality of Life Scale (QOLS-N). Quality of Life Research, 2004, 13, 1001-1009.	1.5	62
125	Prevalence and characteristics of chronic pain in the general Norwegian population. European Journal of Pain, 2004, 8, 555-565.	1.4	192
126	Gender differences in chronic painâ€"findings from a population-based study of Norwegian adults. Pain Management Nursing, 2004, 5, 105-117.	0.4	99

## Anners Lerdal

#	Article	IF	CITATION
127	The Norwegian version of the Herth Hope Index (HHI-N): A psychometric study. Palliative and Supportive Care, 2004, 2, 255-263.	0.6	46
128	Fatigue and its association with sociodemographic variables among multiple sclerosis patients. Multiple Sclerosis Journal, 2003, 9, 509-514.	1.4	99
129	Hope in the general Norwegian population, measured using the Herth Hope Index. Palliative and Supportive Care, 2003, 1, 309-318.	0.6	27
130	A theoretical extension of the concept of energy through an empirical study. Scandinavian Journal of Caring Sciences, 2002, 16, 197-206.	1.0	36
131	A Concept Analysis of Energy: Its Meaning in the Lives of Three Individuals with Chronic Illness. Scandinavian Journal of Caring Sciences, 1998, 12, 3-10.	1.0	17
132	A Concept Analysis of Energy. Scandinavian Journal of Caring Sciences, 1998, 12, 3-10.	1.0	19