

Vincent P Klink

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5825814/publications.pdf>

Version: 2024-02-01

36
papers

1,233
citations

471509

17
h-index

395702

33
g-index

36
all docs

36
docs citations

36
times ranked

732
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Laser capture microdissection (LCM) and comparative microarray expression analysis of syncytial cells isolated from incompatible and compatible soybean (<i>Glycine max</i>) roots infected by the soybean cyst nematode (<i>Heterodera glycines</i>). <i>Planta</i> , 2007, 226, 1389-1409. | 3.2 | 154 |
| 2 | Laser Capture Microdissection (LCM) and Expression Analyses of <i>Glycine max</i> (Soybean) Syncytium Containing Root Regions Formed by the Plant Pathogen <i>Heterodera glycines</i> (Soybean Cyst Nematode). <i>Plant Molecular Biology</i> , 2005, 59, 965-979. | 3.9 | 114 |
| 3 | A correlation between host-mediated expression of parasite genes as tandem inverted repeats and abrogation of development of female <i>Heterodera glycines</i> cyst formation during infection of <i>Glycine max</i> . <i>Planta</i> , 2009, 230, 53-71. | 3.2 | 113 |
| 4 | A time-course comparative microarray analysis of an incompatible and compatible response by <i>Glycine max</i> (soybean) to <i>Heterodera glycines</i> (soybean cyst nematode) infection. <i>Planta</i> , 2007, 226, 1423-1447. | 3.2 | 111 |
| 5 | A gene expression analysis of syncytia laser microdissected from the roots of the <i>Glycine max</i> (soybean) genotype PI 548402 (Peking) undergoing a resistant reaction after infection by <i>Heterodera glycines</i> (soybean cyst nematode). <i>Plant Molecular Biology</i> , 2009, 71, 525-567. | 3.9 | 99 |
| 6 | Syncytium gene expression in <i>Glycine max</i> [PI 88788] roots undergoing a resistant reaction to the parasitic nematode <i>Heterodera glycines</i> . <i>Plant Physiology and Biochemistry</i> , 2010, 48, 176-193. | 5.8 | 74 |
| 7 | The expression of a naturally occurring, truncated allele of an $\hat{\pm}$ -SNAP gene suppresses plant parasitic nematode infection. <i>Plant Molecular Biology</i> , 2012, 80, 131-155. | 3.9 | 65 |
| 8 | Mapping cell fate decisions that occur during soybean defense responses. <i>Plant Molecular Biology</i> , 2011, 77, 513-528. | 3.9 | 59 |
| 9 | Differences in gene expression amplitude overlie a conserved transcriptomic program occurring between the rapid and potent localized resistant reaction at the syncytium of the <i>Glycine max</i> genotype Peking (PI 548402) as compared to the prolonged and potent resistant reaction of PI 88788. <i>Plant Molecular Biology</i> , 2011, 75, 141-165. | 3.9 | 48 |
| 10 | The mitogen activated protein kinase (MAPK) gene family functions as a cohort during the <i>Glycine max</i> defense response to <i>Heterodera glycines</i> . <i>Plant Physiology and Biochemistry</i> , 2019, 137, 25-41. | 5.8 | 44 |
| 11 | Syntaxin 31 functions in <i>Glycine max</i> resistance to the plant parasitic nematode <i>Heterodera glycines</i> . <i>Plant Molecular Biology</i> , 2014, 85, 107-121. | 3.9 | 39 |
| 12 | Microarray Detection Call Methodology as a Means to Identify and Compare Transcripts Expressed within Syncytial Cells from Soybean (<i>Glycine max</i>) Roots Undergoing Resistant and Susceptible Reactions to the Soybean Cyst Nematode (<i>Heterodera glycines</i>). <i>Journal of Biomedicine and Biotechnology</i> , 2010, 2010, 1-30. | 3.0 | 37 |
| 13 | A <i>Glycine max</i> homolog of NON-RACE SPECIFIC DISEASE RESISTANCE 1 (NDR1) alters defense gene expression while functioning during a resistance response to different root pathogens in different genetic backgrounds. <i>Plant Physiology and Biochemistry</i> , 2017, 114, 60-71. | 5.8 | 33 |
| 14 | A harpin elicitor induces the expression of a coiled-coil nucleotide binding leucine rich repeat (CC-NB-LRR) defense signaling gene and others functioning during defense to parasitic nematodes. <i>Plant Physiology and Biochemistry</i> , 2017, 121, 161-175. | 5.8 | 27 |
| 15 | Components of the SNARE-containing regulon are co-regulated in root cells undergoing defense. <i>Plant Signaling and Behavior</i> , 2017, 12, e1274481. | 2.4 | 25 |
| 16 | MiniMax, a new diminutive <i>Glycine max</i> genotype with a rapid life cycle, embryogenic potential and transformation capabilities. <i>Plant Cell, Tissue and Organ Culture</i> , 2008, 92, 183-195. | 2.3 | 24 |
| 17 | Co-regulation of the <i>Glycine max</i> soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptor (SNARE)-containing regulon occurs during defense to a root pathogen. <i>Journal of Plant Interactions</i> , 2016, 11, 74-93. | 2.1 | 24 |
| 18 | Xyloglucan endotransglycosylase/hydrolase increases tightly-bound xyloglucan and chain number but decreases chain length contributing to the defense response that <i>Glycine max</i> has to <i>Heterodera glycines</i> . <i>PLoS ONE</i> , 2021, 16, e0244305. | 2.5 | 19 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Exocyst components promote an incompatible interaction between <i>Glycine max</i> (soybean) and <i>Heterodera glycines</i> (the soybean cyst nematode). <i>Scientific Reports</i> , 2020, 10, 15003. | 3.3 | 18 |
| 20 | Harpin-inducible defense signaling components impair infection by the ascomycete <i>Macrophomina phaseolina</i> . <i>Plant Physiology and Biochemistry</i> , 2018, 129, 331-348. | 5.8 | 14 |
| 21 | A decline in transcript abundance for <i>Heterodera glycines</i> homologs of <i>Caenorhabditis elegans</i> uncoordinated genes accompanies its sedentary parasitic phase. <i>BMC Developmental Biology</i> , 2007, 7, 35. | 2.1 | 13 |
| 22 | The <i>Glycine max</i> Conserved Oligomeric Golgi (COG) Complex Functions During a Defense Response to <i>Heterodera glycines</i> . <i>Frontiers in Plant Science</i> , 2020, 11, 564495. | 3.6 | 11 |
| 23 | Mitogen activated protein kinase (MAPK)-regulated genes with predicted signal peptides function in the <i>Glycine max</i> defense response to the root pathogenic nematode <i>Heterodera glycines</i> . <i>PLoS ONE</i> , 2020, 15, e0241678. | 2.5 | 10 |
| 24 | An expanded role of the SNARE-containing regulon as it relates to the defense process that <i>Glycine max</i> has to <i>Heterodera glycines</i> . <i>Journal of Plant Interactions</i> , 2019, 14, 276-283. | 2.1 | 9 |
| 25 | The heterologous expression of a soybean (<i>Glycine max</i>) xyloglucan endotransglycosylase/hydrolase (XTH) in cotton (<i>Gossypium hirsutum</i>) suppresses parasitism by the root knot nematode <i>Meloidogyne incognita</i> . <i>PLoS ONE</i> , 2020, 15, e0235344. | 2.5 | 9 |
| 26 | A plant transformation system designed for high throughput genomics in <i>Gossypium hirsutum</i> to study root-organism interactions. <i>Journal of Plant Interactions</i> , 2015, 10, 11-20. | 2.1 | 8 |
| 27 | The heterologous expression of a <i>Glycine max</i> homolog of NONEXPRESSOR OF PR1 (NPR1) and β -hydroxynitrile glucosidase suppresses parasitism by the root pathogen <i>Meloidogyne incognita</i> in <i>Gossypium hirsutum</i> . <i>Journal of Plant Interactions</i> , 2016, 11, 41-52. | 2.1 | 7 |
| 28 | MAPKDB: A MAP kinase database for signal transduction element identification. <i>Bioinformatics</i> , 2019, 15, 338-341. | 0.5 | 6 |
| 29 | The impact of pRAP vectors on plant genetic transformation and pathogenesis studies including an analysis of <i>BRI1-ASSOCIATED RECEPTOR KINASE 1 (BAK1)</i> -mediated resistance. <i>Journal of Plant Interactions</i> , 2021, 16, 270-283. | 2.1 | 5 |
| 30 | Conserved oligomeric Golgi (COG) complex genes functioning in defense are expressed in root cells undergoing a defense response to a pathogenic infection and exhibit regulation by MAPKs. <i>PLoS ONE</i> , 2021, 16, e0256472. | 2.5 | 5 |
| 31 | Developing a systems biology approach to study disease progression caused by <i>Heterodera glycines</i> in <i>Glycine max</i> . <i>Gene Regulation and Systems Biology</i> , 2007, 1, 17-33. | 2.3 | 5 |
| 32 | The use of laser capture microdissection to study the infection of <i>Glycine max</i> (soybean) by <i>Heterodera glycines</i> (soybean cyst nematode). <i>Plant Signaling and Behavior</i> , 2008, 3, 105-107. | 2.4 | 3 |
| 33 | The conserved oligomeric Golgi (COG) complex, a window into plant-pathogen interactions. <i>Journal of Plant Interactions</i> , 2022, 17, 344-360. | 2.1 | 1 |
| 34 | <i>Glycine max</i> Homologs of DOESN'T MAKE INFECTIONS 1, 2, and 3 Function to Impair <i>Heterodera glycines</i> Parasitism While Also Regulating Mitogen Activated Protein Kinase Expression. <i>Frontiers in Plant Science</i> , 2022, 13, . | 3.6 | 0 |
| 35 | The central circadian regulator CCA1 functions in <i>Glycine max</i> during defense to a root pathogen, regulating the expression of genes acting in effector triggered immunity (ETI) and cell wall metabolism. <i>Plant Physiology and Biochemistry</i> , 2022, 185, 198-220. | 5.8 | 0 |
| 36 | The heterologous expression of conserved <i>Glycine max</i> (soybean) mitogen activated protein kinase 3 (MAPK3) paralogs suppresses <i>Meloidogyne incognita</i> parasitism in <i>Gossypium hirsutum</i> (upland) <i>Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 5</i> | | |