

Bert van Bavel

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/582842/bert-van-bavel-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

151
papers

7,923
citations

55
h-index

82
g-index

160
ext. papers

8,458
ext. citations

8
avg, IF

5.69
L-index

#	Paper	IF	Citations
151	NordAqua, a Nordic Center of Excellence to develop an algae-based photosynthetic production platform. <i>Physiologia Plantarum</i> , 2021 , 173, 507-513	4.6	2
150	Polymer Type Identification of Marine Plastic Litter Using a Miniature Near-Infrared Spectrometer (MicroNIR). <i>Applied Sciences (Switzerland)</i> , 2020 , 10, 8707	2.6	11
149	Relationships between serum-induced AhR bioactivity or mitochondrial inhibition and circulating polychlorinated biphenyls (PCBs). <i>Scientific Reports</i> , 2017 , 7, 9383	4.9	12
148	The effect of drinking water contaminated with perfluoroalkyl substances on a 10-year longitudinal trend of plasma levels in an elderly Uppsala cohort. <i>Environmental Research</i> , 2017 , 159, 95-102	7.9	19
147	Genetic and methylation variation in the CYP2B6 gene is related to circulating p,p'-dde levels in a population-based sample. <i>Environment International</i> , 2017 , 98, 212-218	12.9	4
146	Circulating levels of perfluoroalkyl substances (PFASs) and carotid artery atherosclerosis. <i>Environmental Research</i> , 2017 , 152, 157-164	7.9	25
145	The metabolic fingerprint of p,p'-DDE and HCB exposure in humans. <i>Environment International</i> , 2016 , 88, 60-66	12.9	47
144	Gas chromatography/atmospheric pressure chemical ionization/mass spectrometry for the analysis of organochlorine pesticides and polychlorinated biphenyls in human serum. <i>Journal of Chromatography A</i> , 2016 , 1453, 88-98	4.5	42
143	Hyperspectral Imaging and Data Analysis for Detecting and Determining Plastic Contamination in Seawater Filtrates. <i>Journal of Near Infrared Spectroscopy</i> , 2016 , 24, 141-149	1.5	49
142	Circulating levels of perfluoroalkyl substances are associated with dietary patterns - A cross sectional study in elderly Swedish men and women. <i>Environmental Research</i> , 2016 , 150, 59-65	7.9	22
141	Changes in serum levels of perfluoroalkyl substances during a 10-year follow-up period in a large population-based cohort. <i>Environment International</i> , 2016 , 95, 86-92	12.9	37
140	Perfluoroalkyl substances (PFAS) including structural PFOS isomers in plasma from elderly men and women from Sweden: Results from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). <i>Environment International</i> , 2015 , 82, 21-7	12.9	22
139	Genome-wide association study of plasma levels of polychlorinated biphenyls disclose an association with the CYP2B6 gene in a population-based sample. <i>Environmental Research</i> , 2015 , 140, 95-101	7.9	10
138	Atmospheric-pressure chemical ionization tandem mass spectrometry (APGC/MS/MS) an alternative to high-resolution mass spectrometry (HRGC/HRMS) for the determination of dioxins. <i>Analytical Chemistry</i> , 2015 , 87, 9047-53	7.8	52
137	Rapid separation of hexabromocyclododecane diastereomers using a novel method combining convergence chromatography and tandem mass spectrometry. <i>Analytical Methods</i> , 2015 , 7, 2950-2958	3.2	10
136	Exposure and body burden of polychlorinated biphenyls (PCB) and metals in a historically contaminated community. <i>Environment International</i> , 2015 , 76, 41-8	12.9	11
135	Circulating levels of environmental contaminants are associated with dietary patterns in older adults. <i>Environment International</i> , 2015 , 75, 93-102	12.9	13

134	Circulating levels of perfluoroalkyl substances and prevalent diabetes in the elderly. <i>Diabetologia</i> , 2014 , 57, 473-9	10.3	83
133	Persistent organic pollutants are related to the change in circulating lipid levels during a 5 year follow-up. <i>Environmental Research</i> , 2014 , 134, 190-7	7.9	31
132	Perfluoroalkyl substances in the blood samples from a male population of Sweden. <i>Science Bulletin</i> , 2014 , 59, 388-395		6
131	Persistent organic pollutants and liver dysfunction biomarkers in a population-based human sample of men and women. <i>Environmental Research</i> , 2014 , 134, 251-6	7.9	31
130	Case-control study on perfluorinated alkyl acids (PFAAs) and the risk of prostate cancer. <i>Environment International</i> , 2014 , 63, 35-9	12.9	45
129	Influence of persistent organic pollutants on the complement system in a population-based human sample. <i>Environment International</i> , 2014 , 71, 94-100	12.9	20
128	Genetic variation in the CYP2B6 gene is related to circulating 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) concentrations: an observational population-based study. <i>Environmental Health</i> , 2014 , 13, 34	6	8
127	In vitro bioassays for detecting dioxin-like activity--application potentials and limits of detection, a review. <i>Science of the Total Environment</i> , 2014 , 487, 37-48	10.2	67
126	Persistent organic pollutants distribution in lipoprotein fractions in relation to cardiovascular disease and cancer. <i>Environment International</i> , 2014 , 65, 93-9	12.9	56
125	Circulating levels of p,p'-DDE are related to prevalent hypertension in the elderly. <i>Environmental Research</i> , 2014 , 129, 27-31	7.9	42
124	The metabolically healthy but obese phenotype is associated with lower plasma levels of persistent organic pollutants as compared to the metabolically abnormal obese phenotype. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E1061-6	5.6	42
123	Genetic variation in the CYP1A1 gene is related to circulating PCB118 levels in a population-based sample. <i>Environmental Research</i> , 2014 , 133, 135-40	7.9	10
122	Influence of persistent organic pollutants on oxidative stress in population-based samples. <i>Chemosphere</i> , 2014 , 114, 303-9	8.4	27
121	Does mortality risk of cigarette smoking depend on serum concentrations of persistent organic pollutants? Prospective investigation of the vasculature in Uppsala seniors (PIVUS) study. <i>PLoS ONE</i> , 2014 , 9, e95937	3.7	8
120	Persistent organic pollutants and inflammatory markers in a cross-sectional study of elderly Swedish people: the PIVUS cohort. <i>Environmental Health Perspectives</i> , 2014 , 122, 977-83	8.4	37
119	The dioRAMA project: assessment of dioxin-like activity in sediments and fish (<i>Rutilus rutilus</i>) in support of the ecotoxicological characterization of sediments. <i>Journal of Soils and Sediments</i> , 2013 , 13, 770-774	3.4	7
118	An environmental wide association study (EWAS) approach to the metabolic syndrome. <i>Environment International</i> , 2013 , 55, 1-8	12.9	49
117	A rapid method for the determination of perfluoroalkyl substances including structural isomers of perfluorooctane sulfonic acid in human serum using 96-well plates and column-switching ultra-high performance liquid chromatography tandem mass spectrometry. <i>Journal of Chromatography A</i> , 2013 , 1285, 164-70	4.5	46

116	Professional ski waxers' exposure to PFAS and aerosol concentrations in gas phase and different particle size fractions. <i>Environmental Sciences: Processes and Impacts</i> , 2013 , 15, 814-22	4.3	17
115	Circulating levels of persistent organic pollutants are related to retrospective assessment of life-time weight change. <i>Chemosphere</i> , 2013 , 90, 998-1004	8.4	17
114	Biotransformation of fluorotelomer compound to perfluorocarboxylates in humans. <i>Environment International</i> , 2013 , 51, 8-12	12.9	51
113	Global DNA hypermethylation is associated with high serum levels of persistent organic pollutants in an elderly population. <i>Environment International</i> , 2013 , 59, 456-61	12.9	63
112	Perfluoroalkyl acids in subarctic wild male mink (<i>Neovison vison</i>) in relation to age, season and geographical area. <i>Environment International</i> , 2013 , 59, 425-30	12.9	25
111	Influence of age, season, body condition and geographical area on concentrations of chlorinated and brominated contaminants in wild mink (<i>Neovison vison</i>) in Sweden. <i>Chemosphere</i> , 2013 , 90, 1664-71	8.4	12
110	The need for capacity building and first results for the Stockholm Convention Global Monitoring Plan. <i>TrAC - Trends in Analytical Chemistry</i> , 2013 , 46, 72-84	14.6	34
109	Circulating levels of persistent organic pollutants (POPs) are associated with left ventricular systolic and diastolic dysfunction in the elderly. <i>Environmental Research</i> , 2013 , 123, 39-45	7.9	25
108	Circulating levels of persistent organic pollutants in relation to visceral and subcutaneous adipose tissue by abdominal MRI. <i>Obesity</i> , 2013 , 21, 413-8	8	39
107	Chemical and bioanalytical characterisation of PAHs in risk assessment of remediated PAH-contaminated soils. <i>Environmental Science and Pollution Research</i> , 2013 , 20, 8511-20	5.1	34
106	Capacity building for persistent organic pollutant (POP) analysis in the Pacific and POP trends in the Pacific Islands. <i>TrAC - Trends in Analytical Chemistry</i> , 2013 , 46, 173-177	14.6	3
105	Worldwide distribution of persistent organic pollutants in air, including results of air monitoring by passive air sampling in five continents. <i>TrAC - Trends in Analytical Chemistry</i> , 2013 , 46, 150-161	14.6	97
104	Persistent organic pollutants and abnormal geometry of the left ventricle in the elderly. <i>Journal of Hypertension</i> , 2013 , 31, 1547-53	1.9	14
103	Polychlorinated naphthalenes (PCNs) in sub-Arctic and Arctic marine mammals, 1986-2009. <i>Environmental Pollution</i> , 2012 , 164, 118-24	9.3	10
102	Methoxylated polybrominated diphenyl ethers (MeO-PBDEs) are major contributors to the persistent organobromine load in sub-Arctic and Arctic marine mammals, 1986-2009. <i>Science of the Total Environment</i> , 2012 , 416, 482-9	10.2	23
101	Polybrominated diphenyl ethers (PBDEs) in marine mammals from Arctic and North Atlantic regions, 1986-2009. <i>Environment International</i> , 2012 , 40, 102-109	12.9	57
100	Associations of persistent organic pollutants with abdominal obesity in the elderly: The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. <i>Environment International</i> , 2012 , 40, 170-178	12.9	99
99	Per- and polyfluorinated compounds (PFCs) in house dust and indoor air in Catalonia, Spain: implications for human exposure. <i>Environment International</i> , 2012 , 39, 172-80	12.9	91

98	Circulating levels of persistent organic pollutants (POPs) among elderly men and women from Sweden: results from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS). <i>Environment International</i> , 2012 , 44, 59-67	12.9	36
97	Background exposure to persistent organic pollutants predicts stroke in the elderly. <i>Environment International</i> , 2012 , 47, 115-20	12.9	56
96	Human dietary exposure to perfluoroalkyl substances in Catalonia, Spain. Temporal trend. <i>Food Chemistry</i> , 2012 , 135, 1575-82	8.5	97
95	Human exposure to perfluorinated compounds in Catalonia, Spain: contribution of drinking water and fish and shellfish. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 4408-15	5.7	72
94	Increasing levels of long-chain perfluorocarboxylic acids (PFCAs) in Arctic and North Atlantic marine mammals, 1984-2009. <i>Chemosphere</i> , 2012 , 86, 278-85	8.4	76
93	A rapid method for screening of the Stockholm Convention POPs in small amounts of human plasma using SPE and HRGC/HRMS. <i>Chemosphere</i> , 2012 , 86, 747-53	8.4	55
92	Circulating levels of persistent organic pollutants (POPs) and carotid atherosclerosis in the elderly. <i>Environmental Health Perspectives</i> , 2012 , 120, 38-43	8.4	83
91	An investigation of the co-variation in circulating levels of a large number of environmental contaminants. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2012 , 22, 476-82	6.7	19
90	Circulating levels of persistent organic pollutants associate in divergent ways to fat mass measured by DXA in humans. <i>Chemosphere</i> , 2011 , 85, 335-43	8.4	58
89	Polychlorinated biphenyls and organochlorine pesticides in plasma predict development of type 2 diabetes in the elderly: the prospective investigation of the vasculature in Uppsala Seniors (PIVUS) study. <i>Diabetes Care</i> , 2011 , 34, 1778-84	14.6	189
88	Inhalation exposure to fluorotelomer alcohols yield perfluorocarboxylates in human blood?. <i>Environmental Science & Technology</i> , 2010 , 44, 7717-22	10.3	48
87	Analysis of POPs in human samples reveal a contribution of brominated dioxin of up to 15% of the total dioxin TEQ. <i>Chemosphere</i> , 2010 , 78, 113-20	8.4	53
86	A time trend study of significantly elevated perfluorocarboxylate levels in humans after using fluorinated ski wax. <i>Environmental Science & Technology</i> , 2010 , 44, 2150-5	10.3	86
85	Biomonitoring perfluorinated compounds in Catalonia, Spain: concentrations and trends in human liver and milk samples. <i>Environmental Science and Pollution Research</i> , 2010 , 17, 750-8	5.1	117
84	Time trends of persistent organic pollutants in Sweden during 1993-2007 and relation to age, gender, body mass index, breast-feeding and parity. <i>Science of the Total Environment</i> , 2010 , 408, 4412-9	10.2	84
83	Levels in food and beverages and daily intake of perfluorinated compounds in Norway. <i>Chemosphere</i> , 2010 , 80, 1137-43	8.4	148
82	Tissue-specific contaminant accumulation and associated effects on hepatic serum analytes and cytochrome P450 enzyme activities in hooded seals (<i>Cystophora cristata</i>) from the Gulf of St. Lawrence. <i>Archives of Environmental Contamination and Toxicology</i> , 2009 , 56, 360-70	3.2	14
81	Accuracy and precision in the determination of perfluorinated chemicals in human blood verified by interlaboratory comparisons. <i>Journal of Chromatography A</i> , 2009 , 1216, 394-400	4.5	37

80	Polybrominated diphenyl ethers and their hydroxylated analogues in ringed seals (<i>Phoca hispida</i>) from Svalbard and the Baltic Sea. <i>Environmental Science & Technology</i> , 2009 , 43, 3494-9	10.3	68
79	Concentrations, patterns and metabolites of organochlorine pesticides in relation to xenobiotic phase I and II enzyme activities in ringed seals (<i>Phoca hispida</i>) from Svalbard and the Baltic Sea. <i>Environmental Pollution</i> , 2009 , 157, 2428-34	9.3	28
78	Levels of perfluorinated chemicals in municipal drinking water from Catalonia, Spain: public health implications. <i>Archives of Environmental Contamination and Toxicology</i> , 2009 , 57, 631-8	3.2	87
77	Concentrations of organohalogen compounds and titres of antibodies to Epstein-Barr virus antigens and the risk for non-Hodgkin lymphoma. <i>Oncology Reports</i> , 2009 , 21, 1567-76	3.5	30
76	Human exposure to perfluorinated chemicals through the diet: intake of perfluorinated compounds in foods from the Catalan (Spain) market. <i>Journal of Agricultural and Food Chemistry</i> , 2008 , 56, 1787-94	5.7	224
75	Novel evidence for natural formation of dioxins in ball clay. <i>Chemosphere</i> , 2008 , 70, 1280-9	8.4	42
74	Biotransformation of PCBs in relation to phase I and II xenobiotic-metabolizing enzyme activities in ringed seals (<i>Phoca hispida</i>) from Svalbard and the Baltic Sea. <i>Environmental Science & Technology</i> , 2008 , 42, 8952-8	10.3	78
73	Long-term worldwide QA/QC of dioxins and dioxin-like PCBs in environmental samples. <i>Analytical Chemistry</i> , 2008 , 80, 3956-64	7.8	18
72	Biomarker responses and decreasing contaminant levels in ringed seals (<i>Pusa hispida</i>) from Svalbard, Norway. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2008 , 71, 1009-18	3.8	22
71	Increased concentrations of certain persistent organic pollutants in subjects with self-reported electromagnetic hypersensitivity—a pilot study. <i>Electromagnetic Biology and Medicine</i> , 2008 , 27, 197-203	2.2	7
70	Activities and identification of aryl hydrocarbon receptor agonists in sediments from the Danube river. <i>Analytical and Bioanalytical Chemistry</i> , 2008 , 390, 2009-19	4.4	83
69	Levels of perfluorochemicals in water samples from Catalonia, Spain: is drinking water a significant contribution to human exposure?. <i>Environmental Science and Pollution Research</i> , 2008 , 15, 614-9	5.1	123
68	Changes in toxicity and Ah receptor agonist activity of suspended particulate matter during flood events at the rivers Neckar and Rhine - a mass balance approach using in vitro methods and chemical analysis. <i>Environmental Science and Pollution Research</i> , 2008 , 15, 536-53	5.1	80
67	Exposure of perfluorinated chemicals through lactation: levels of matched human milk and serum and a temporal trend, 1996-2004, in Sweden. <i>Environmental Health Perspectives</i> , 2007 , 115, 226-30	8.4	365
66	Degradation of polycyclic aromatic hydrocarbons (PAHs) in contaminated soils by Fenton's reagent: a multivariate evaluation of the importance of soil characteristics and PAH properties. <i>Journal of Hazardous Materials</i> , 2007 , 149, 86-96	12.8	133
65	Accumulation and transfer of contaminants in killer whales (<i>Orcinus orca</i>) from Norway: indications for contaminant metabolism. <i>Environmental Toxicology and Chemistry</i> , 2007 , 26, 1582-90	3.8	36
64	Relative differences in aryl hydrocarbon receptor-mediated response for 18 polybrominated and mixed halogenated dibenzo-p-dioxins and -furans in cell lines from four different species. <i>Environmental Toxicology and Chemistry</i> , 2007 , 26, 2448-54	3.8	89
63	Reed beds receiving industrial sludge containing nitroaromatic compounds. Effects of outgoing water and bed material extracts in the umu-c genotoxicity assay, DR-CALUX assay and on early life stage development in zebrafish (<i>Danio rerio</i>). <i>Environmental Science and Pollution Research</i> , 2007 , 14, 2002-11	5.1	38

62	Fractionation and Determination of Ah Receptor (AhR) Agonists in Organic Waste After Anaerobic Biodegradation and in Batch Experiments with PCB and decaBDE (8 pp). <i>Environmental Science and Pollution Research</i> , 2007 , 14 Suppl 1, 36-43	5.1	7
61	Levels of brominated flame retardants in blood in relation to levels in household air and dust. <i>Environment International</i> , 2007 , 33, 62-9	12.9	174
60	Perfluorinated chemicals in blood of residents in Catalonia (Spain) in relation to age and gender: a pilot study. <i>Environment International</i> , 2007 , 33, 616-23	12.9	126
59	Identification and pattern of perfluorooctane sulfonate (PFOS) isomers in human serum and plasma. <i>Environment International</i> , 2007 , 33, 782-8	12.9	93
58	Halogenated organic contaminants and mercury in northern fulmars (<i>Fulmarus glacialis</i>): levels, relationships to dietary descriptors and blood to liver comparison. <i>Environmental Pollution</i> , 2007 , 146, 25-33	9.3	38
57	Decreased survival in pancreatic cancer patients with high concentrations of organochlorines in adipose tissue. <i>Biomedicine and Pharmacotherapy</i> , 2007 , 61, 659-64	7.5	23
56	Degradation of toxaphene by <i>Bjerkandera</i> sp. strain BOL13 using waste biomass as a cosubstrate. <i>Applied Microbiology and Biotechnology</i> , 2006 , 71, 549-54	5.7	14
55	Levels of brominated flame retardants in Northern Fulmar (<i>Fulmarus glacialis</i>) eggs from the Faroe Islands. <i>Science of the Total Environment</i> , 2006 , 367, 840-6	10.2	47
54	Evaluation of total lipids using enzymatic methods for the normalization of persistent organic pollutant levels in serum. <i>Science of the Total Environment</i> , 2006 , 366, 361-6	10.2	94
53	Congener-specific accumulation and patterns of chlorinated and brominated contaminants in adult male walrus from Svalbard, Norway: indications for individual-specific prey selection. <i>Science of the Total Environment</i> , 2006 , 370, 70-9	10.2	25
52	Levels of 12 perfluorinated chemicals in pooled Australian serum, collected 2002-2003, in relation to age, gender, and region. <i>Environmental Science & Technology</i> , 2006 , 40, 3742-8	10.3	133
51	Struggle for quality in determination of perfluorinated contaminants in environmental and human samples. <i>Environmental Science & Technology</i> , 2006 , 40, 7854-60	10.3	110
50	Degradation of toxaphene in aged and freshly contaminated soil. <i>Chemosphere</i> , 2006 , 63, 609-15	8.4	5
49	Perfluorinated chemicals in relation to other persistent organic pollutants in human blood. <i>Chemosphere</i> , 2006 , 64, 1582-91	8.4	130
48	Chemical and toxicological characterisation of PBDFs from photolytic decomposition of decaBDE in toluene. <i>Environment International</i> , 2006 , 32, 851-7	12.9	35
47	Tissue-specific accumulation and lactational transfer of polychlorinated biphenyls, chlorinated pesticides, and brominated flame retardants in hooded seals (<i>Cistophora cristata</i>) from the Gulf of St. Lawrence: applications for monitoring. <i>Environmental Pollution</i> , 2006 , 142, 476-86	9.3	40
46	Adipose tissue concentrations of persistent organic pollutants and the risk of prostate cancer. <i>Journal of Occupational and Environmental Medicine</i> , 2006 , 48, 700-7	2	61
45	In utero exposure to persistent organic pollutants in relation to testicular cancer risk. <i>Journal of Developmental and Physical Disabilities</i> , 2006 , 29, 228-34		114

44	Ah receptor agonists in UV-exposed toluene solutions of decabromodiphenyl ether (decaBDE) and in soils contaminated with polybrominated diphenyl ethers (PBDEs). <i>Environmental Science and Pollution Research</i> , 2006 , 13, 161-9	5.1	15
43	Comparison of Fenton's Reagent and Ozone Oxidation of Polycyclic Aromatic Hydrocarbons in Aged Contaminated Soils (7 pp). <i>Journal of Soils and Sediments</i> , 2006 , 6, 208-214	3.4	41
42	High concentrations of organochlorines in a patient with kidney cancer and anorexia-cachexia syndrome. <i>Medicinal Chemistry</i> , 2006 , 2, 607-10	1.8	6
41	Reply to Comment on: Personal air sampling and analysis of polybrominated diphenyl ethers and other bromine containing compounds at an electronics recycling facility in Sweden by M. L. Hardy, JEM, 2005, 7, DOI: 10.1039/b418857h. <i>Journal of Environmental Monitoring</i> , 2005 , 7, 644		
40	Development of a solid-phase extraction-HPLC/single quadrupole MS method for quantification of perfluorochemicals in whole blood. <i>Analytical Chemistry</i> , 2005 , 77, 864-70	7.8	80
39	Dioxin emissions after installation of a polishing wet scrubber in a hazardous waste incineration facility. <i>Chemosphere</i> , 2005 , 61, 405-12	8.4	38
38	A toxaphene-degrading bacterium related to <i>Enterobacter cloacae</i> , strain D1 isolated from aged contaminated soil in Nicaragua. <i>Systematic and Applied Microbiology</i> , 2005 , 28, 632-9	4.2	11
37	Distribution of brominated flame retardants in different dust fractions in air from an electronics recycling facility. <i>Science of the Total Environment</i> , 2005 , 350, 151-60	10.2	58
36	Polybrominated diphenyl ethers--plasma levels and thyroid status of workers at an electronic recycling facility. <i>International Archives of Occupational and Environmental Health</i> , 2005 , 78, 584-92	3.2	59
35	Concentrations of polychlorinated biphenyls in blood and the risk for testicular cancer. <i>Journal of Developmental and Physical Disabilities</i> , 2004 , 27, 282-90		58
34	Adipose tissue concentrations of p,p'-DDE and the risk for endometrial cancer. <i>Gynecologic Oncology</i> , 2004 , 95, 706-11	4.9	31
33	Polybrominated diphenylethers and methoxylated tetrabromodiphenylethers in cetaceans from the Mediterranean Sea. <i>Archives of Environmental Contamination and Toxicology</i> , 2004 , 47, 542-50	3.2	68
32	Personal air sampling and analysis of polybrominated diphenyl ethers and other bromine containing compounds at an electronic recycling facility in Sweden. <i>Journal of Environmental Monitoring</i> , 2004 , 6, 874-80		57
31	Congener-specific accumulation and food chain transfer of polybrominated diphenyl ethers in two arctic food chains. <i>Environmental Science & Technology</i> , 2004 , 38, 1667-74	10.3	144
30	Analytical challenges hamper perfluoroalkyl research. <i>Environmental Science & Technology</i> , 2004 , 38, 248A-255A	10.3	167
29	Increased concentrations of polychlorinated biphenyls, hexachlorobenzene, and chlordanes in mothers of men with testicular cancer. <i>Environmental Health Perspectives</i> , 2003 , 111, 930-4	8.4	157
28	Levels and patterns of polycyclic aromatic hydrocarbons in incineration ashes. <i>Science of the Total Environment</i> , 2003 , 311, 221-31	10.2	63
27	Time trends of selected persistent organic pollutants in lake sediments from Greenland. <i>Environmental Science & Technology</i> , 2003 , 37, 4319-24	10.3	41

26	Multivariate characterization of polycyclic aromatic hydrocarbons using semi-empirical molecule orbital calculations and physical data. <i>Chemosphere</i> , 2003 , 50, 627-37	8.4	14
25	Polycyclic aromatic hydrocarbons in weathered bottom ash from incineration of municipal solid waste. <i>Chemosphere</i> , 2003 , 53, 123-8	8.4	37
24	Flux estimates and sedimentation of polychlorinated naphthalenes in the northern part of the Baltic Sea. <i>Environmental Pollution</i> , 2003 , 126, 93-105	9.3	26
23	Development of a high-performance liquid chromatography carbon column based method for the fractionation of dioxin-like polychlorinated biphenyls. <i>Journal of Chromatography A</i> , 2002 , 962, 79-93	4.5	19
22	Biological and chemical determination of dioxin-like compounds in sediments by means of a sediment triad approach in the catchment area of the river Neckar. <i>Ecotoxicology</i> , 2002 , 11, 323-36	2.9	62
21	Polychlorinated naphthalene levels, distribution, and biomagnification in a benthic food chain in the Baltic Sea. <i>Environmental Science & Technology</i> , 2002 , 36, 5005-13	10.3	36
20	Is DDT exposure during fetal period and breast-feeding associated with neurological impairment?. <i>Environmental Research</i> , 2002 , 88, 141-4	7.9	24
19	Case-control study on concentrations of organohalogen compounds and titers of antibodies to Epstein-Barr virus antigens in the etiology of non-Hodgkin lymphoma. <i>Leukemia and Lymphoma</i> , 2001 , 42, 619-29	1.9	75
18	Adipose tissue concentrations of dioxins and dibenzofurans, titers of antibodies to Epstein-Barr virus early antigen and the risk for non-Hodgkin lymphoma. <i>Environmental Research</i> , 2001 , 87, 99-107	7.9	26
17	Analytical techniques for the determination of POPs. <i>European Journal of Lipid Science and Technology</i> , 2000 , 102, 50-52	3	1
16	Pressurised liquid extraction of polycyclic aromatic hydrocarbons from contaminated soils. <i>Journal of Chromatography A</i> , 2000 , 883, 151-62	4.5	86
15	Potential pathways and exposure to explain the human body burden of organochlorine compounds: a multivariate statistical analysis of human monitoring in Wetzburg, Germany. <i>Chemosphere</i> , 2000 , 40, 1075-82	8.4	10
14	Multivariate data evaluation of PCB and dioxin profiles in the general population in Sweden and Spain. <i>Chemosphere</i> , 2000 , 40, 1083-8	8.4	41
13	Organochlorine compounds in the Gulf of Bothnia: sediment and benthic species. <i>Chemosphere</i> , 2000 , 40, 1205-11	8.4	21
12	Pressurised hot water/steam extraction of polychlorinated dibenzofurans and naphthalenes from industrial soil. <i>Analyst</i> , 1999 , 124, 1351-1354	5	44
11	. <i>Environmental Toxicology and Chemistry</i> , 1999 , 18, 1157	3.8	2
10	Results from the second round of the international intercalibration on PCDDs, PCDFs and PCBs in a fly ash extract. <i>Chemosphere</i> , 1998 , 37, 2077-2089	8.4	5
9	Concentrations and spatial variations of cyclodienes and other organochlorines in herring and perch from the Baltic Sea. <i>Science of the Total Environment</i> , 1998 , 215, 69-83	10.2	36

8	Concentrations, biomagnification and spatial variation of organochlorine compounds in a pelagic food web in the northern part of the Baltic Sea. <i>Science of the Total Environment</i> , 1998 , 217, 143-54	10.2	71
7	Some Aspects of the Etiology of Non-Hodgkin's Lymphoma. <i>Environmental Health Perspectives</i> , 1998 , 106, 679	8.4	5
6	Occurrence, Sedimentation, and Spatial Variations of Organochlorine Contaminants in Settling Particulate Matter and Sediments in the Northern Part of the Baltic Sea. <i>Environmental Science & Technology</i> , 1998 , 32, 1754-1759	10.3	102
5	Development of a Solid Phase Carbon Trap for Simultaneous Determination of PCDDs, PCDFs, PCBs, and Pesticides in Environmental Samples Using SFE-LC. <i>Analytical Chemistry</i> , 1996 , 68, 1279-83	7.8	55
4	Levels of PCBs in the aquatic environment of the Gulf of Bothnia: Benthic species and sediments. <i>Marine Pollution Bulletin</i> , 1996 , 32, 210-218	6.7	29
3	Supercritical fluid extraction of PCBs from human adipose tissue for HRGC/LRMS analysis. <i>Chemosphere</i> , 1995 , 30, 1229-36	8.4	20
2	Influence of combustion parameters on the formation of polychlorinated dibenzo-p-dioxins, dibenzofurans, benzenes, and biphenyls and polyaromatic hydrocarbons in a pilot incinerator. <i>Environmental Science & Technology</i> , 1993 , 27, 1602-1610	10.3	104
1	Circulating Levels of Persistent Organic Pollutants in Relation to Visceral and Subcutaneous Adipose Tissue by Abdominal MRI. <i>Obesity</i> ,	8	3