Dino Gnecco

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5830352/publications.pdf

Version: 2024-02-01

567144 610775 84 776 15 24 h-index citations g-index papers 96 96 96 837 citing authors all docs docs citations times ranked

#	Article	IF	CITATIONS
1	Diastereoselective synthesis of new zwitterionic bicyclic lactams, scaffolds for construction of 2-substituted-4-hydroxy piperidine and its pipecolic acid derivatives. RSC Advances, 2022, 12, 4187-4190.	1.7	O
2	Curcumin induces cortico-hippocampal neuronal reshaping and memory improvements in aged mice. Journal of Chemical Neuroanatomy, 2022, 121, 102091.	1.0	7
3	Diastereoselective Functionalization of Chiral Nâ€Acylâ€1,3â€oxazolidines and Their Applications in the Synthesis of Bioactive Molecules. European Journal of Organic Chemistry, 2022, 2022, .	1.2	3
4	Synthesis of (+)- and (\hat{a}^{\prime})-Geissman-Waiss lactone from chiral sulfonium salts. Tetrahedron Letters, 2020, 61, 151697.	0.7	3
5	Stereoconvergent synthesis of N-Boc-(2R,3S)-3-hydroxy-2-phenylpiperidine. Tetrahedron Letters, 2019, 60, 820-824.	0.7	1
6	Diastereoselective Synthesis of 3-Alkylindoloquinolizine Derivatives via Regiospecific Oxidative Cyclization. Heterocycles, 2019, 98, 509.	0.4	4
7	Divergent Synthesis of 5,6- and 3,6-Dihydropyridin-2(1H)-one via Intramolecular Knoevenagel Condensation. Heterocycles, 2019, 98, 96.	0.4	O
8	Efficient Synthesis of Chiral 5-Methoxycarbonylpyridin-2(1H)-ones and 3-Bromo-5-methoxycarbonylpyridin-2(1H)-ones. Heterocycles, 2019, 98, 215.	0.4	0
9	Highly Regioselective Ring Opening of a Common N,N-Dialkylaziridinium Ion by Carboxylic Acids. Heterocycles, 2018, 96, 219.	0.4	1
10	Diastereospecific Intramolecular Cyclopropanation of Enantiopure 8-Bromo-3-phenylhexahydrooxazolo[3,2-a]pyridin-5-ones. Heterocycles, 2018, 96, 152.	0.4	1
11	Preparation of Chiral \hat{I}^2 -Enamino Esters from Methyl Propiolate: Synthesis of Chiral Methyl 1-Substituted 6-Oxo-1,4,5,6-tetrahydropyridine-3-carboxylates. Heterocycles, 2018, 96, 895.	0.4	1
12	Curcuma treatment prevents cognitive deficit and alteration of neuronal morphology in the limbic system of aging rats. Synapse, 2017, 71, e21952.	0.6	30
13	Asymmetric Tandem Conjugate Addition–Aldol Condensation with <i>N</i> à€Acryloyloxazolidines Derived from 2â€Phenylglycinol. Asian Journal of Organic Chemistry, 2017, 6, 67-70.	1.3	8
14	The influence of sulfur configuration in ¹ H NMR chemical shifts of diasteromeric fiveâ€membered cyclic sulfites. Magnetic Resonance in Chemistry, 2017, 55, 233-238.	1.1	1
15	Retro-Curcuminoids as Mimics of Dehydrozingerone and Curcumin: Synthesis, NMR, X-ray, and Cytotoxic Activity. Molecules, 2017, 22, 33.	1.7	9
16	Chiral cyclic zwitterionic bipyridinium-4-olates for the diastereoselective synthesis of (R,S)- and (S,R)-trozamicol. Tetrahedron Letters, 2016, 57, 1683-1686.	0.7	3
17	Crystal structures of two chiral piperidine derivatives: $1-[(1R)-2-hydroxy-1-phenylethyl]$ piperidin-4-one and $8-[(1S)-1-phenylethyl]-1,4-dioxa-8-azaspiro[4.5]$ decane-7-thione. Acta Crystallographica Section E: Crystallographic Communications, 2015, 71, 1207-1211.	0.2	O
18	Diasterospecific Etherification and Diasteroselective Monobromination of (R)-(â€")-1-(2-Hydroxy-1-phenylethyl)-3,4-dihydropyridin-2(1H)-one. Heterocycles, 2015, 91, 1042.	0.4	1

#	Article	IF	CITATIONS
19	7-endo cyclization of 2,3-epoxyamides and 2,3-aziridine carboxamides by intramolecular Friedel–Crafts reaction. Tetrahedron: Asymmetry, 2015, 26, 95-101.	1.8	12
20	Oxidation and Aromatization of the Enantiopure Piperidine Derived from (R)-(-)-2-Phenylglycinol to (1'R)-(-)-1-(2'-Hydroxy-1'-phenylethyl)-1H-pyridin-2-one. Heterocycles, 2014, 89, 725.	0.4	4
21	Investigation of Three Diasteromeric Chalcone Epoxides Derivatives by NMR Spectroscopy and X-ray Crystallography. Journal of Chemical Crystallography, 2014, 44, 512-519.	0.5	2
22	Diastereoselective Approach to <i>cis</i> -4-Methyl/thiol-Pipecolic Esters Based on RCM Reaction and Conjugate Michael Addition. Synthetic Communications, 2014, 44, 2838-2847.	1.1	7
23	Oxazolidine Sulfur Ylides Derived from Phenylglycinol for the Specific and Highly Diastereoselective Synthesis of Aryl and Alkyl <i>trans</i> \$\text{i} \times \text{\text{E}} \text{poxyamides.} European Journal of Organic Chemistry, 2013, 2013, 5561-5565.	1.2	15
24	Synthesis of the indoloazocine derivatives from a chiral indol amide-stabilized sulfur ylide. Tetrahedron Letters, 2013, 54, 2729-2732.	0.7	9
25	4-Hydroxy-1,1′-bis[(S)-1-phenylethyl]-5,5′,6,6′-tetrahydro-3,4′-bipyridine-2,2′(1H,1′H)-dione. Acta Crystallographica Section E: Structure Reports Online, 2013, 69, o408-o409.	³ 0.2	1
26	Diastereoselective synthesis of aryl and alkyl trans-glycidic amides from pseudoephedrine-derived sulfonium salt. Chemospecific exo-tet ring closure for morpholin-3-ones. Tetrahedron, 2012, 68, 10252-10256.	1.0	11
27	¹ H and ¹³ C NMR characterization of new cycloartane triterpenes from <i>Mangifera indica</i> . Magnetic Resonance in Chemistry, 2012, 50, 52-57.	1.1	51
28	Acetate Bridged Trinuclear Zn, Ca and Mg Metal Complexes with 2- and 4-Substituted Pyridines. Journal of Chemical Crystallography, 2012, 42, 794-802.	0.5	8
29	Diastereoselective arylation of enantiopure 3-bromopiperidin-2-one derived from (R)-(â°')-2-phenylglycinol with organocuprate reagents. Tetrahedron Letters, 2011, 52, 5947-5950.	0.7	9
30	Chemodivergent Synthesis of 7-Aryl/alkyl-6-hydroxy-1,4-oxazepan-5-ones and 2-[Aryl/alkyl(hydroxy)methyl]morpholin-3-ones from a Common Epoxyamide Precursor. Synthesis, 2011, 2011, 2310-2320.	1.2	1
31	(1′S,2R,3R)-(â^²)-2-Hydroxy-3-morpholino-3-phenyl-N-(1′-phenylethyl)propionamide. Acta Crystallographica Section E: Structure Reports Online, 2009, 65, o743-o743.	0.2	0
32	Application of amide-stabilized sulfur ylide reactivity to the stereodivergent synthesis of (R,S)- and (S,R)-reboxetine. Tetrahedron: Asymmetry, 2009, 20, 2764-2768.	1.8	23
33	New cyclic zwitterionic building blocks for the synthesis of piperidine-2,4-dione and pyridine-2-one compounds. Tetrahedron Letters, 2009, 50, 4208-4211.	0.7	13
34	Crystal Structure of an Unexpected Derivative of Curcumin: 2-[2-(4-Acetoxy-3-methoxyphenyl)ethyl]benzothiazole. X-ray Structure Analysis Online, 2009, 25, 97-98.	0.1	1
35	A Short Synthesis of Indolizidine (+)-209B from (3R,6S,8AS)-(-)-6-Methyl-3-phenyl-hexahydrooxazolo[3,2-a]pyridin-5-one. Heterocycles, 2009, 78, 2589.	0.4	12
36	Preparation of (R)-(+)-3-Phenyl-2,3,5,6,7,8-hexahydrooxazolo[3,2-a]pyridin-4-ylium Bromide: Synthesis of (S)-(+)-Coniine, (R)-(-)-Coniceine and (R)-(+)-Anabasine. Heterocycles, 2007, 71, 2699.	0.4	20

#	Article	IF	CITATIONS
37	Reactivity of $(1\hat{a} \in S)$ -1- $(1\hat{a} \in S)$ -phenyl-ethyl)-4-hydroxy-piperidin-2-one with Lawesson's reagent. Journal of Sulfur Chemistry, 2007, 28, 239-243.	1.0	6
38	X-ray crystal structures of new chiral enaminones from 2,4-pentanedione and their heterocyclic derivatives. Journal of Chemical Crystallography, 2007, 37, 119-133.	0.5	1
39	Heterocyclic Derivatives of Curcumin: Crystal Structure of 3,5-Bis[.BETA(4-acetoxy-3-methoxyphenyl)ethyl]isoxazol. Analytical Sciences: X-ray Structure Analysis Online, 2006, 22, X165-X166.	0.1	0
40	N-Benzoyl-N,N′-dicyclohexylurea. Acta Crystallographica Section E: Structure Reports Online, 2006, 62, o2922-o2923.	0.2	5
41	Study of minimum energy conformers of N-substituted derivatives of piperidine and pyrrolidine. Evidence of weak H-bonding by theoretical correlation with experimental NMR data. Journal of Molecular Structure, 2006, 786, 53-64.	1.8	7
42	Regioselective Endocyclic Oxidation of Enantiopure 3â€Alkylpiperidines: Synthesis of (3\$,5\$)â€(â€)â€3â€Ethylâ€5â€Methylpiperidine. Synthetic Communications, 2006, 36, 935-942.	1.1	3
43	Crystal Structures and Synthesis of 5-Hydroxy-1,7-bis(4-hydroxy-3-methoxyphenyl)-hept-4,6-dien-3-one. Analytical Sciences: X-ray Structure Analysis Online, 2005, 21, X59-X60.	0.1	0
44	Efficient preparation of $(1\hat{a} \in ^2R)$ - (\hat{a}°) -1- $(2\hat{a} \in ^2$ -hydroxy- $1\hat{a} \in ^2$ -phenylethyl)piperidin-2-one: synthesis of $(2\hat{a} \in ^2S, 3R)$ - $(+)$ -stenusine. Tetrahedron: Asymmetry, 2005, 16, 949-952.	1.8	14
45	(–)-1-Benzyl-4-[1(S)-phenylethylamino]-5,6-dihydropyridin-2(1H)-one. Acta Crystallographica Section E: Structure Reports Online, 2005, 61, o2924-o2926.	0.2	1
46	(–)-1-Benzyl-4-[(1R)-2-hydroxy-1-phenylethylamino]-5,6-dihydropyridin-2(1H)-one. Acta Crystallographica Section E: Structure Reports Online, 2005, 61, o2927-o2929.	0.2	1
47	Synthesis and Structure of New Heterocyclic Derivatives of Curcumin. Heterocycles, 2005, 65, 49.	0.4	9
48	(1′R,3S)-2-Oxo-1-(1′-phenylethyl)piperidine-3-carboxylic acid: a case of a very strong intramolecular hydrogen bond. Acta Crystallographica Section E: Structure Reports Online, 2004, 60, o2110-o2112.	0.2	0
49	Syntheses of pyridin-4-ylium chirons: applications in a synthesis of (+)-coniine. Tetrahedron: Asymmetry, 2004, 15, 847-850.	1.8	13
50	Controlled reduction of 5-alkyl-3-phenyl-2,3,5,6,7,8-hexahydro-oxazolo[3,2-a]pyridin-4-ylium iodide: enantioselective synthesis of (â^2)-dihydropinidine and (+)-indolizidine 167B. Tetrahedron: Asymmetry, 2004, 15, 3393-3395.	1.8	23
51	Isolation and Characterization of Five New Tetrasaccharide Glycosides from the Roots oflpomoeastansand Their Cytotoxic Activity. Journal of Natural Products, 2004, 67, 1552-1556.	1.5	19
52	An Enantioselective Access to 1-Alkyl-1,2,3,4-tetrahydroisoquinolines. Application to a New Synthesis of (â^')-Argemonineâ€. Journal of Organic Chemistry, 2004, 69, 2737-2740.	1.7	33
53	Crystal Structure of the Curcumin Derivative, Acetic Acid 4-[7-(4-Acetoxy-3-methoxyphenyl)-3,5-dioxoheptyl]-2-methoxyphenyl ester. Analytical Sciences: X-ray Structure Analysis Online, 2004, 20, X91-X92.	0.1	0
54	Crystal Structure of {Acetic acid 4-[7-(4-acetoxy-3-methoxyphenyl)-3,5-dioxoheptyl]-2-methoxy ester-03.05}-boron difluoride: A Boron Complex of Acetylated Tetrahydrocurcumin Derivative. Analytical Sciences: X-ray Structure Analysis Online, 2004, 20, X167-X168.	0.1	1

#	Article	IF	CITATIONS
55	Solution1H and 13C NMR of new chiral 1,4-oxazepinium heterocycles and their intermediates from the reaction of 2,4-pentanedione with \hat{l}_{\pm} -L-amino acids and (R)-(-)-2-phenylglycinol. Magnetic Resonance in Chemistry, 2003, 41, 975-982.	1.1	4
56	(-)-(1′S,4aS,7R,8aR)-4a-Ethyl-7-hydroxy-1- (1′-phenylethyl)perhydroquinolinium Bromide ChemInform, 2003 34, no.	0.1	0
57	trans-(3R,2aS)-(â^')-3-Phenyl-2,3,5,6,7,8-hexahydro-oxazolo[3,2-a]pyridine-5-thione. Acta Crystallographica Section E: Structure Reports Online, 2003, 59, o519-o521.	0.2	3
58	Crystal Structure of trans(3R,2aS)-(-)-3-Phenyl-hexahydro-oxazolo[3,2-a]-pyridin-5-one. Analytical Sciences, 2003, 19, 1223-1224.	0.8	2
59	Synthesis and characterization of a new (phthalocyani-nato)bis(carboxylate) silicon(IV) compound with increased solubility. Journal of Porphyrins and Phthalocyanines, 2002, 06, 198-202.	0.4	11
60	(â€")-(1 â€ 2 S,4aS,7R,8aR)-4a-Ethyl-7-hydroxy-1-(1 â€ 2 -phenylethyl)perhydroquinolinium bromide. Acta Crystallographica Section C: Crystal Structure Communications, 2002, 58, o591-o592.	0.4	1
61	Crystal Structure of (+)-(R)-3-Methyl-1-(1'-phenyl-ethyl)-1H-pyridin-2-one Analytical Sciences, 2001, 17, 1247-1248.	0.8	1
62	Crystal Structure of (-)-(1'R)-1-(2'-Hydroxy-1'-phenyl-ethyl)-1H-pyridin-2-one Analytical Sciences, 2001, 17, 1139-1140.	0.8	1
63	New methodology for the synthesis of enantiopure (3R,2aR)-(â^')-3-phenyl-hexahydro-oxazolo[3,2-a]-pyridin-5-one: a synthesis of (S)-(+)-coniine. Tetrahedron: Asymmetry, 2001, 12, 357-360.	1.8	11
64	Synthesis of $(\hat{a}^{})$ - $(1\hat{a}\in ^2S, 4aS, 8aR)$ - and $(+)$ - $(1\hat{a}\in ^2S, 4aR, 8aS)$ -4a-ethyl-1- $(1\hat{a}\in ^2$ -phenylethyl)-octahydroquinolin-7-on Tetrahedron: Asymmetry, 2001, 12, 2099-2102.	es. 1.8	7
65	Unexpected retro-Michael reaction of (â^')-(1â€2S,4aS,8aR)- and (+)-(1â€2S,4aR,8aS)-4a-ethyl-1-(1-phenylethyl)octahydroquinolin-7-ones. Tetrahedron: Asymmetry, 2001, 12, 3209-3211.	1.8	7
66	Synthesis of New Chiral Aziridinoalcohols. Synthetic Communications, 2000, 30, 1303-1309.	1.1	4
67	Total assignment of the 1H and 13C NMR spectra of casimiroedine and its peracetylated derivative. Magnetic Resonance in Chemistry, 2000, 38, 366-369.	1.1	1
68	Regioselective Oxidation of 3-Substituted Pyridinium Salts. Molecules, 2000, 5, 1175-1181.	1.7	2
69	Regiospecific and Enantiospecific Ring Opening of Methyl (+)-(1'R, 2R)- and (-)-(1'R,) Tj ETQq1 1 0.784314 rgBT /C	Dverlock 1 1.7	.0 ₂₀ 50 182
70	The Zincke's Reaction: A New Alternative for the Preparation of L-[2-(3-Indol)Ethyl]-Alkylpyridinium Chloride Derivatives. Synthetic Communications, 1999, 29, 281-287.	1.1	8
71	Title is missing!. Journal of Chemical Crystallography, 1998, 28, 529-537.	0.5	9
72	A short enantioselective access to 2,3,6-trialkylpiperidines and 5,8-dialkylindolizidines. Tetrahedron, 1998, 54, 9357-9372.	1.0	14

#	Article	IF	CITATIONS
73	Oxidation of chiral non-racemic pyridinium salts to enantiopure 2-pyridone and 3-alkyl-2-pyridones. Tetrahedron: Asymmetry, 1998, 9, 2027-2029.	1.8	24
74	Comparison of crystal and solution structures and 1H and 13C chemical shifts for grandiflorenic acid, kaurenoic acid, and monoginoic acid. Canadian Journal of Chemistry, 1997, 75, 342-347.	0.6	18
75	Pyridine-Derived Oxazolidines as Chiral 3-Alkyl-4,5-dihydropyridinium and 3-Alkyl-3,4,5,6-tetrahydropyridinium Salt Equivalents. Journal of Organic Chemistry, 1997, 62, 729-733.	1.7	50
76	Synthesis of α-phenyl-1-(R)-(â^')-piperidineacetic esters. Tetrahedron: Asymmetry, 1997, 8, 203-206.	1.8	11
77	Synthesis of (2R,3S)-(â^')-2-phenyl-3-methylaziridine. Tetrahedron: Asymmetry, 1997, 8, 2877-2879.	1.8	18
78	AN IMPROVED PREPARATION OF 1-METHYL-4-CYANO-4-PHENYLPIPERIDINE. Organic Preparations and Procedures International, 1996, 28, 478-480.	0.6	1
79	The Unambiguous Detection of Kaurenic Derivatives in Aqueous Infusions of Montanoa tomentosaby GC-MS and 2D-NMR Spectroscopy: An Answer to Contradictory Reports. Planta Medica, 1996, 62, 569-571.	0.7	9
80	Detailed characterization by, two-dimensional NMR of two unusual bicyclo [2.2.2]octenedione derivatives produced by the reaction of perezone with thiourea. Magnetic Resonance in Chemistry, 1995, 33, 3-7.	1.1	5
81	The Reaction of Perezone and Isoperezone with Hydroxylamine: A Surprisingly Facile Method for Introducing an NH2Group into the Quinone Functionality. Natural Product Research, 1995, 6, 103-109.	0.4	5
82	1,4-dihydropyridines from dithionite reduction of pyridinium salts without electron-withdrawing groups as substituents. Tetrahedron Letters, 1994, 35, 707-710.	0.7	43
83	Zincke's Reaction with Chiral Primary Amines: A Practical Entry to Pyridinium Salts of Interest in Asymmetric Synthesis. Synlett, 1992, 1992, 431-434.	1.0	53
84	Asymmetric synthesis from pyridines: use of new chiral 1,4-dihydropyridines in a short synthesis of 5,8-disubstituted indolizidine (+)-209B. Journal of the Chemical Society Chemical Communications, 1991, , 625.	2.0	25