
Jun Okabe

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5839457/publications.pdf Version: 2024-02-01



LUN OKARE

#	Article	IF	CITATIONS
1	Macrophage ubiquitin-specific protease 2 contributes to motility, hyperactivation, capacitation, and in vitro fertilization activity of mouse sperm. Cellular and Molecular Life Sciences, 2021, 78, 2929-2948.	5.4	11
2	Epigenetic evidence of an Ac/Dc axis by VPA and SAHA. Clinical Epigenetics, 2021, 13, 58.	4.1	13
3	Branched-chain amino acids and l-carnitine attenuate lipotoxic hepatocellular damage in rat cirrhotic liver. Biomedicine and Pharmacotherapy, 2021, 135, 111181.	5.6	12
4	SAHA attenuates Takotsubo-like myocardial injury by targeting an epigenetic Ac/Dc axis. Signal Transduction and Targeted Therapy, 2021, 6, 159.	17.1	14
5	Valproic acid influences the expression of genes implicated with hyperglycaemia-induced complement and coagulation pathways. Scientific Reports, 2021, 11, 2163.	3.3	18
6	Targeting Treatment Refractory <i>NET</i> by EZH2 Inhibition in Postural Tachycardia Syndrome. Circulation Research, 2020, 126, 1058-1060.	4.5	5
7	The Set7 Lysine Methyltransferase Regulates Plasticity in Oxidative Phosphorylation Necessary for Trained Immunity Induced by β-Glucan. Cell Reports, 2020, 31, 107548.	6.4	76
8	Sex-Based Mhrt Methylation Chromatinizes MeCP2 in the Heart. IScience, 2019, 17, 288-301.	4.1	8
9	Systems approach to the pharmacological actions of HDAC inhibitors reveals EP300 activities and convergent mechanisms of regulation in diabetes. Epigenetics, 2017, 12, 991-1003.	2.7	20
10	NADPH Oxidase Nox5 Accelerates Renal Injury in Diabetic Nephropathy. Diabetes, 2017, 66, 2691-2703.	0.6	119
11	Ubiquitin-Specific Protease 2 Modulates the Lipopolysaccharide-Elicited Expression of Proinflammatory Cytokines in Macrophage-like HL-60 Cells. Mediators of Inflammation, 2017, 2017, 1-15.	3.0	23
12	NET silencing by let-7i in postural tachycardia syndrome. JCI Insight, 2017, 2, e90183.	5.0	17
13	Current perspectives in Set7 mediated stem cell differentiation. Non-coding RNA, 2016, 2, 14.	2.6	2
14	Pharmacological inhibition of arginine and lysine methyltransferases induces nuclear abnormalities and suppresses angiogenesis in human endothelial cells. Biochemical Pharmacology, 2016, 121, 18-32.	4.4	17
15	Set7 mediated interactions regulate transcriptional networks in embryonic stem cells. Nucleic Acids Research, 2016, 44, gkw621.	14.5	15
16	Reactive Oxygen Species Can Provide Atheroprotection via NOX4-Dependent Inhibition of Inflammation and Vascular Remodeling. Arteriosclerosis, Thrombosis, and Vascular Biology, 2016, 36, 295-307.	2.4	147
17	The primary microRNA-208b interacts with Polycomb-group protein, Ezh2, to regulate gene expression in the heart. Nucleic Acids Research, 2014, 42, 790-803.	14.5	57
18	Vascular histone deacetylation by pharmacological HDAC inhibition. Genome Research, 2014, 24, 1271-1284.	5.5	79

Jun Okabe

#	Article	IF	CITATIONS
19	Deep sequencing reveals novel Set7 networks. Cellular and Molecular Life Sciences, 2014, 71, 4471-4486.	5.4	26
20	Genetic Targeting or Pharmacologic Inhibition of NADPH Oxidase Nox4 Provides Renoprotection in Long-Term Diabetic Nephropathy. Journal of the American Society of Nephrology: JASN, 2014, 25, 1237-1254.	6.1	301
21	Endothelial Transcriptome in Response to Pharmacological Methyltransferase Inhibition. ChemMedChem, 2014, 9, 1755-1762.	3.2	8
22	NADPH Oxidase 1 Plays a Key Role in Diabetes Mellitus–Accelerated Atherosclerosis. Circulation, 2013, 127, 1888-1902.	1.6	325
23	Ubiquitinâ€specific protease 2â€69 in macrophages potentially modulates metainflammation. FASEB Journal, 2013, 27, 4940-4953.	0.5	31
24	Distinguishing Hyperglycemic Changes by Set7 in Vascular Endothelial Cells. Circulation Research, 2012, 110, 1067-1076.	4.5	147
25	Analysis of the IGF2/H19 imprinting control region uncovers new genetic defects, including mutations of OCT-binding sequences, in patients with 11p15 fetal growth disorders. Human Molecular Genetics, 2010, 19, 803-814.	2.9	106
26	Epigenetic phenomena linked to diabetic complications. Nature Reviews Endocrinology, 2010, 6, 665-675.	9.6	202
27	Trichostatin A accentuates doxorubicin-induced hypertrophy in cardiac myocytes. Aging, 2010, 2, 659-668.	3.1	42
28	Hyperglycemia Induces a Dynamic Cooperativity of Histone Methylase and Demethylase Enzymes Associated With Gene-Activating Epigenetic Marks That Coexist on the Lysine Tail. Diabetes, 2009, 58, 1229-1236.	0.6	468
29	Sendai Virus-Mediated Gene Delivery into Hepatocytes via Isolated Hepatic Perfusion. Biological and Pharmaceutical Bulletin, 2006, 29, 1728-1734.	1.4	7
30	Reply to "Testing for association between MeCP2 and the brahma-associated SWI/SNF chromatin-remodeling complex― Nature Genetics, 2006, 38, 964-967.	21.4	16
31	Limited capacity of the nuclear matrix to bind telomere repeat binding factor TRF1 may restrict the proliferation of mortal human fibroblasts. Human Molecular Genetics, 2003, 13, 285-293.	2.9	14
32	Basic Peptides as Functional Components of Non-viral Gene Transfer Vehicles. Current Protein and Peptide Science, 2003, 4, 141-150.	1.4	43
33	TRF1 is a critical trans-acting factor required for de novo telomere formation in human cells. Human Molecular Genetics, 2000, 9, 2639-2650.	2.9	33
34	Identification and Characterization of Cell Lines with a Defect in a Post-adsorption Stage of Sendai Virus-mediated Membrane Fusion. Journal of Biological Chemistry, 2000, 275, 17549-17555.	3.4	13
35	Gene delivery systems using the Sendai virus. Molecular Membrane Biology, 1999, 16, 123-127.	2.0	17
36	Gene transfer vectors based on Sendai virus. Journal of Controlled Release, 1998, 54, 61-68.	9.9	24