

Sha Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5839917/publications.pdf>

Version: 2024-02-01

52
papers

1,511
citations

361413

20
h-index

345221

36
g-index

52
all docs

52
docs citations

52
times ranked

1755
citing authors

#	ARTICLE	IF	CITATIONS
1	Vesicle trafficking in <i>Arabidopsis</i> pollen tubes. <i>FEBS Letters</i> , 2022, , .	2.8	5
2	<i>Arabidopsis</i> RAN GTPases are critical for mitosis during male and female gametogenesis. <i>FEBS Letters</i> , 2022, 596, 1892-1903.	2.8	1
3	A positive feedback circuit for ROP-mediated polar growth. <i>Molecular Plant</i> , 2021, 14, 395-410.	8.3	19
4	Spliceosome component JANUS fulfills a role of mediator in transcriptional regulation during <i>Arabidopsis</i> development. <i>Plant Signaling and Behavior</i> , 2021, 16, 1841974.	2.4	2
5	The <i>Arabidopsis</i> SNARE protein YKT61 is essential for gametophyte development. <i>Journal of Integrative Plant Biology</i> , 2021, 63, 676-694.	8.5	13
6	<i>Arabidopsis</i> CBP60b is a central transcriptional activator of immunity. <i>Plant Physiology</i> , 2021, 186, 1645-1659.	4.8	30
7	The canonical $\hat{\pm}$ -SNAP is essential for gametophytic development in <i>Arabidopsis</i> . <i>PLoS Genetics</i> , 2021, 17, e1009505.	3.5	9
8	Functions of plant importin $\hat{2}$ proteins beyond nucleocytoplasmic transport. <i>Journal of Experimental Botany</i> , 2021, 72, 6140-6149.	4.8	7
9	<i>Arabidopsis</i> ADP-RIBOSYLATION FACTORs mediate tapetum-controlled pollen development. <i>Plant Journal</i> , 2021, 108, 268-280.	5.7	3
10	<i>S</i> -acylation of CBL10/SCaBP8 by PAT10 is crucial for its tonoplast association and function in salt tolerance. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 718-722.	8.5	15
11	Transcriptional Regulation of <i>PLETHORA1</i> in the Root Meristem Through an Importin and Its Two Antagonistic Cargos. <i>Plant Cell</i> , 2020, 32, 3812-3824.	6.6	15
12	Nitrate deficiency induces differential endocytosis in roots through NRT1.1. <i>Plant Signaling and Behavior</i> , 2020, 15, 1794394.	2.4	1
13	UV RESISTANCE LOCUS8 mediates ultraviolet-B-induced stomatal closure in an ethylene-dependent manner. <i>Plant Science</i> , 2020, 301, 110679.	3.6	14
14	FERONIA mediates root nutating growth. <i>Plant Journal</i> , 2020, 104, 1105-1116.	5.7	13
15	NRT1.1-Mediated Nitrate Suppression of Root Coiling Relies on PIN2- and AUX1-Mediated Auxin Transport. <i>Frontiers in Plant Science</i> , 2020, 11, 671.	3.6	19
16	Targeting and signaling of Rho of plants guanosine triphosphatases require synergistic interaction between guanine nucleotide inhibitor and vesicular trafficking. <i>Journal of Integrative Plant Biology</i> , 2020, 62, 1484-1499.	8.5	11
17	<i>Arabidopsis</i> <i>KETCH1</i> Is Critical for the Nuclear Accumulation of Ribosomal Proteins and Gametogenesis. <i>Plant Cell</i> , 2020, 32, 1270-1284.	6.6	22
18	SF3b4: A Versatile Player in Eukaryotic Cells. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 14.	3.7	14

#	ARTICLE	IF	CITATIONS
19	COPII Components Sar1b and Sar1c Play Distinct Yet Interchangeable Roles in Pollen Development. <i>Plant Physiology</i> , 2020, 183, 974-985.	4.8	19
20	HUA ENHANCER1 Mediates Ovule Development. <i>Frontiers in Plant Science</i> , 2020, 11, 397.	3.6	9
21	Arabidopsis Chloroplast protein for Growth and Fertility1 (CGF1) and CGF2 are essential for chloroplast development and female gametogenesis. <i>BMC Plant Biology</i> , 2020, 20, 172.	3.6	4
22	Downregulating VAC14 in Guard Cells Causes Drought Hypersensitivity by Inhibiting Stomatal Closure. <i>Frontiers in Plant Science</i> , 2020, 11, 602701.	3.6	1
23	Arabidopsis JANUS Regulates Embryonic Pattern Formation through Pol II-Mediated Transcription of WOXP2 and PIN7. <i>IScience</i> , 2019, 19, 1179-1188.	4.1	20
24	Importin Î24 Mediates Nuclear Import of GRF-Interacting Factors to Control Ovule Development in Arabidopsis. <i>Plant Physiology</i> , 2019, 179, 1080-1092.	4.8	42
25	Free-base porphyrins as CEST MRI contrast agents with highly upfield shifted labile protons. <i>Magnetic Resonance in Medicine</i> , 2019, 82, 577-585.	3.0	14
26	Vacuolar trafficking in pollen tube growth and guidance. <i>Plant Signaling and Behavior</i> , 2018, 13, e1464854.	2.4	2
27	The ADAPTOR PROTEIN-3 Complex Mediates Pollen Tube Growth by Coordinating Vacuolar Targeting and Organization. <i>Plant Physiology</i> , 2018, 177, 216-225.	4.8	25
28	AGC1.5 Kinase Phosphorylates RopGEFs to Control Pollen Tube Growth. <i>Molecular Plant</i> , 2018, 11, 1198-1209.	8.3	43
29	A Tonoplast-Associated Calcium-Signaling Module Dampens ABA Signaling during Stomatal Movement. <i>Plant Physiology</i> , 2018, 177, 1666-1678.	4.8	47
30	Arabidopsis VAC14 Is Critical for Pollen Development through Mediating Vacuolar Organization. <i>Plant Physiology</i> , 2018, 177, 1529-1538.	4.8	22
31	Arabidopsis PROTEIN S-ACYL TRANSFERASE4 mediates root hair growth. <i>Plant Journal</i> , 2017, 90, 249-260.	5.7	31
32	Tonoplast targeting of VHA3 relies on a Rab5-mediated but Rab7-independent vacuolar trafficking route. <i>Journal of Integrative Plant Biology</i> , 2017, 59, 230-233.	8.5	22
33	Arabidopsis adaptor protein 1G is critical for pollen development. <i>Journal of Integrative Plant Biology</i> , 2017, 59, 594-599.	8.5	10
34	AP1G mediates vacuolar acidification during synergid-controlled pollen tube reception. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E4877-E4883.	7.1	22
35	Adaptor Protein-3-Dependent Vacuolar Trafficking Involves a Subpopulation of COPII and HOPS Tethering Proteins. <i>Plant Physiology</i> , 2017, 174, 1609-1620.	4.8	42
36	Protein S-acyl transferase 4 controls nucleus position during root hair tip growth. <i>Plant Signaling and Behavior</i> , 2017, 12, e1311438.	2.4	7

#	ARTICLE	IF	CITATIONS
37	Update on adaptor protein-3 in Arabidopsis. <i>Plant Signaling and Behavior</i> , 2017, 12, e1356969.	2.4	2
38	Reactive oxygen species mediate tapetal programmed cell death in tobacco and tomato. <i>BMC Plant Biology</i> , 2017, 17, 76.	3.6	48
39	GPS in pollen tubes: Found!. <i>Science China Life Sciences</i> , 2016, 59, 438-439.	4.9	1
40	Precocious leaf senescence by functional loss of PROTEIN S-ACYL TRANSFERASE14 involves the NPR1-dependent salicylic acid signaling. <i>Scientific Reports</i> , 2016, 6, 20309.	3.3	32
41	<i>PLURIPETALA</i> mediates <i>ROP</i> 2 localization and stability in parallel to <i>SCN</i> 1 but synergistically with <i>TIP</i> 1 in root hairs. <i>Plant Journal</i> , 2016, 86, 413-425.	5.7	25
42	Arabidopsis RhoGDIs Are Critical for Cellular Homeostasis of Pollen Tubes. <i>Plant Physiology</i> , 2016, 170, 841-856.	4.8	43
43	HAPLESS13-Mediated Trafficking of STRUBBELIG Is Critical for Ovule Development in Arabidopsis. <i>PLoS Genetics</i> , 2016, 12, e1006269.	3.5	36
44	Protein palmitoylation is critical for the polar growth of root hairs in Arabidopsis. <i>BMC Plant Biology</i> , 2015, 15, 50.	3.6	32
45	To Grow or Not to Grow: FERONIA Has Her Say. <i>Molecular Plant</i> , 2014, 7, 1261-1263.	8.3	8
46	Spatiotemporal Production of Reactive Oxygen Species by NADPH Oxidase Is Critical for Tapetal Programmed Cell Death and Pollen Development in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2014, 26, 2007-2023.	6.6	247
47	Arabidopsis <i>Rop</i> GEF4 and <i>Rop</i> GEF10 are important for <i>FERONIA</i> -mediated developmental but not environmental regulation of root hair growth. <i>New Phytologist</i> , 2013, 200, 1089-1101.	7.3	81
48	Arabidopsis <i>COBRA</i> - <i>LIKE</i> 10, a GPI-anchored protein, mediates directional growth of pollen tubes. <i>Plant Journal</i> , 2013, 74, 486-497.	5.7	105
49	HAPLESS13, the Arabidopsis μ 1 Adaptin, Is Essential for Protein Sorting at the trans-Golgi Network/Early Endosome. <i>Plant Physiology</i> , 2013, 162, 1897-1910.	4.8	77
50	The juxtamembrane and carboxy-terminal domains of Arabidopsis PRK2 are critical for ROP-induced growth in pollen tubes. <i>Journal of Experimental Botany</i> , 2013, 64, 5599-5610.	4.8	30
51	PROTEIN S-ACYL TRANSFERASE10 Is Critical for Development and Salt Tolerance in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2013, 25, 1093-1107.	6.6	131
52	The C-Terminal Hypervariable Domain Targets Arabidopsis ROP9 to the Invaginated Pollen Tube Plasma Membrane. <i>Molecular Plant</i> , 2013, 6, 1362-1364.	8.3	18