

# Vidal de Freitas Mansano

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5843360/publications.pdf>

Version: 2024-02-01

93

papers

2,618

citations

623734

14

h-index

214800

47

g-index

94

all docs

94

docs citations

94

times ranked

2794

citing authors

#	ARTICLE	IF	CITATIONS
1	Untangling nomenclatural issues of some Amazonian trees of <i>Eperua</i> Aubl. (Leguminosae) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	0.3	
2	Brazilian Flora 2020: Leveraging the power of a collaborative scientific network. <i>Taxon</i> , 2022, 71, 178-198.	0.7	68
3	Evolution of the Anther Gland in Early-Branching Papilionoids (ADA Clade, Papilioideae,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	3.5	2
4	A Taxonomic Revision of the Amazonian Genus <i>Dicorynia</i> (Fabaceae: Dialioideae). <i>Phytotaxa</i> , 2022, 554, 1-31.	0.3	0
5	Bloodwood: the composition and secreting-site of the characteristic red exudate that gives the name to the <i>Swartzia</i> species (Fabaceae). <i>Journal of Plant Research</i> , 2021, 134, 127-139.	2.4	2
6	Resolving the non- <i>papilionaceous</i> flower of <i>Camoensia scandens</i> , a papilionoid legume of the core genistoid clade: development, glands and insights into the pollination and systematics of the group. <i>Journal of Plant Research</i> , 2021, 134, 823-839.	2.4	3
7	(2818) Proposal to conserve the name <i>Ficus trigona</i> (<i>Moraceae</i>) with a conserved type. <i>Taxon</i> , 2021, 70, 678-679.	0.7	1
8	Typification of <i>Ficus</i> sect. <i>Americanae</i> (Moraceae): <i>F. aurea</i> and <i>F. pertusa</i> complexes. <i>Phytotaxa</i> , 2021, 514, 149-157.	0.3	0
9	A Taxonomic Revision of the Genus <i>Poeppigia</i> (Fabaceae: Dialioideae). <i>Phytotaxa</i> , 2021, 513, 175-202.	0.3	0
10	A sophisticated case of division of labour in the trimorphic stamens of the <i>Cassia fistula</i> (Leguminosae) flower. <i>AoB PLANTS</i> , 2021, 13, plab054.	2.3	9
11	Molecular phylogenetic insights into the evolution of <i>Eriosema</i> (Fabaceae): a recent tropical savanna-adapted genus. <i>Botanical Journal of the Linnean Society</i> , 2020, 194, 439-459.	1.6	3
12	<i>Peltogyne barbata</i> (Leguminosae, Detarioideae), a new species endemic to the Trombetas River area, Brazil. <i>Kew Bulletin</i> , 2020, 75, 1.	0.9	0
13	Development of inflorescences and flowers in Fabaceae subfamily Dialioideae: an evolutionary overview and complete ontogenetic series for <i>Apuleia</i> and <i>Martiodendron</i> . <i>Botanical Journal of the Linnean Society</i> , 2020, 193, 19-46.	1.6	5
14	Phylogenetic implications of the anatomical study of the Amburaneae clade (Fabaceae: Faboideae). <i>Botanical Journal of the Linnean Society</i> , 2020, 194, 69-83.	1.6	7
15	<strong>Nomenclatural revision of the <em>Ficus</em> sect. <em>Americanae</em> (Moraceae): Typification of <em>F. citrifolia</em> and allied species</strong>. <i>Phytotaxa</i> , 2020, 474, 145-153.	0.3	1
16	Environmental filters structure plant communities in the Brazilian Chaco. <i>Acta Botanica Brasilica</i> , 2020, 34, 746-754.	0.8	3
17	<p><strong><em>Dialium</em></strong><strong> <em>heterophyllum</em> (Fabaceae: Dialioideae), a new tree species from the Amazon</strong></p>. <i>Phytotaxa</i> , 2020, 477, 47-59.	0.3	2
18	&lt;p&gt;&lt;strong&gt;<em>Parkinsonia glauca</em>&lt;/strong&gt;&lt;strong&gt; <em>Caesalpinioideae</em>, Leguminosae), a new combination and status&lt;/strong&gt;&lt;/p&gt;. <i>Phytotaxa</i> , 2020, 435, 248-250.	0.3	1

#	ARTICLE	IF	CITATIONS
19	Karyological traits related to phylogenetic signal and environmental conditions within the Hymenaea clade (Leguminosae, Detarioideae). <i>Perspectives in Plant Ecology, Evolution and Systematics</i> , 2019, 39, 125462.	2.7	10
20	<p><strong>Taxonomic Synopsis of <em>Eriosema</em> (Leguminosae: Papilionoideae, Phaseoleae) in Brazil</strong></p>. <i>Phytotaxa</i> , 2019, 416, 91-137.	0.3	5
21	Floral development of <i>Hymenaea verrucosa</i> : an ontogenetic approach to the unusual flower of Fabaceae subfamily Detarioideae. <i>Botanical Journal of the Linnean Society</i> , 2018, 187, 46-58.	1.6	14
22	Floral development of Moraceae species with emphasis on the perianth and androecium. <i>Flora: Morphology, Distribution, Functional Ecology of Plants</i> , 2018, 240, 116-132.	1.2	16
23	A revision of the neotropical <i>Mucuna</i> species (Leguminosaeâ€”Papilionoideae). <i>Phytotaxa</i> , 2018, 337, 1.	0.3	3
24	The role of biogeographical barriers and bridges in determining divergent lineages in <i>Ficus</i> (Moraceae). <i>Botanical Journal of the Linnean Society</i> , 2018, 187, 594-613.	1.6	15
25	Nomenclatural revision of <i>Ficus</i> sect. <i>Americanae</i> (Moraceae): typification of <i>Ficus americana</i> and allied species. <i>Phytotaxa</i> , 2018, 361, 244.	0.3	3
26	A new combination in <i>Parkinsonia</i> (Caesalpinoideae/Fabaceae): <i>Parkinsonia andicola</i> . <i>Phytotaxa</i> , 2018, 344, 295.	0.3	1
27	A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny: The Legume Phylogeny Working Group (LPWG). <i>Taxon</i> , 2017, 66, 44-77.	0.7	803
28	<i>Ficus latipedunculata</i> (Moraceae), a New Species from Brazil, and Taxonomic Key for <i>Ficus</i> sect. <i>Pharmacosycea</i> Occurring in Atlantic Forest. <i>Systematic Botany</i> , 2017, 42, 185-190.	0.5	1
29	Genetic conservation of small populations of the endemic tree <i>Swartzia glazioviana</i> (Taub.) Glaz. (Leguminosae) in the Atlantic Forest. <i>Conservation Genetics</i> , 2017, 18, 1105-1117.	1.5	12
30	On the â€œCangaÃ§Ã£oâ€•route: a new species of <i>Hymenaea</i> (Leguminosae) from the Brazilian Caatinga. <i>Kew Bulletin</i> , 2017, 72, 1.	0.9	5
31	Using legumes as indicators in the seasonally dry vegetation types in South America. <i>Ecological Indicators</i> , 2017, 73, 708-715.	6.3	4
32	Typification of names in <i>Ficus</i> sect. <i>Pharmacosycea</i> (Moraceae). <i>Phytotaxa</i> , 2017, 312, 298.	0.3	0
33	A taxonomic revision of the South American genus <i>Discolobium</i> (Leguminosae, Papilionoideae). <i>Phytotaxa</i> , 2017, 308, 1.	0.3	4
34	Secretory spaces in species of the clade Dipterygeae (Leguminosae, Papilionoideae). <i>Acta Botanica Brasilica</i> , 2017, 31, 374-381.	0.8	14
35	REVISITING THE TAXONOMIC DIVERSITY OF GUIBOURTIA IN THE NEOTROPICS (LEGUMINOSAE) Tj ETQq1 1 0.784314 rgBT /Overlock 10.3	0.3	1
36	Taxonomic synopsis of the <i>Ficus</i> sect. <i>Pharmacosycea</i> (Moraceae) from Colombia. <i>Phytotaxa</i> , 2017, 313, 1.	0.3	3

#	ARTICLE	IF	CITATIONS
37	DNA microsatellite markers for <i>Swartzia glazioviana</i> (Fabaceae), a threatened species from the Brazilian Atlantic Forest. <i>Applications in Plant Sciences</i> , 2016, 4, 1500081.	2.1	1
38	Hijmania, a replacement name for Maria (Moraceae). <i>Phytotaxa</i> , 2016, 247, 97.	0.3	3
39	<i>Swartzia hilareana</i> (Leguminosae), an "old" new species from the state of Minas Gerais, Brazil. <i>Phytotaxa</i> , 2016, 253, 156.	0.3	0
40	Evidence for Division of Labor and Division of Function Related to the Pollen Release in Papilionoideae (Leguminosae) with a Heteromorphic Androecium. <i>International Journal of Plant Sciences</i> , 2016, 177, 590-607.	1.3	17
41	High developmental lability in the perianth of <i>Inga</i> (Fabales, Fabaceae): a Neotropical woody rosid with gamopetalous corolla. <i>Botanical Journal of the Linnean Society</i> , 2016, , .	1.6	3
42	(25) Proposal to add Glaziou's "Plantae Brasiliae centralis a Glaziou lectae" to the list of suppressed works in Appendix VI. <i>Taxon</i> , 2016, 65, 1181-1182.	0.7	4
43	A New Species of <i>Eriosema</i> (Leguminosae, Papilionoideae, Phaseoleae) from Mato Grosso do Sul, Brazil, with a Secretory Structure Novel to the Genus. <i>Phytotaxa</i> , 2016, 263, 122.	0.3	8
44	A Taxonomic Revision of the genus <i>Dialium</i> (Leguminosae: Dialiinae) in the Netotropics. <i>Phytotaxa</i> , 2016, 283, 123.	0.3	2
45	A Molecular Phylogeny and New Infrageneric Classification of <i>Mucuna</i> Adans. (Leguminosae-Papilionoideae) including Insights from Morphology and Hypotheses about Biogeography. <i>International Journal of Plant Sciences</i> , 2016, 177, 76-89.	1.3	20
46	Evidence for a conserved karyotype in <i>Swartzia</i> (Fabaceae, Papilionoideae): Implications for the taxonomy and evolutionary diversification of a species-rich neotropical tree genus. <i>Brittonia</i> , 2016, 68, 93-101.	0.2	4
47	A TAXONOMIC REVIEW AND A NEW SPECIES OF THE SOUTH AMERICAN WOODY GENUS <i>AMBURANA</i> (LEGUMINOSAE, PAPILIONOIDEAE). <i>Phytotaxa</i> , 2015, 212, 249.	0.3	9
48	A checklist of woody Leguminosae in the South American Corridor of Dry Vegetation. <i>Phytotaxa</i> , 2015, 207, 1.	0.3	13
49	Reestablishment of <i>Hymenaea travassii</i> (Leguminosae, Caesalpinioideae), a species endemic to the Bolivian Chaco. <i>Phytotaxa</i> , 2015, 219, 96.	0.3	4
50	Growing knowledge: an overview of Seed Plant diversity in Brazil. <i>Rodriguesia</i> , 2015, 66, 1085-1113.	0.9	1,082
51	A revision of the genus <i>Myroxylon</i> (Leguminosae: Papilionoideae). <i>Kew Bulletin</i> , 2015, 70, 1.	0.9	7
52	Floral Development of the Early-Branching Papilionoid Legume <i>Amburana cearensis</i> Reveals Rare and Novel Characters. <i>International Journal of Plant Sciences</i> , 2015, 176, 94-106.	1.3	24
53	An overview of the infrageneric nomenclature of <i>Ficus</i> (Moraceae). <i>Taxon</i> , 2015, 64, 589-594.	0.7	15
54	Miscellaneous additions to <i>Swartzia</i> (Fabaceae) from Chocoan and Andean Colombia. <i>Brittonia</i> , 2015, 67, 298-310.	0.2	1

#	ARTICLE	IF	CITATIONS
55	Molecular Phylogenetics of &lt;i&gt; <i>Ficus</i> &lt;/i&gt; Section &lt;i&gt; <i>Pharmacosycea</i> &lt;/i&gt; and the Description of &lt;i&gt; <i>Ficus</i> &lt;/i&gt; Subsection &lt;i&gt; <i>Carautaea</i> &lt;/i&gt; (Moraceae). Systematic Botany, 2015, 40, 504-509.	0.5	11
56	Rodrigu��sia: 80 years disseminating Botanical Science. Rodriguesia, 2015, 66, 1-3.	0.9	13
57	Taxonomic Studies in &lt;i&gt; <i>Mucuna</i> &lt;/i&gt; Adans. (Leguminosae - Papilionoideae) from Peru. Systematic Botany, 2014, 39, 884-896.	0.5	3
58	Comparative development of rare cases of a polycarpellate gynoecium in an otherwise monocarpellate family, Leguminosae. American Journal of Botany, 2014, 101, 572-586.	1.7	26
59	Floral ontogeny in Dipterygeae (Fabaceae) reveals new insights into one of the earliest branching tribes in papilionoid legumes. Botanical Journal of the Linnean Society, 2014, 174, 529-550.	1.6	33
60	(2283) Proposal to reject the name <i>Dolichos altissimus</i> (<i>Leguminosae</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542 Td (<i>P	0.7	
61	Morphological study of fruits, seeds and embryo in the tropical tribe Dipterygeae (Leguminosae-Papilionoideae). Rodriguesia, 2014, 65, 89-97.	0.9	9
62	Increments to the genus <i>Swartzia</i> (Leguminosae) from the southern Amazonian Craton. Kew Bulletin, 2013, 68, 269-284.	0.9	4
63	Towards a new classification system for legumes: Progress report from the 6th International Legume Conference. South African Journal of Botany, 2013, 89, 3-9.	2.5	51
64	Elucidating the unusual floral features of <i>Swartzia dipetala</i> (Fabaceae). Botanical Journal of the Linnean Society, 2013, 173, 303-320.	1.6	33
65	<i>Mucuna globulifera</i> (Leguminosae: Papilionoideae), a new species from Costa Rica, Panama and Colombia. Kew Bulletin, 2013, 68, 151-155.	0.9	7
66	Three new species of <i>Mucuna</i> (Leguminosae: Papilionoideae: Phaseoleae) from South America. Kew Bulletin, 2013, 68, 143-150.	0.9	6
67	A new species of <i>Casearia</i> (Salicaceae) from Brazil. Journal of Systematics and Evolution, 2013, 51, 228-229.	3.1	10
68	A Taxonomic Revision of <i>Mucuna</i> (Fabaceae: Papilionoideae: Phaseoleae) in Brazil. Systematic Botany, 2013, 38, 631-637.	0.5	7
69	<p class="Default">Dorstenia acangatara (Moraceae), a new and threatened species from Southeastern Brazil. Phytotaxa, 2013, 118, 29.	0.3	2
70	&lt;i&gt; <i>Mucuna jarocha</i> &lt;/i&gt; (Leguminosae-Papilionoideae-Phaseoleae), a new species from Mexico. Phytotaxa, 2013, 89, 43.	0.3	6
71	Ericaceae do Parque Nacional do Itatiaia, RJ, Brasil. Hoehnea (revista), 2013, 40, 115-130.	0.2	6
72	Taxonomic Revision of the <i>Casearia ulmifolia</i> Complex (Salicaceae). Novon, 2012, 22, 196-206.	0.3	6

#	ARTICLE	IF	CITATIONS
73	Updates to the taxonomy of <i>Swartzia</i> (Leguminosae) in extra-Amazonian Brazil, with descriptions of five new species and a regional key to the genus. <i>Brittonia</i> , 2012, 64, 119-138.	0.2	14
74	Coexistence and geographical distribution of Leguminosae in an area of Atlantic forest in the semiárid region of Brazil. <i>Journal of Systematics and Evolution</i> , 2012, 50, 25-35.	3.1	10
75	Richness and diversity of Leguminosae in an altitudinal gradient in the tropical semiárid zone of Brazil. <i>Journal of Systematics and Evolution</i> , 2012, 50, 433-442.	3.1	2
76	A família Leguminosae na Serra de Baturité, Ceará, uma Área de Floresta Atlântica no semiárido brasileiro. <i>Rodriguesia</i> , 2011, 62, 563-613.	0.9	13
77	A taxonomic reappraisal of the South American genus <i>Holocalyx</i> (Leguminosae, Papilionoideae). <i>Brittonia</i> , 2010, 62, 110-115.	0.2	3
78	Comparações florísticas e taxonomia da família Gesneriaceae no Parque Nacional do Itatiaia, Brasil. <i>Hoehnea</i> (revista), 2010, 37, 131-145.	0.2	6
79	A New Species of <i>Casearia</i> (Salicaceae) from Southeastern Brazil. <i>Novon</i> , 2010, 20, 179-181.	0.3	8
80	A phylogenetically based sectional classification of <i>Swartzia</i> (Leguminosae-Papilionoideae). <i>Taxon</i> , 2009, 58, 913-924.	0.7	15
81	Floral anatomy of the Lecointea clade (Leguminosae, Papilionoideae, Swartzieae sensu lato). <i>Plant Systematics and Evolution</i> , 2008, 273, 201-209.	0.9	12
82	O gênero <i>Swartzia</i> Schreb. (Leguminosae, Papilionoideae) no estado do Rio de Janeiro. <i>Rodriguesia</i> , 2007, 58, 469-483.	0.9	5
83	A new <i>Swartzia</i> (Leguminosae: Papilionoideae: Swartzieae) species with trimorphic stamens from Amazonian Brazil. <i>Botanical Journal of the Linnean Society</i> , 2005, 147, 235-238.	1.6	7
84	Composition of the Lecointeaclade (Leguminosae, Papilionoideae, Swartzieae), a reevaluation based on combined evidence from morphology and molecular data. <i>Taxon</i> , 2004, 53, 1007-1018.	0.7	21
85	A new species of <i>Exostyles</i> (Leguminosae, Papilionoideae, Swartzieae s.l.), from Paraná State, Brazil. <i>Botanical Journal of the Linnean Society</i> , 2004, 146, 103-106.	1.6	1
86	A Revision of the Genus <i>Exostyles</i> Schott (Leguminosae: Papilionoideae). <i>Kew Bulletin</i> , 2004, 59, 521.	0.9	4
87	Floral ontogeny of <i>Lecointea</i>, <i>Zollernia</i>, <i>Exostyles</i>, and <i>Harleyodendron</i> (Leguminosae: Papilionoideae: Swartzieae s.l.). <i>American Journal of Botany</i> , 2002, 89, 1553-1569.	1.7	51
88	<i>Swartzia</i> Schreb. (Leguminosae: Papilionoideae: Swartzieae): A Taxonomic Study of the <i>Swartzia acutifolia</i> Complex including a New Name and a New Species from Southeastern Brazil. <i>Kew Bulletin</i> , 2001, 56, 917.	0.9	16
89	Phytogeographic relationships of the species of Leguminosae presents in an area of the Atlantic forest domain in the semi-arid region of Brazil. <i>Rodriguesia</i> , 0, 72, .	0.9	0
90	Phylogeny of <i>Dorstenia</i> (Moraceae) reveals the polyphyletic nature of its neotropical sections. <i>Rodriguesia</i> , 0, 72, .	0.9	1

#	ARTICLE	IF	CITATIONS
91	Taxonomic review of the species of <i>Parkinsonia</i> (Leguminosae, Caesalpinoideae) from the Americas. <i>Rodriguesia</i> , 0, 72, .	0.9	3
92	Flora of Espírito Santo: Capparaceae. <i>Rodriguesia</i> , 0, 73, .	0.9	0
93	Deguelia tenuiflora (Leguminosae, Papilioideae), a remarkable new species from the Brazilian Amazon. <i>Rodriguesia</i> , 0, 73, .	0.9	1