Takeshi Ozeki

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/584425/publications.pdf

Version: 2024-02-01

1163117 1474206 9 702 8 9 citations h-index g-index papers 9 9 9 1124 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Genome-wide association study identifies HLA-A*3101 allele as a genetic risk factor for carbamazepine-induced cutaneous adverse drug reactions in Japanese population. Human Molecular Genetics, 2011, 20, 1034-1041.	2.9	454
2	Specific HLA types are associated with antiepileptic drug-induced Stevens–Johnson syndrome and toxic epidermal necrolysis in Japanese subjects. Pharmacogenomics, 2013, 14, 1821-1831.	1.3	60
3	Association of HLA-A*31:01 Screening With the Incidence of Carbamazepine-Induced Cutaneous Adverse Reactions in a Japanese Population. JAMA Neurology, 2018, 75, 842.	9.0	52
4	Empirical evaluation of variant calling accuracy using ultra-deep whole-genome sequencing data. Scientific Reports, 2019, 9, 1784.	3.3	46
5	Variants at HLA-A, HLA-C, and HLA-DQB1 Confer Risk of Psoriasis Vulgaris in Japanese. Journal of Investigative Dermatology, 2018, 138, 542-548.	0.7	39
6	Comparison of effects of UGT1A1*6 and UGT1A1*28 on irinotecan-induced adverse reactions in the Japanese population: analysis of the Biobank Japan Project. Journal of Human Genetics, 2019, 64, 1195-1202.	2.3	19
7	Association of HLA-A*11:01 with Sulfonamide-Related Severe Cutaneous Adverse Reactions in Japanese Patients. Journal of Investigative Dermatology, 2020, 140, 1659-1662.e6.	0.7	18
8	<i>HLAâ€B*51:01</i> and <i>CYP2C9*3</i> Are Risk Factors for Phenytoinâ€Induced Eruption in the Japanese Population: Analysis of Data From the Biobank Japan Project. Clinical Pharmacology and Therapeutics, 2020, 107, 1170-1178.	4.7	13
9	Characteristics of adverse drug reactions associated with antiepileptics at a tertiary children's hospital in Japan: A retrospective observational cohort study. Epilepsy Research, 2021, 173, 106614.	1.6	1