

Maria Pascual

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5845835/publications.pdf>

Version: 2024-02-01

16
papers

1,531
citations

687363

13
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

1850
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Expanded Adipose-Derived Stem Cells for the Treatment of Complex Perianal Fistula. <i>Diseases of the Colon and Rectum</i> , 2009, 52, 79-86. | 1.3 | 694 |
| 2 | An assessment of the incidence of fistula-in-ano in four countries of the European Union. <i>International Journal of Colorectal Disease</i> , 2007, 22, 1459-1462. | 2.2 | 190 |
| 3 | Treatment of enterocutaneous fistula in Crohn's Disease with adipose-derived stem cells: a comparison of protocols with and without cell expansion. <i>International Journal of Colorectal Disease</i> , 2009, 24, 27-30. | 2.2 | 143 |
| 4 | Involvement of Fc γ 3 receptor IIIA genotypes in susceptibility to rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2000, 43, 735. | 6.7 | 130 |
| 5 | IL-6 promoter polymorphisms in rheumatoid arthritis. <i>Genes and Immunity</i> , 2000, 1, 338-340. | 4.1 | 86 |
| 6 | The telomeric part of the HLA region predisposes to rheumatoid arthritis independently of the class II loci. <i>Human Immunology</i> , 2001, 62, 75-84. | 2.4 | 63 |
| 7 | Genetic polymorphisms in Spanish rheumatoid arthritis patients: an association and linkage study. <i>Genes and Immunity</i> , 2003, 4, 117-121. | 4.1 | 38 |
| 8 | A poly(ADP-ribose) polymerase haplotype spanning the promoter region confers susceptibility to rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2003, 48, 638-641. | 6.7 | 37 |
| 9 | A new allele within the transmembrane region of the human MICA gene with seven GCT repeats. <i>Tissue Antigens</i> , 2002, 60, 526-528. | 1.0 | 33 |
| 10 | Distribution of HERV-LTR elements in the 5' flanking region of HLA-DQB1 and association with autoimmunity. <i>Immunogenetics</i> , 2001, 53, 114-118. | 2.4 | 29 |
| 11 | Association of MICA-A5.1 allele with susceptibility to celiac disease in a family study. <i>American Journal of Gastroenterology</i> , 2003, 98, 359-362. | 0.4 | 28 |
| 12 | HLA haplotypes and susceptibility to rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2002, 31, 275-278. | 1.1 | 24 |
| 13 | Genetic determinants of rheumatoid arthritis: the inducible nitric oxide synthase (NOS2) gene promoter polymorphism. <i>Genes and Immunity</i> , 2002, 3, 299-301. | 4.1 | 24 |
| 14 | Polymorphism of the inducible nitric oxide synthase gene in celiac disease. <i>Human Immunology</i> , 2002, 63, 1062-1065. | 2.4 | 8 |
| 15 | Association of Fc γ 3 receptor IIIA polymorphism with rheumatoid arthritis: Comment on the article by Morgan et al. <i>Arthritis and Rheumatism</i> , 2002, 46, 556-557. | 6.7 | 3 |
| 16 | Distribution of HERV-LTR elements in HLA-DQB1 alleles: comments on the article by Bieda et al.. <i>Diabetologia</i> , 2003, 46, 869-870. | 6.3 | 1 |