Maria Pascual

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5845835/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Treatment of enterocutaneous fistula in Crohn's Disease with adipose-derived stem cells: a comparison of protocols with and without cell expansion. International Journal of Colorectal Disease, 2009, 24, 27-30.	2.2	143
2	Expanded Adipose-Derived Stem Cells for the Treatment of Complex Perianal Fistula. Diseases of the Colon and Rectum, 2009, 52, 79-86.	1.3	694
3	An assessment of the incidence of fistula-in-ano in four countries of the European Union. International Journal of Colorectal Disease, 2007, 22, 1459-1462.	2.2	190
4	Distribution of HERV-LTR elements in HLA-DQB1 alleles: comments on the article by Bieda et al Diabetologia, 2003, 46, 869-870.	6.3	1
5	A poly(ADPâ€ribose) polymerase haplotype spanning the promoter region confers susceptibility to rheumatoid arthritis. Arthritis and Rheumatism, 2003, 48, 638-641.	6.7	37
6	Genetic polymorphisms in Spanish rheumatoid arthritis patients: an association and linkage study. Genes and Immunity, 2003, 4, 117-121.	4.1	38
7	Association of MICA-A5.1 allele with susceptibility to celiac disease in a family study. American Journal of Gastroenterology, 2003, 98, 359-362.	0.4	28
8	HLA haplotypes and susceptibility to rheumatoid arthritis. Scandinavian Journal of Rheumatology, 2002, 31, 275-278.	1.1	24
9	Polymorphism of the inducible nitric oxide synthase gene in celiac disease. Human Immunology, 2002, 63, 1062-1065.	2.4	8
10	Association of Fc ^î 3 receptor IIIA polymorphism with rheumatoid arthritis: Comment on the article by Morgan et al. Arthritis and Rheumatism, 2002, 46, 556-557.	6.7	3
11	A new allele within the transmembrane region of the human MICA gene with seven GCT repeats. Tissue Antigens, 2002, 60, 526-528.	1.0	33
12	Genetic determinants of rheumatoid arthritis: the inducible nitric oxide synthase (NOS2) gene promoter polymorphism. Genes and Immunity, 2002, 3, 299-301.	4.1	24
13	The telomeric part of the HLA region predisposes to rheumatoid arthritis independently of the class II loci. Human Immunology, 2001, 62, 75-84.	2.4	63
14	Distribution of HERV-LTR elements in the 5′-flanking region of HLA-DQB1 and association with autoimmunity. Immunogenetics, 2001, 53, 114-118.	2.4	29
15	Involvement of FcÎ ³ receptor IIIA genotypes in susceptibility to rheumatoid arthritis. Arthritis and Rheumatism, 2000, 43, 735.	6.7	130
16	IL-6 promoter polymorphisms in rheumatoid arthritis. Genes and Immunity, 2000, 1, 338-340.	4.1	86